
Complex Analysis I:

Problem Set VIII

Youngduck Choi
CILVR Lab
New York University
yc1104@nyu.edu

Abstract

This work contains the solutions to the problem set VIII of Complex Analysis I 2015 at Courant Institute of Mathematical Sciences.

Question 273-12.

Solution. (a) As $\exp(iz^2)$ is entire, by Cauchy-Goursat theorem, we have

$$\int_{C_1} \exp(iz^2) dz + \int_{C_R} \exp(iz^2) dz + \int_{C_2} \exp(iz^2) dz = 0,$$

where C_1 refers to the segment from 0 to R , and C_2 refers to the segment from $Re^{i\frac{\pi}{4}}$ to 0. Observe that we can parametrize points on C_1 as $z = x(0 \leq x \leq R)$, and points on C_2 as $z = re^{i\frac{\pi}{4}}(0 \leq r \leq R)$. Therefore, the above equality can be written as

$$\int_0^R \exp(ix^2) dx = e^{i\frac{\pi}{4}} \int_0^R e^{-r^2} dr - \int_{C_R} e^{iz^2} dz.$$

Equating the real and imaginary parts separately from the above equation yields

$$\begin{aligned} \int_0^R \cos(x^2) dx &= \frac{1}{\sqrt{2}} \int_0^R e^{-r^2} dr - \operatorname{Re} \int_{C_R} e^{iz^2} dz, \\ \int_0^R \sin(x^2) dx &= \frac{1}{\sqrt{2}} \int_0^R e^{-r^2} dr - \operatorname{Im} \int_{C_R} e^{iz^2} dz. \end{aligned}$$

□

(b) We can parametrize the points on C_R as $z = Re^{i\theta}(0 \leq \theta \leq \frac{\pi}{4})$. It follows that

$$\begin{aligned} \int_{C_R} e^{iz^2} &= \int_0^{\frac{\pi}{4}} e^{iR^2 e^{i2\theta}} Rie^{i\theta} d\theta \\ &= iR \int_0^{\frac{\pi}{4}} e^{-R^2 \sin(2\theta)} e^{iR^2 \cos(2\theta)} e^{i\theta} d\theta. \end{aligned}$$

Consequently, we have

$$\begin{aligned}
\left| \int_{C_R} e^{iz^2} \right| &= \left| iR \int_0^{\frac{\pi}{4}} e^{-R^2 \sin(2\theta)} e^{iR^2 \cos(2\theta)} e^{i\theta} d\theta \right| \\
&\leq R \int_0^{\frac{\pi}{4}} \left| e^{-R^2 \sin(2\theta)} \right| \left| e^{iR^2 \cos(2\theta)} \right| \left| e^{i\theta} \right| d\theta \\
&= R \int_0^{\frac{\pi}{4}} \left| e^{-R^2 \sin(2\theta)} \right| d\theta \\
&= R \int_0^{\frac{\pi}{4}} e^{-R^2 \sin(2\theta)} d\theta \\
&= \frac{R}{2} \int_0^{\frac{\pi}{2}} e^{-R^2 \sin(\phi)} d\theta,
\end{aligned}$$

by change of variable $\phi = 2\theta$. From Jordan's lemma in section 88, pg. 271, continues to

$$\begin{aligned}
\left| \int_{C_R} e^{iz^2} \right| &\leq \frac{R}{2} \frac{\pi}{2R^2} \\
&= \frac{\pi}{4R},
\end{aligned}$$

which limits 0 as $R \rightarrow \infty$. Hence, the integral limits 0 as $R \rightarrow \infty$. □

(c) Combining the result from (a),(b) and the given formula, we have

$$\begin{aligned}
\int_0^\infty \cos(x^2) dx &= \frac{1}{\sqrt{2}} \int_0^\infty e^{-x^2} dx \\
&= \frac{1}{2} \sqrt{\frac{\pi}{2}}, \\
\int_0^\infty \sin(x^2) dx &= \frac{1}{\sqrt{2}} \int_0^\infty e^{-x^2} dx \\
&= \frac{1}{2} \sqrt{\frac{\pi}{2}},
\end{aligned}$$

as desired. □

Question 278-1.

Solution. Let $f(z) = \frac{e^{iaz} - e^{ibz}}{z^2}$. On the indented contour in figure 108, by Cauchy-Goursat theorem, we obtain

$$\int_{L_1} f(z) dz + \int_{L_2} f(z) dz = - \int_{C_\rho} f(z) dz - \int_{C_R} f(z) dz.$$

Observe that on L_1 and L_2 , we have the following parametric representations:

$$L_1 : z = re^{i0} = r(\rho \leq r \leq R) \text{ and } -L_2 : z = re^{i\pi} = -r(\rho \leq r \leq R).$$

It follows that

$$\begin{aligned}
\int_{L_1} f(z) dz + \int_{L_2} f(z) dz &= \int_{L_1} f(z) dz - \int_{-L_2} f(z) dz \\
&= \int_\rho^R \frac{e^{iar} - e^{ibr}}{r^2} dr + \int_\rho^R \frac{e^{-iar} - e^{-ibr}}{r^2} dr \\
&= \int_\rho^R \frac{(e^{iar} + e^{-iar}) - (e^{ibr} + e^{-ibr})}{r^2} dr \\
&= 2 \int_\rho^R \frac{\cos(ar) - \cos(br)}{r^2} dr.
\end{aligned}$$

Now, writing the Laurent series of $f(z)$ gives

$$\begin{aligned} f(z) &= \frac{e^{iaz} - e^{ibz}}{z^2} \\ &= \frac{1}{z^2} \left[\left(1 + \frac{iaz}{1!} + \frac{(iaz)^2}{2!} \dots \right) + \left(1 + \frac{ibz}{1!} + \frac{(ibz)^2}{2!} \dots \right) \right] \\ &= \frac{i(a-b)}{z} + \dots \end{aligned}$$

Therefore, we see that $z = 0$ is a simple pole of $f(z)$ and the residue B_0 equals $i(a-b)$. Consequently, we obtain that

$$\lim_{\rho \rightarrow 0} \int_{C_\rho} f(z) dz = -i(a-b)\pi i = \pi(a-b).$$

Observe that

$$|f(z)| \leq \frac{|e^{iaz}| + |e^{ibz}|}{|z|^2}.$$

For z on C_R , we have

$$|f(z)| \leq \frac{2}{R^2}.$$

Therefore, by ML inequality, it follows that

$$\begin{aligned} \left| \int_{C_R} f(z) dz \right| &\leq \int_{C_R} |f(z)| dz \\ &\leq \frac{2}{R^2} \pi R \\ &= \frac{2\pi}{R}. \end{aligned}$$

Therefore, as $R \rightarrow \infty$ $\frac{2\pi}{R} \rightarrow 0$, the integral $\int_{C_R} f(z) dz$ tends to 0 as $R \rightarrow \infty$. Consequently,

$$\int_0^\infty \frac{\cos(ar) - \cos(br)}{r^2} dr = \frac{\pi}{2}(b-a).$$

Setting $b = 2$ and $a = 0$ and using the identity $1 - \cos(2x) = 2\sin^2(x)$, we finally obtain

$$\int_0^\infty \frac{\sin^2(x)}{x^2} dx = \frac{\pi}{2},$$

as desired. \square

Question 287-1.

Solution. Observe that the following equality holds, by the linearity of integration:

$$\int_0^{2\pi} \frac{d\theta}{5 + 4 \sin \theta} = \frac{1}{5} \int_0^{2\pi} \frac{d\theta}{1 + \frac{4}{5} \sin \theta}$$

From the example 1 from pg.285 in the section 92, it follows that

$$\begin{aligned} \int_0^{2\pi} \frac{d\theta}{5 + 4 \sin \theta} &= \frac{1}{5} \int_0^{2\pi} \frac{d\theta}{1 + \frac{4}{5} \sin \theta} \\ &= \frac{1}{5} \frac{2\pi}{\sqrt{1 - \frac{4}{5}^2}} \\ &= \frac{1}{5} \frac{2\pi}{\frac{3}{5}} \\ &= \frac{2}{3}\pi. \end{aligned}$$

\square

Question 287-6.

Solution. As $|\sin(-\pi + \theta)| = |\sin(\theta)|$, we have $\sin^{2n}(\theta) = \sin^{2n}(-\pi + \theta)$, and $\int_0^\pi \sin^{2n}(\theta) d\theta = \int_{-\pi}^0 \sin^{2n}(\theta) d\theta$. Hence, it follows that

$$\int_0^\pi \sin^{2n}(\theta) d\theta = \frac{1}{2} \int_C \sin^{2n}(\theta),$$

where C is the positively oriented unit circle $|z| = 1$. Substituting $\frac{z - z^{-1}}{2i}$ for $\sin(\theta)$, using the binomial formula, and linearity of integration, we obtain

$$\begin{aligned} \frac{1}{2} \int_C \sin^{2n}(\theta) &= \frac{1}{2} \int_C \left(\frac{z - z^{-1}}{2i} \right)^{2n} \frac{dz}{iz} \\ &= \frac{1}{2^{2n+1}(-1)^ni} \int_C \frac{(z - z^{-1})^{2n}}{z} dz \\ &= \frac{1}{2^{2n+1}(-1)^ni} \int_C \sum_{k=0}^n \binom{2n}{k} z^k z^{2n-k} (-z^{-1})^k z^{-1} dz \\ &= \frac{1}{2^{2n+1}(-1)^ni} \sum_{k=0}^n \binom{2n}{k} (-1)^k \int_C z^{2n-2k-1} dz. \end{aligned}$$

Observe that we only get non-zero integral value for $k = n$ case, and $\int_C z^{-1} dz = 2\pi i$. Therefore, it follows that

$$\begin{aligned} \int_0^\pi \sin^{2n}(\theta) d\theta &= \frac{1}{2^{2n+1}(-1)^ni} \frac{(2n)!}{n!n!} (-1)^n 2\pi i \\ &= \frac{2n!}{2^{2n}(n!)^2} \pi, \end{aligned}$$

as desired. □

Question 293-6.

Solution. (a) Inside the circle $|z| = 1$, write

$$f(z) = -5z^4 \text{ and } g(z) = z^6 + z^3 - 2z.$$

Then, observe that when $|z| = 1$,

$$|f(z)| = 5|z|^4 = 5 \text{ and } |g(z)| \leq |z|^6 + |z|^3 + 2|z| = 4.$$

The conditions of Rouché's theorem are thus satisfied. Consequently, since $f(z)$ has 4 zeroes, counting multiplicities, inside the circle $|z| = 1$, $f(z) + g(z)$ has 4 zeroes, inside the circle $|z| = 1$. Therefore, the polynomial $z^6 - 5z^4 + z^3 - 2z$ has 4 zeroes, inside the circle $|z| = 1$. □

(b) Inside the circle $|z| = 1$, write

$$f(z) = 9 \text{ and } g(z) = 2z^4 - 2z^3 + 2z^2 - 2z.$$

Then, observe that when $|z| = 1$,

$$|f(z)| = 9 \text{ and } |g(z)| = 2|z|^4 + 2|z|^3 + 2|z|^2 + 2|z| = 8.$$

The conditions of Rouché's theorem are thus satisfied. Consequently, since $f(z)$ has 0 zero, counting multiplicities, inside the circle $|z| = 1$, $f(z) + g(z)$ has 0 zero, inside the circle $|z| = 1$. Therefore, the polynomial $2z^4 - 2z^3 + 2z^2 - 2z + 9$ has 0 zero, inside the circle $|z| = 1$. □

(c) Inside the circle $|z| = 1$, write

$$f(z) = -4z^3 \text{ and } g(z) = z^7 + z - 1.$$

Then, observe that when $|z| = 1$,

$$|f(z)| = 4|z|^3 = 4 \text{ and } |g(z)| \leq |z|^7 + |z| - 1 = 1.$$

The conditions of Rouché's theorem are thus satisfied. Consequently, since $f(z)$ has 3 zeroes, counting multiplicities, inside the circle $|z| = 1$, $f(z) + g(z)$ has 3 zeroes. Therefore, the polynomial $z^7 - 4z^3 + z - 1$ has 3 zeroes inside the circle $|z| = 1$. □

Question 293-8.

Solution. Inside the circle $|z| = 2$, write

$$f(z) = 2z^5 \text{ and } g(z) = 6z^2 + z + 1.$$

Then, observe that when $|z| = 2$,

$$|f(z)| = 2|z|^5 = 64 \text{ and } |g(z)| \leq 6|z|^2 + |z| + |1| = 8.$$

The conditions of Rouché's theorem are thus satisfied. Consequently, since $f(z)$ has 5 zeroes, counting multiplicities, inside the circle $|z| = 2$, $f(z) + g(z)$ has 5 zeroes. On the other hand, inside the circle $|z| = 1$, write

$$f(z) = -6z^2 \text{ and } g(z) = 2z^5 + z + 1.$$

Then, observe that when $|z| = 1$,

$$|f(z)| = 6|z|^2 = 6 \text{ and } |g(z)| \leq 2|z|^5 + |z| + 1 = 4.$$

The conditions of Rouché's theorem are thus satisfied. Consequently, since $f(z)$ has 2 zeroes, counting multiplicities, inside the circle $|z| = 1$, $f(z) + g(z)$ has 2 zeroes. Therefore, we have shown that in the annulus $1 \leq |z| \leq 2$, we have $5 - 2 = 3$ zeroes. \square