

---

# Real Variables: Problem Set X

---

**Youngduck Choi**  
Courant Institute of Mathematical Sciences  
New York University  
yc1104@nyu.edu

## Abstract

This work contains solutions to the problem set X of Real Variables 2015 at NYU.

## 1 Solutions

**Question Royden 17-6.**

6. Let  $(X, \mathcal{M}, \mu)$  be a measure space and  $X_0$  belong to  $\mathcal{M}$ . Define  $\mathcal{M}_0$  to be the collection of sets in  $\mathcal{M}$  that are subsets of  $X_0$  and  $\mu_0$  the restriction of  $\mu$  to  $\mathcal{M}_0$ . Show that  $(X_0, \mathcal{M}_0, \mu_0)$  is a measure space.

**Solution.** We first show that  $\| \cdot \| : X \rightarrow \mathbb{R}$  given is a norm on  $X$ . First of all,

**Question Royden 17-15.**

15. Show that if  $\nu_1$  and  $\nu_2$  are any two finite signed measures, then so is  $\alpha\nu_1 + \beta\nu_2$ , where  $\alpha$  and  $\beta$  are real numbers. Show that

$$|\alpha\nu| = |\alpha||\nu| \text{ and } |\nu_1 + \nu_2| \leq |\nu_1| + |\nu_2|,$$

where  $\nu \leq \mu$  means  $\nu(E) \leq \mu(E)$  for all measurable sets  $E$ .

**Solution.**

**Question Royden 17-17.**

17. Let  $\mu$  and  $\nu$  be finite signed measures. Define  $\mu \wedge \nu = \frac{1}{2}(\mu + \nu - |\mu - \nu|)$  and  $\mu \vee \nu = \mu + \nu - \mu \wedge \nu$ .
- (i) Show that the signed measure  $\mu \wedge \nu$  is smaller than  $\mu$  and  $\nu$  but larger than any other signed measure that is smaller than  $\mu$  and  $\nu$ .
  - (ii) Show that the signed measure  $\mu \vee \nu$  is larger than  $\mu$  and  $\nu$  but smaller than any other measure that is larger than  $\mu$  and  $\nu$ .
  - (iii) If  $\mu$  and  $\nu$  are positive measures, show that they are mutually singular if and only if  $\mu \wedge \nu = 0$ .

**Solution.**

**Question Royden 18-50.**

50. Establish the uniqueness of the function  $f$  in the Radon-Nikodym Theorem.

**Solution.**

**Question Royden 18-54.**

54. Let  $\mu$ ,  $\nu$ , and  $\lambda$  be  $\sigma$ -finite measures on the measurable space  $(X, \mathcal{M})$ .

- (i) If  $\nu \ll \mu$  and  $f$  is a nonnegative function on  $X$  that is measurable with respect to  $\mathcal{M}$ , show that

$$\int_X f \, d\nu = \int_X f \left[ \frac{d\nu}{d\mu} \right] d\mu.$$

- (ii) If  $\nu \ll \mu$  and  $\lambda \ll \mu$ , show that

$$\frac{d(\nu + \lambda)}{d\mu} = \frac{d\nu}{d\mu} + \frac{d\lambda}{d\mu} \text{ a.e. } [\mu].$$

- (iii) If  $\nu \ll \mu \ll \lambda$ , show that

$$\frac{d\nu}{d\lambda} = \frac{d\nu}{d\mu} \cdot \frac{d\mu}{d\lambda} \text{ a.e. } [\lambda].$$

**Solution.**

**Question Royden 18-55.**

55. Let  $\mu$ ,  $\nu$ ,  $\nu_1$ , and  $\nu_2$  be measures on the measurable space  $(X, \mathcal{M})$ .

- (i) Show that if  $\nu \perp \mu$  and  $\nu \ll \mu$ , then  $\nu = 0$ .
- (ii) Show that if  $\nu_1$  and  $\nu_2$  are singular with respect to  $\mu$ , then, for any  $\alpha \geq 0, \beta \geq 0$ , so is the measure  $\alpha\nu_1 + \beta\nu_2$ .
- (iii) Show that if  $\nu_1$  and  $\nu_2$  are absolutely continuous with respect to  $\mu$ , then, for any  $\alpha \geq 0, \beta \geq 0$ , so is the measure  $\alpha\nu_1 + \beta\nu_2$ .
- (iv) Prove the uniqueness assertion in the Lebesgue decomposition.

**Solution.**