

---

# Real Variables: Problem Set VI

---

Youngduck Choi  
Courant Institute of Mathematical Sciences  
New York University  
yc1104@nyu.edu

## Abstract

This work contains solutions to the problem set VI of Real Variables 2015 at NYU.

## 1 Solutions

### Question 9.10.

**Solution.** Let  $\{X_n, \rho_n\}_{n=1}^{\infty}$  be a countable collection of metric spaces. We now define  $(\prod_{n=1}^{\infty} X_n, p_*) = (X, p_*)$  such that for  $x, y \in X$ ,

$$p_*(x, y) = \sum_{n=1}^{\infty} \frac{1}{2^n} \cdot \frac{p_n(x_n, y_n)}{1 + p_n(x_n, y_n)}.$$

First, we can show that  $p_*$  is well-defined via comparison test with the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ , as

$$0 \leq \frac{p_n(x_n, y_n)}{1 + p_n(x_n, y_n)} \leq 1 \text{ for all } n.$$

As  $p_n(x_n, y_n) \geq 0$  for all  $n$ , we have  $p_*(x, y) \geq 0$  for all  $x, y \in X$ . If  $p_*(x, y) = 0$ , then  $p_n(x_n, y_n) = 0$  for all  $n$ . As each  $p_n$  is a metric space  $x_n = y_n$  for all  $n$ . Therefore,  $x = y$ . If  $x = y$ , then  $x_n = y_n$  for all  $n$ . As each  $p_n$  is a metric space,  $p_n(x_n, y_n) = 0$  for all  $n$ . Therefore,  $p_*(x, y) = 0$ .

Since  $p_n(x_n, y_n) = p_n(y_n, x_n)$  for all  $n$ , for  $x, y \in X$ , we

$$\begin{aligned} p_*(x, y) &= \sum_{n=1}^{\infty} \frac{1}{2^n} \cdot \frac{p_n(x_n, y_n)}{1 + p_n(x_n, y_n)} \\ &= \sum_{n=1}^{\infty} \frac{1}{2^n} \cdot \frac{p_n(y_n, x_n)}{1 + p_n(y_n, x_n)} \\ &= p_*(y, x). \end{aligned}$$

Let  $x, y, z \in X$ . By the problem 6 and the triangle inequality of each metric space  $X_n$ , which gives  $p_n(x_n, z_n) \leq p_n(x_n, y_n) + p_n(y_n, z_n)$  for each  $n$ , we have

$$\frac{p_n(x_n, z_n)}{1 + p_n(x_n, z_n)} \leq \frac{p_n(x_n, y_n)}{1 + p_n(x_n, y_n)} + \frac{p_n(y_n, z_n)}{1 + p_n(y_n, z_n)},$$

for all  $n$ . Hence, we have

$$\sum_{n=1}^{\infty} \frac{p_n(x_n, z_n)}{1 + p_n(x_n, z_n)} \leq \sum_{n=1}^{\infty} \frac{p_n(x_n, y_n)}{1 + p_n(x_n, y_n)} + \frac{p_n(y_n, z_n)}{1 + p_n(y_n, z_n)},$$

which can be written as

$$p_*(x, z) \leq p_*(x, y) + p_*(y, z).$$

Therefore, we have shown that all required properties of a metric space hold for  $(X, p_*)$ .  $(X, p_*)$  is a metric space.  $\square$

### Question 9.20.

**Solution.** Let  $E$  be a subset of a metric space  $X$ , and let  $\text{int}E$  be the interior of  $E$ . We first show that  $\text{int}E \subseteq E$ . If  $x \in X \setminus E$ , then every ball of  $x$  contains a point in  $X \setminus E$ . Hence,  $x \notin E$ . Therefore,  $\text{int}E \subseteq E$ .

Now, we wish to show that  $\text{int}E$  is open. For the first case, assume that  $E = \text{int}E$ . Let  $x \in \text{int}E$ . Since  $x$  is an interior point of  $E$ , there exists an open ball  $B(x, r)$  contained in  $E$ . Since  $E = \text{int}E$ , the open ball  $B(x, r)$  is contained in  $\text{int}E$  as well. Hence,  $\text{int}E$  is open in this case. For the remaining case, assume that  $E \setminus \text{int}E \neq \emptyset$ . Let  $x \in \text{int}E$ . Since  $x$  is an interior point of  $E$ , there exists an open ball  $B(x, r)$  contained in  $E$ . Suppose that there exists  $y \in B(x, r) \cap E \setminus \text{int}E$ . Then, we have  $d(x, y) < r$ . Consider  $B(y, r - d(x, y))$ , which is valid since  $r - d(x, y) > 0$ . By the triangle inequality, for any point  $z \in B(y, r - d(x, y))$ ,

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< r. \end{aligned}$$

Hence,  $B(y, r - d(x, y))$  is an open ball contained in  $B(x, r)$ , which is again contained in  $E$ , which contradicts the fact that  $y \in E \setminus \text{int}E$ . Hence,  $B(x, r)$  is contained in  $\text{int}E$ . Therefore,  $\text{int}E$  is open. As we covered all cases,  $\text{int}E$  for any subset  $E$  of a metric space  $X$  is open.

Assume  $E$  is open. Let  $x \in E$ . As  $E$  is open, there exists an open ball around  $x$  contained in  $E$ . Therefore,  $x \in \text{int}E$ . Hence,  $E \subseteq \text{int}E$ . As we have  $\text{int}E \subseteq E$  from above, we have shown that  $E = \text{int}E$ .

Assume  $E = \text{int}E$ . Let  $x \in E$ . Then, as  $E = \text{int}E$ ,  $x \in \text{int}E$ . By the definition of interior point, there exists an open ball around  $x$  contained in  $E$ . Hence,  $E$  is open.  $\square$

### Question 9.32.

**Solution.**

,

### Question 9.43.

**Solution.**

### Question 9.72.

**Solution.** Assume  $A \cap B \neq \emptyset$ . Then, there exists  $x \in A \cap B$ . Since  $\rho(x, x) = 0$ , we have  $\text{dist}(A, B) = 0$ . By contrapositive, we have shown that if  $\text{dist}(A, B) > 0$ , then  $A \cap B = \emptyset$ . Assume that  $A \cap B = \emptyset$ .

### Question 9.77.

**Solution.** Let  $X$  and  $Y$  be separable metric spaces. Consider the standard product metric on  $X \times Y$ . Then, there exist a countable dense subset  $D_X$  in  $X$  and countable dense subset  $D_Y$  in  $Y$ . Observe that  $D_X \times D_Y$  is countable. We claim that  $D_X \times D_Y$  is a dense subset in  $X \times Y$ . Therefore,  $X \times Y$  is separable.  $\square$