
Real Variables: Problem Set V

Youngduck Choi
Courant Institute of Mathematical Sciences
New York University
yc1104@nyu.edu

Abstract

This work contains solutions to the problem set V of Real Variables 2015 at NYU.

1 Solutions

Question 6.10.

Solution. Let $x_1, x_2 \in (a, b)$, such that $x_1 < x_2$. Then, we have

$$\begin{aligned} f(x_1) &= \sum_{k=1}^{\infty} l((c_k, d_k) \cap (-\infty, x_1)) \\ f(x_2) &= \sum_{k=1}^{\infty} l((c_k, d_k) \cap (-\infty, x_2)). \end{aligned}$$

As $x_1 < x_2$, by the monotonicity property of the length function, we have

$$l((c_k, d_k) \cap (-\infty, x_1)) \leq l((c_k, d_k) \cap (-\infty, x_2)),$$

for all $k \geq 1$. It follows that $f(x_1) \leq f(x_2)$. Hence, f is increasing. We show that f fails to be differentiable at each point in E , which is a set of measure zero contained in the open interval (a, b) . Let $x \in E$. Then, by the preceding problem, there exist a countable collection of open intervals contained in (a, b) , $\{(c_k, d_k)\}_{k=1}^{\infty}$ such that each point in E belongs to infinitely many intervals in the collection and $\sum_{k=1}^{\infty} d_k - c_k < \infty$. Let $\{(c_{k_i}, d_{k_i})\}_{i=1}^{\infty}$ be the sub-collection such that $x \in (c_{k_i}, d_{k_i})$ for all i . Then, there exist a finite sub-cover $\{(c_{k_i}, d_{k_i})\}_{i=1}^n$ that x belongs to. Since, n is finite, as intersection of finite open sets is open, there exists a_n such that

$$B(x, a_n) \in \cup_{k=1}^n (c_{k_i}, d_{k_i}),$$

such that (B, a_n) denotes the ball of radius a_n , centered at x . Observe that

$$\begin{aligned} f(x + a_n) - f(x) &\geq \sum_{i=1}^n l((c_{k_i}, d_{k_i}) \cap (x, x + a_n)) \\ &= na_n. \end{aligned}$$

It follows that

$$\begin{aligned} \bar{D}f(x) &= \lim_{h \rightarrow 0} \sup_{0 < |t| \leq h} \left\{ \frac{f(x+t) - f(x)}{t} \right\} \\ &= \lim_{h \rightarrow 0} \sup_{0 < |t| \leq h} \frac{na_n}{a_n} \\ &\geq m. \end{aligned}$$

Since m was arbitrary, we have that

$$\bar{D}f(x) = \infty,$$

which is not finite, and by definition, x is not differentiable at x . Therefore, f fails to be differentiable at each point in E . \square

Question 6.33.

Solution. Let $\{f_n\}$ be a sequence of real-valued functions on $[a, b]$ that converges pointwise on $[a, b]$ to the real-valued function f . We wish to show that $TV(f) \leq \liminf TV(f_n)$. Fix $P = \{x_0, \dots, x_m\}$ be a partition of $[a, b]$. As $f_n \rightarrow f$ pointwise, we have

$$\begin{aligned} V(f, P) &= \sum_{k=0}^{m-1} |f(x_{k+1}) - f(x_k)| \\ &= \lim_{n \rightarrow \infty} \sum_{k=0}^{m-1} |f_n(x_{k+1}) - f_n(x_k)| \\ &= \lim_{n \rightarrow \infty} V(f_n, P). \end{aligned}$$

By the definition of total variation, it follows that

$$V(f_n, P) \leq TV(f_n),$$

for all n . Consequently, we obtain

$$V(f, P) \leq \liminf_{n \rightarrow \infty} TV(f_n),$$

and since P was arbitrary, we finally have that

$$TV(f) \leq \liminf_{n \rightarrow \infty} TV(f_n),$$

as desired. \square

Question 4. Royden 6.42.

Solution. Let f and g be real-valued functions, that are absolutely continuous functions on $[a, b]$. We wish to show that $f + g$ is absolutely continuous on $[a, b]$. Fix $\epsilon > 0$. As f and g are both absolutely continuous on $[a, b]$, there exist $\delta_f, \delta_g > 0$, such that for any finite disjoint open intervals $\{(a_k, b_k)\}_{k=1}^n$ in (a, b) ,

$$\begin{aligned} \sum_{k=1}^n [b_k - a_k] < \delta_f &\implies \sum_{k=1}^n |f(b_k) - f(a_k)| < \frac{\epsilon}{2} \\ \sum_{k=1}^n [b_k - a_k] < \delta_g &\implies \sum_{k=1}^n |g(b_k) - g(a_k)| < \frac{\epsilon}{2}. \end{aligned}$$

Define $\delta = \min(\delta_f, \delta_g)$. Let $\{(a_k, b_k)\}_{k=1}^n$ be a finite disjoint open intervals in (a, b) , such that $\sum_{k=1}^n [b_k - a_k] < \delta$. It follows that

$$\begin{aligned} \sum_{k=1}^n |f + g(b_k) - f + g(a_k)| &\leq \sum_{k=1}^n |f(b_k) - f(a_k)| + |g(b_k) - g(a_k)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Since ϵ was arbitrary, we have shown that $f + g$ is absolutely continuous on $[a, b]$.

Let f be a real-valued function, that is absolutely continuous on $[a, b]$. We show that cf , for any $c \in \mathbb{R}$, is absolutely continuous on $[a, b]$. Let $c = 0$. Then $cf = 0$, which can trivially be shown to be

absolutely continuous, as $f(c) = 0$ for any $c \in [a, b]$. Suppose $c \neq 0$. As f is absolutely continuous on $[a, b]$, there exists $\delta_f > 0$, such that for any finite disjoint open intervals $\{(a_k, b_k)\}_{k=1}^n$ in (a, b) ,

$$\sum_{k=1}^n |b_k - a_k| < \delta_f \implies \sum_{k=1}^n |f(b_k) - f(a_k)| < \frac{\epsilon}{|c|}.$$

Define $\delta = \delta_f$. Let $\{(a_k, b_k)\}_{k=1}^n$ be a finite disjoint open intervals in (a, b) such that $\sum_{k=1}^n [b_k - a_k] < \delta$. It follows that

$$\begin{aligned} \sum_{k=1}^n |cf(b_k) - cf(a_k)| &= |c| \sum_{k=1}^n |f(b_k) - f(a_k)| \\ &< |c| \frac{\epsilon}{|c|} = \epsilon. \end{aligned}$$

Since ϵ was arbitrary, combined with the $c = 0$ case, we have shown that cf , for any $c \in \mathbb{R}$, is absolutely continuous on $[a, b]$.

Let f be a real-valued function, that is absolutely continuous on $[a, b]$. We wish to show that f^2 is absolutely continuous on $[a, b]$. As f is absolutely continuous, f is continuous on $[a, b]$. Hence, by the Extreme Value Theorem, there exists M such that $|f| \leq M$ on $[a, b]$. Fix $\epsilon > 0$. As f is absolutely continuous on $[a, b]$, there exists $\delta_f > 0$, such that for any finite disjoint open intervals $\{(a_k, b_k)\}_{k=1}^n$ in (a, b) ,

$$\sum_{k=1}^n |b_k - a_k| < \delta_f \implies \sum_{k=1}^n |f(b_k) - f(a_k)| < \frac{\epsilon}{2M}.$$

Define $\delta = \delta_f$. Let $\{(a_k, b_k)\}_{k=1}^n$ be a finite disjoint open intervals in (a, b) such that $\sum_{k=1}^n [b_k - a_k] < \delta$. It follows that

$$\begin{aligned} \sum_{k=1}^n |f^2(b_k) - f^2(a_k)| &= \sum_{k=1}^n |f(b_k) - f(a_k)| |f(b_k) + f(a_k)| \\ &\leq 2M \sum_{k=1}^n |f(b_k) - f(a_k)| \\ &< 2M \frac{\epsilon}{2M} = \epsilon. \end{aligned}$$

Since ϵ was arbitrary, we have shown that f^2 is absolutely continuous on $[a, b]$.

Let f and g be real-valued functions, that are absolutely continuous on $[a, b]$. We wish to show that fg is absolutely continuous on $[a, b]$. Observe that

$$(f + g)^2 = f^2 + g^2 + 2fg,$$

which simplifies to

$$fg = -\frac{1}{2}((f + g)^2 + (-f^2) + (-g^2)).$$

As we have previously shown that a square of an absolutely continuous function is absolutely continuous, and a scalar multiple of an absolutely continuous function is absolutely continuous, we can deduce that fg is absolutely continuous on $[a, b]$. This completes the proof. \square

Question 4. 6.45.

Solution. Let f be a real-valued function, that is absolutely continuous on \mathbb{R} . Let g be a real-valued function, that is absolutely continuous and strictly monotone on $[a, b]$. Without the loss of generality, we assume that g is strictly increasing. We wish to show that $f \circ g$ is absolutely continuous on $[a, b]$. Fix $\epsilon > 0$. As f is absolutely continuous on \mathbb{R} , it is also absolutely continuous on $[g(a), g(b)]$, which

is a non-degenerate closed interval, as g is strictly increasing. there exists δ_f , such that for any finite disjoint open intervals $\{(a_k, b_k)\}_{k=1}^\infty$ in $(g(a), g(b))$,

$$\sum_{k=1}^n [b_k - a_k] < \delta_f \implies \sum_{k=1}^n |f(b_k) - f(a_k)| < \epsilon \quad (*).$$

As g is absolutely continuous, there exists δ_g , such that for any finite disjoint open intervals $\{(a_k, b_k)\}_{k=1}^\infty$ in (a, b) ,

$$\sum_{k=1}^n [b_k - a_k] < \delta_g \implies \sum_{k=1}^n |g(b_k) - g(a_k)| < \delta_f.$$

Define $\delta = \delta_g$. Let $\{(a_k, b_k)\}_{k=1}^n$ be a finite disjoint open intervals in (a, b) such that $\sum_{k=1}^n [b_k - a_k] < \delta_g$. It follows that $\sum_{k=1}^n [g(b_k) - g(a_k)] < \delta_f$. As g is strictly increasing, we observe that $\{(g(a_k), g(b_k))\}_{k=1}^n$ form a finite disjoint open intervals in $(g(a), g(b))$. Therefore, from $(*)$ it follows that

$$\sum_{k=1}^n |f \circ g(b_k) - f \circ g(a_k)| < \epsilon.$$

Since ϵ was arbitrary, we have shown that $f \circ g$ is absolutely continuous on $[a, b]$. \square

Question 6.56.

Solution. Let g be strictly increasing and absolutely continuous on $[a, b]$.

(i) Let O be an open subset of (a, b) . Then, O can be represented as a countable union of disjoint intervals in (a, b) :

$$O = \cup_{k=1}^\infty (a_k, b_k),$$

and since g is strictly increasing, we have $\{(g(a_k), g(b_k))\}_{k=1}^\infty$ forms a collection of disjoint intervals, and

$$g(O) = \cup_{k=1}^\infty (g(a_k), g(b_k)).$$

Therefore, by the countable additivity of measure, it follows that

$$m(g(O)) = \sum_{k=1}^\infty g(b_k) - g(a_k).$$

On the other hand, by the countable additivity of integration, and as g is absolutely continuous, we have

$$\begin{aligned} \int_O g'(x) dx &= \sum_{k=1}^\infty \int_{g(a_k)}^{g(b_k)} g'(x) dx \\ &= \sum_{k=1}^\infty g(b_k) - g(a_k). \end{aligned}$$

Therefore, $m(g(O)) = \int_O g'(x) dx$, as desired.

(ii)

(iii) Let E be a measure zero set. We have previously shown that a continuous map carries a measure zero set to a measure zero set. Hence, $g(E)$ has measure zero. Furthermore, we have that an integral over a measure zero set is zero. Therefore, we have that $m(g(E)) = 0 = \int_E g'(x) dx$ as desired.

(iv) Let A be any measurable set of $[a, b]$. By the outer approximation of a measurable set via $G - \delta$ set, there exists $G - \delta$ set G such that $m(G \setminus A) = 0$ and $A \subseteq G$. Then, by the finite additivity of measure we have

$$\begin{aligned} m(g(A)) &= m(g(G \setminus A \cup A)) \\ &= m(g(G \setminus A)) + m(g(A)) \\ &= m(A) \end{aligned}$$

(v)

(vi)

(vii)