

Diff Geo II: Problem Set II

Youngduck Choi *

Abstract

This work contains solutions for the problem set II.

Question 1-1.

1. In class, we used the fact that the curvature tensor can be expressed in terms of parallel transport. That is, we claimed that if $q \in M$ and if $\alpha: \mathbb{R}^2 \rightarrow M$ is a smooth map such that $\frac{\partial \alpha}{\partial x} = X$, $\frac{\partial \alpha}{\partial y} = Y$, then for all $Z \in T_q M$, we have

$$R(X, Y)Z = \lim_{s \rightarrow 0} \frac{Z - p_{\gamma_s}(Z)}{s^2},$$

where $\gamma_s: [0, 4] \rightarrow M$ is the image under α of the boundary of an $s \times s$ square, i.e.,

$$\gamma_s(t) = \begin{cases} \alpha(st, 0) & t \in [0, 1] \\ \alpha(s, s(t-1)) & t \in [1, 2] \\ \alpha(s(3-t), s) & t \in [2, 3] \\ \alpha(0, s(4-t)) & t \in [3, 4]. \end{cases}$$

Prove this fact.

(Hint: Construct a frame of vector fields $V_1, \dots, V_m \in \mathbf{V}(\alpha)$ such that $\nabla_X V_i = 0$ and $\nabla_Y V_i(u_1, 0) = 0$. Any vector field W along γ_s can be expressed as a linear combination of the V_i — when is W parallel?)

Solution.

Let $s > 0$. Set

$$V_i(\alpha(s_1, s_2)) = P_{\alpha(0, s_2), \alpha(s_1, s_2)} \circ P_{q, \alpha(0, s_2)}(V_i(q))$$

for all $1 \leq i \leq m$ and $(s_1, s_2) \in [0, s] \times [0, s]$. As $\frac{\partial \alpha}{\partial x} = X$, $\frac{\partial \alpha}{\partial y} = Y$, we have

$$\nabla_X V_i = 0 \quad \text{and} \quad \nabla_Y V_i(u_1, 0) = 0$$

for all i . Then, we see that any $W \in V(\gamma)$ can only change along 2nd part of the path. We compute

$$\begin{aligned} V_i(q)(P_\gamma W(q) - W(q)) &= \langle V_i, W \rangle \gamma_s(2) + \langle V_i, W \rangle \gamma_s(1) \\ &= \int_0^s \partial_t \langle V_i, W \rangle dt = \int_0^s V_i \cdot \nabla_Y W(1, t) dt \\ &= \int_0^s V_i \cdot \nabla_Y W(1, t) dt - \int_0^s \partial_r (V_i \cdot \nabla_Y W(r, t)) dr dt \\ &= - \int_0^s \int_0^s V_i \cdot \nabla_X \nabla_Y W(r, t) dr dt \end{aligned}$$

*Department of Mathematics, Courant Institute of Mathematical Sciences, yc1104@nyu.edu; If you find an error and want to share with me, you can reach me via email.

for all i . Now, with $W(0,0) = Z$, and $[X,Y] = 0$, letting $s \downarrow 0$, gives

$$\lim_{s \downarrow 0} \frac{\langle Z - P_\gamma(Z), V_i \rangle}{s^2} = \langle R(X,Y)Z, V_i \rangle$$

for all i , and hence,

$$R(X,Y)Z = \lim_{s \downarrow 0} \frac{Z - P_\gamma(Z)}{s^2}.$$

□

Question 1-2.

2. Suppose that M is a two-dimensional Riemannian manifold with curvature tensor R .

(a) Use the symmetries of the curvature tensor to show that if $p \in M$, $V, W \in T_p M$, then

$$K = \frac{\langle R(V, W)W, V \rangle}{\|V\|^2\|W\|^2 - 2\langle V, W \rangle^2}$$

is independent of V and W .

(b) Prove that if K is as above, then

$$R(X, Y)Z = K(\langle Y, Z \rangle X - \langle X, Z \rangle Y)$$

for all $X, Y, Z \in T_p M$.

Solution.

I believe there is a typo in the problem: 2 in the denominator should be gone.

(a) Let E_1, E_2 be orthonormal frame, so $V = V^1 E_1 + V^2 E_2$ and $W = W^1 E_1 + W^2 E_2$. Then,

$$\begin{aligned} \langle R(V, W)W, V \rangle &= \langle R(V^1 E_1 + V^2 E_2, W^1 E_1 + W^2 E_2)W, V \rangle \\ &= \langle V^1 W^2 R(E_1, E_2)W, V \rangle + \langle V^2 W^1 R(E_2, E_1)W, V \rangle \\ &= (V^1 W^2 - V^2 W^1)(W^1 V^2 \langle R(E_1, E_2)E_1, E_2 \rangle + W^2 V^1 \langle R(E_1, E_2)E_2, E_1 \rangle) \\ &= -(V^1 W^2 - V^2 W^1)^2 \langle R(E_1, E_2)E_1, E_2 \rangle \end{aligned} \tag{1}$$

where (??), in particular, holds by $R(E_1, E_1)W = R(E_2, E_2)W = 0$. Furthermore,

$$\|V\|^2\|W\|^2 - \langle V, W \rangle^2 = (V^1 W^2 - V^2 W^1)^2$$

and hence

$$\begin{aligned} K &= \frac{-(V^1 W^2 - V^2 W^1)^2 \langle R(E_1, E_2)E_1, E_2 \rangle}{(V^1 W^2 - V^2 W^1)^2} \\ &= \langle R(E_1, E_2)E_2, E_1 \rangle \end{aligned}$$

Therefore, we see that K only depend on the chosen frame, so K is independent of V and W .

(b) We compute

$$\begin{aligned} R(X, Y)Z &= \sum_{i,j,k=1}^2 X^i Y^j Z^k R(E_i, E_j)E_k \\ &= (X^1 Y^2 Z^1 - X^2 Y^1 Z^1)R(E_1, E_2)E_1 + (X^1 Y^2 Z^2 - X^2 Y^1 Z^2)R(E_1, E_2)E_2 \end{aligned}$$

and

$$\begin{aligned}
K(<Y, Z>X - <X, Z>Y) &= <R(E_1, E_2)E_2, E_1>X(Y^1Z^1 + Y^2Z^2) \\
&- <R(E_1, E_2)E_2, E_1>Y(X^1Z^1 + X^2Z^2) \\
&= <R(E_1, E_2)E_2, E_1>(X^1Y^2Z^2 - X^2Z^2Y^1)E_1 \\
&+ <R(E_1, E_2)E_2, E_1>(Y^1Z^1X^2 - X^1Y^2Z^1)E_2
\end{aligned}$$

Therefore,

$$\begin{aligned}
<R(X, Y)Z, E_1> &= <R(E_1, E_2)E_2, E_1>(X^1Y^2Z^2 - X^2Y^1Z^2) \\
&= <K(<Y, Z>X - <X, Z>Y), E_1>
\end{aligned}$$

Similarly,

$$<R(X, Y)Z, E_2> = <K(<Y, Z>X - <X, Z>Y), E_2>.$$

As the equality holds for the basis elements, we have

$$R(X, Y)Z = K(<Y, Z>X - <X, Z>Y)$$

as required. □

Question 1-3.

3. Find a sequence $\gamma_i: [0, 1] \rightarrow S^2$ of piecewise-smooth curves with the same endpoints such that the γ_i converge to a piecewise-smooth curve γ , but the parallel transport maps $P_{\gamma_i}: T_p M \rightarrow T_q M$ do not converge to the map P_γ .

Solution.

All triangles in the discussion are great circle triangles. From class, we saw that the parallel transport along a path of triangle, starting at a vertex and coming back to the chosen vertex, is given by a rotation of $\theta - \pi$ where θ is the sum of three angles of the triangle, formed by the path. To discuss pointwise convergence, we choose the geodesic distance on S^2 . Let p be the north pole of S^2 . Then, choose a triangle path, starting at p and coming back to p such that $\theta - \pi > 0$. Now, for each $i \in \mathbb{N}$, choose a triangle path, γ_i^* , whose angle sum is θ' , yet $\theta - \pi = k\theta'$ for some $k \in \mathbb{N}$, and the path of the triangle is contained in $B(p, \frac{1}{i})$, and set $\gamma_i = (\gamma_i^*)^k$, which is the standard path composition of γ_i^* k times. Then, set γ be the trivial path from p to p itself. Then, we see that γ_i converges to γ pointwise, as for any $t \in [0, 1]$, for any $\epsilon > 0$, there exists i such that $d(\gamma_i(t), \gamma(t)) < \epsilon$, whenever $n \geq i$. Furthermore, P_{γ_i} is given by a rotation of $\theta - \pi > 0$, yet $P_\gamma = I$, so we see that P_{γ_i} does not converge to P_γ as a linear map. Hence, we have the desired construction. \square

Question 1-4.

Let $M \subset \mathbb{R}^3$ be the cone

$$M = \{(r \cos \theta \sin \phi, r \sin \theta \sin \phi, r \cos \phi) \mid r > 0, \theta \in [0, 2\pi)\}$$

as in the first problem set.

4. Construct an isometry from the complement of the line $\theta = 0$ to a subset of \mathbb{R}^2 . Use this isometry to calculate the parallel transport map $P_{0,2\pi}$ from problem 5 of the first problem set.

Solution.

Cutting the cone and unfolding it, and using elementary geometry tell us that, in order to have an isometry, we must satisfy $r \sin(\phi)\theta = r\theta'$, because the radius of the circle changes from $r \sin(\phi)$ to r , by unfolding. As a parallel transport along any curve in \mathbb{R}^2 is an identity, we have that a parallel transport on the image is given by $(\cos(\theta'), -r \sin(\theta'); \sin(\theta'), -r \cos(\theta'))$.