

---

# ProbLimI: Problem Set VIII

---

Youngduck Choi  
CIMS  
New York University  
yc1104@nyu.edu

## Abstract

This work contains solutions to the exercises of the problem set VIII. The chosen problems are 1, 2, and 3.

### Question 1.

1. Let  $X_k$  be i.i.d. random variables and let  $S_n = \sum_{k=1}^n X_k$ . Show that if  $S_n/n$  converges a.s. as  $n \rightarrow \infty$ , then  $X_1$  is necessarily integrable.

### Solution.

Suppose for sake of contradiction that  $\mathbb{E}|X_1| = \infty$ . By Tonelli,

$$\begin{aligned}\mathbb{E}|X_1| &= \int_{\Omega} |X_1| d\mathbb{P} = \int_{\Omega} \int_0^{|X_1|} 1 dy d\mathbb{P} = \int_{\Omega} \int_0^{\infty} 1_{\{|X_1| > y\}} dy d\mathbb{P} \\ &= \int_0^{\infty} \int_{\Omega} 1_{\{|X_1| > y\}} d\mathbb{P} dy = \int_0^{\infty} \mathbb{P}(|X_1| > y) dy.\end{aligned}$$

Since

$$\int_0^n \mathbb{P}(|X_1| > y) dy \leq \sum_{k=0}^n \mathbb{P}(|X_1| > k)$$

for all  $n \geq 1$ , from (??),

$$\sum_{k=0}^{\infty} \mathbb{P}(|X_1| > k) = \infty.$$

Therefore, by Borel Cantelli II,

$$\mathbb{P}(|X_n| \geq n \text{ i.o.}) = 1.$$

Now, observe that

$$\frac{S_n}{n} - \frac{S_{n+1}}{n+1} = \frac{S_n}{n(n+1)} - \frac{X_{n+1}}{n+1}$$

and hence, by reverse triangle inequality,

$$\left| \frac{S_n}{n} - \frac{S_{n+1}}{n+1} \right| \geq \left\| \frac{S_n}{n(n+1)} - \frac{X_{n+1}}{n+1} \right\|$$

for all  $n \geq 1$ . Suppose

$$w \in \left\{ \frac{S_n}{n} \text{ converges} \right\} \cap \{ |X_n| \geq n \text{ i.o.} \}.$$

Then, by the convergence of  $\frac{S_n(w)}{n}$ , there exists  $0 < \delta < 1$  such that

$$\left| \frac{S_n(w)}{n(n+1)} \right| < \delta$$

for all  $n$  large enough, and hence

$$\left| \frac{S_n(w)}{n} - \frac{S_{n+1}(w)}{n+1} \right| > 1 - \delta \text{ i.o.,}$$

which contradicts the fact that  $\frac{S_n(w)}{n}$  converges. Hence,

$$\left\{ \frac{S_n}{n} \text{ converges} \right\} \cap \{ |X_n| \geq n \text{ i.o.} \} = \emptyset$$

so

$$\mathbb{P}\left(\frac{S_n}{n} \text{ converges}\right) = 0$$

which contradicts that  $\frac{S_n}{n}$  converges a.s. Hence,  $\mathbb{E}|X_1| < \infty$ , i.e.  $X_1$  is integrable. □

## Question 2.

2. Let  $S_n = \sum_{k=1}^n X_k$  for i.i.d. r.v.'s  $X_k$ .

- (a) Prove that if  $\frac{d}{dt}\Phi_{X_1}(0) = a + ib \in \mathbb{C}$  then  $a = 0$  and  $S_n/n \xrightarrow{P} b$  as  $n \rightarrow \infty$ .
- (b) Prove that  $S_n/n \xrightarrow{P} b \in \mathbb{R}$  implies that  $\Phi_{X_1}(x_k)^{\lfloor t/x_k \rfloor} \rightarrow \exp(ib t)$  as  $x_k \downarrow 0$  and  $\Phi_{X_1}(x_k)^{\lfloor t/x_k \rfloor} \rightarrow \exp(-ib t)$  as  $x_k \uparrow 0$  for all  $t > 0$ . Deduce that  $\frac{d}{dt}\Phi_{X_1}(0) = ib$ , and conclude that  $S_n/n \xrightarrow{P} b \in \mathbb{R}$  if and only if  $\Phi_{X_1}(t)$  is differentiable at  $t = 0$ .
- (c) Give an example of a random variable  $X$  with  $\Phi_X(t)$  differentiable at  $t = 0$  yet  $\mathbb{E}|X| = \infty$ .

## Solution.

(a) Let  $\frac{d}{dt}\Phi_{X_1}(0) = a + ib$ . Then, there exists  $t_0 > 0$  small enough such that

$$\Phi_{X_1}(t_0) = 1 + (a + ib)t_0 + o(|t_0|) \quad \text{and} \quad \Phi_{X_1}(-t_0) = 1 - (a + ib)t_0 + o(|t_0|). \quad (1)$$

As  $\Phi_{X_1}(t) = \overline{\Phi_{X_1}(-t)}$  for any  $t \in \mathbb{R}$ , from (??),

$$(1 + at_0) + bt_0i = (1 - at_0) + bt_0i$$

which implies  $a = 0$ . Observe that

$$\Phi_{n^{-1}S_n}(t) = \Phi_{X_1}\left(\frac{t}{n}\right)^n = \left(1 + \frac{bti}{n} + o\left(\frac{1}{n}\right)\right)^n$$

for any  $t \in \mathbb{R}$  and  $n \in \mathbb{N}$ . Taking  $n \rightarrow \infty$ , we see that

$$\Phi_{n^{-1}S_n}(t) = e^{ibt}$$

so by continuity theorem,  $\frac{S_n}{n} \rightarrow_D b$  as  $n \rightarrow \infty$ . From pset 4-1,  $\frac{S_n}{n} \rightarrow_p b$  as  $n \rightarrow \infty$ .

(b) Since convergence in probability implies convergence in distribution, and by continuity theorem,

$$\Phi_{X_1}\left(\frac{t}{n}\right)^n \rightarrow e^{ibt}$$

for all  $t \in \mathbb{R}$ . Let  $t \in \mathbb{R}$ , and  $\epsilon > 0$ . By uniform continuity of  $\Phi_{X_1}$ , there exists  $\delta > 0$ ,

$$\left|\frac{t}{n_k} - x_k\right| < \delta \implies |\Phi_{X_1}(x_k) - \Phi_{X_1}\left(\frac{t}{n_k}\right)| < \epsilon.$$

where  $\{n_k\}$  is any subsequence of  $\{n\}$ . Since  $x_k \rightarrow 0$  and  $n_k \rightarrow \infty$ ,

$$|\Phi_{X_1}(x_k) - \Phi_{X_1}\left(\frac{t}{n_k}\right)| < \epsilon$$

and

$$\left(\Phi_{X_1}\left(\frac{t}{n_k}\right) - \epsilon\right)^{n_k - n_k + \lfloor \frac{t}{x_k} \rfloor} \leq \Phi_{X_1}(x_k)^{\lfloor \frac{t}{x_k} \rfloor} \leq \left(\Phi_{X_1}\left(\frac{t}{n_k}\right) + \epsilon\right)^{n_k - n_k + \lfloor \frac{t}{x_k} \rfloor}$$

for all  $k$  large enough. Since  $\epsilon > 0$  was arbitrary, we can take  $\{\epsilon_k\}$  such that  $\epsilon_k \rightarrow 0$ . Then, by the givens,

$$\Phi_{X_1}(x_k)^{\lfloor \frac{t}{x_k} \rfloor} \rightarrow e^{ibt}$$

as  $k \rightarrow \infty$ . For the negative part, consider the conjugate relation as part a, then carry out the exact same limit process.

To conclude, conversely, suppose  $\Phi\left(\frac{t}{n}\right)^n \rightarrow e^{ibt}$ , then by taking logs  $n \log(\Phi\left(\frac{t}{n}\right)) \rightarrow ibt$ . By differentiability of  $\log(z)$  for  $z = 1$  on the complex plane,  $n(\Phi\left(\frac{t}{n}\right) - 1) \rightarrow ibt$ . Hence, by the above result and (a), we have that  $\Phi(t)$  is differentiable at  $t = 0$ .

(c) Define each  $X$  such that

$$\mathbb{P}(X = (-1)^k k) = C \frac{k}{k^2 \log(k)}$$

for all  $k \geq 2$ , where  $C$  is the normalization constant. Then,

$$\mathbb{E}|X| = \sum_{k=2}^{\infty} \frac{C}{k \log(k)} = \infty.$$

Furthermore,

$$n\mathbb{P}(|X| > n) = n \sum_{k=n+1}^{\infty} \frac{C}{k^2 \log(k)} \rightarrow 0$$

and

$$\mathbb{E}(X 1_{\{|X| \leq n\}}) = \sum_{k=2}^n (-1)^k \frac{C}{k \log(k)} \text{ converges}$$

as  $n \rightarrow \infty$ . Hence, from thm 2.2.7 in Durrett,

$$\frac{S_n}{n} \rightarrow_P \sum_{k=2}^{\infty} (-1)^k \frac{C}{k \log(k)}$$

and by (b),  $\Phi_X$  is differentiable at  $t = 0$ . Therefore,  $X$  is the desired construction. □

**Question 4.**

4. Define  $f_n(x) = e^{ix} - \sum_{k=0}^n (ix)^k / k!$  for  $x \in \mathbb{R}$  and  $n \geq 0$ .

(a) Show that  $|f_n(x)| \leq \min \{2|x|^n/n!, |x|^{n+1}/(n+1)!\}$  for all  $x$  and  $n$ .

(b) Use this to show that if  $\mathbb{E}|X|^n < \infty$  then

$$\left| \Phi_X(t) - \sum_{k=0}^n (it)^k \mathbb{E}[X^k] / k! \right| \leq |t|^n \mathbb{E} \left[ \min \{2|X|^n/n!, |t||X|^{n+1}/(n+1)!\} \right].$$

Explain the implication this has for a CLT for i.i.d. r.v.'s  $X_k$  with  $\mathbb{E}|X_k|^n < \infty$ .

**Solution.**

Let  $x \in \mathbb{R}$ . By integration by parts,

$$\int_0^x (x-s)^n e^{is} ds = \frac{x^{n+1}}{n+1} + \frac{i}{n+1} \int_0^x (x-s)^{n+1} e^{is} ds \quad (2)$$

for each  $n \geq 0$ . If  $n = 0$ , then

$$x + i \int_0^x (x-s) e^{is} ds = \int_0^x e^{is} ds = \frac{e^{ix} - 1}{i}$$

and hence

$$e^{ix} = 1 + ix + i^2 \int_0^x (x-s) e^{is} ds.$$

Suppose for some  $n > 0$

$$e^{ix} - \sum_{k=0}^n \frac{(ix)^k}{k!} = \frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds \quad (3)$$

Then, combined with (??),

$$\begin{aligned} e^{ix} - \sum_{k=0}^{n+1} \frac{(ix)^k}{k!} &= \frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds - \frac{(ix)^{n+1}}{(n+1)!} \\ &= \frac{i^{n+1}}{n!} \left( \int_0^x (x-s)^n e^{is} ds - \frac{x^{n+1}}{(n+1)} \right) \\ &= \frac{i^{n+2}}{(n+1)!} \int_0^x (x-s)^{n+1} e^{is} ds. \end{aligned}$$

Hence, by induction, (??) holds for all  $n \geq 0$ . If  $x \geq 0$ , then

$$\left| \frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds \right| \leq \frac{1}{n!} \int_0^x |(x-s)^n| ds = \frac{1}{n!} \int_0^x (x-s)^n ds = \frac{1}{(n+1)!} |x|^{n+1}.$$

If  $x < 0$ , then

$$\begin{aligned} \left| \frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds \right| &= \left| \frac{i^{n+1}}{n!} \int_x^0 (x-s)^n e^{is} ds \right| \leq \frac{1}{n!} \int_x^0 |(x-s)^n e^{is}| ds \\ &\leq \frac{1}{n!} \int_x^0 (s-x)^n ds = \frac{1}{(n+1)!} (-x)^{n+1} = \frac{1}{(n+1)!} |x|^{n+1}. \end{aligned}$$

Therefore,

$$|f_n(x)| \leq \frac{1}{(n+1)!} |x|^{n+1} \quad (4)$$

for any  $n \geq 0$ . Now, again by integration by parts,

$$\frac{i}{n} \int_0^x (x-s)^n e^{is} ds = -\frac{x^n}{n} + \int_0^x (x-s)^{n-1} e^{is} ds$$

and hence

$$\frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds = \frac{i^n}{(n-1)!} \int_0^x (x-s)^{n-1} (e^{is} - 1) ds$$

for any  $n \geq 1$ . If  $x \geq 0$ , then

$$\begin{aligned} \left| \frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds \right| &= \left| \frac{i^n}{(n-1)!} \int_0^x (x-s)^{n-1} (e^{is} - 1) ds \right| \\ &\leq \frac{1}{(n-1)!} \int_0^x |(x-s)^{n-1}| |e^{is} - 1| ds \\ &\leq \frac{2}{(n-1)!} \int_0^x (x-s)^{n-1} ds = \frac{2}{n!} |x|^n. \end{aligned}$$

If  $x < 0$ , then

$$\begin{aligned} \left| \frac{i^{n+1}}{n!} \int_0^x (x-s)^n e^{is} ds \right| &= \left| \frac{i^n}{(n-1)!} \int_0^x (x-s)^{n-1} (e^{is} - 1) ds \right| \\ &\leq \frac{1}{(n-1)!} \int_0^x |(x-s)^{n-1}| |e^{is} - 1| ds \\ &\leq \frac{2}{(n-1)!} \int_0^x (x-s)^{n-1} ds = \frac{2}{n!} |x|^n. \end{aligned}$$

Therefore, combined with (??),

$$|f_n(x)| \leq \min\left(\frac{2|x|^n}{n!}, \frac{|x|^{n+1}}{(n+1)!}\right)$$

for all  $x \in \mathbb{R}$  and  $n \geq 1$ .

(ii) From the above estimate,

$$\left| e^{itX} - \sum_{k=0}^n \frac{(itX)^k}{k!} \right| \leq \min\left(\frac{2|tX|^n}{n!}, \frac{|tX|^{n+1}}{(n+1)!}\right)$$

and hence, by Jensen's inequality, and linearity of expectation, which is granted by  $\mathbb{E}|X|^n < \infty$ ,

$$\begin{aligned} \left| \Phi_X(t) - \sum_{k=0}^n \frac{\mathbb{E}(itX)^k}{k!} \right| &\leq \mathbb{E} \left| e^{itX} - \sum_{k=0}^n \frac{(itX)^k}{k!} \right| \\ &\leq \mathbb{E} \left[ \min\left(\frac{2|tX|^n}{n!}, \frac{|tX|^{n+1}}{(n+1)!}\right) \right] \\ &= |t|^n \mathbb{E} \left[ \min\left(\frac{2|X|^n}{n!}, \frac{|t||X|^{n+1}}{(n+1)!}\right) \right] \end{aligned}$$

for all  $t \in \mathbb{R}$ .

Suppose  $\mathbb{E}|X|^2 < \infty$ . Let  $t \in \mathbb{R}$ . Then,

$$\Phi_X(t) = 1 + it\mathbb{E}X - t^2\mathbb{E}\frac{X^2}{2} + \Psi(t)$$

for some constant  $\Psi$ , dependent on  $t$ . By the established result,

$$|\Psi(t)| \leq t^2 \mathbb{E}(|t||X|^3 \wedge 2|X|^2)$$

where denotes the minimum operation. Observe that as  $t \rightarrow 0$ ,

$$|t||X|^3 \wedge 2|X|^2 \rightarrow 0 \text{ everywhere and } |t||X|^3 \wedge 2|X|^2 \leq 2|X|^2 \text{ everywhere.}$$

Since  $\mathbb{E}|X|^2 < \infty$ , by DCT,

$$\frac{|\Psi(t)|}{t^2} \rightarrow 0$$

so

$$\Phi_X(t) = 1 + it\mathbb{E}X - t^2\mathbb{E}\frac{X^2}{2} + o(t^2).$$

This shows that we can prove the CLT for the iid case without assuming the finite third moment.  
 $\square$