

Reactive MD-force field: 2013 C

```

39      ! Number of general parameters
50.0000 !p(boc1)
 9.5469 !p(boc2)
26.5405 !p(coa2)
 1.9940 !p(trip4)
 2.3157 !p(trip3)
70.0000 !kc2
 1.0588 !p(ovun6)
 4.6000 !p(trip2)
12.1176 !p(ovun7)
13.3056 !p(ovun8)
-101.4874 !p(trip1)
 0.0000 !Lower Taper-radius (swa)
10.0000 !Upper Taper-radius (swb)
 0.0000 !not used
33.8667 !p(val7)
 6.0891 !p(lp1)
 1.0563 !p(val9)
 2.0384 !p(val10)
 6.1431 !not used
 6.9290 !p(pen2)
 0.3989 !p(pen3)
 3.9954 !p(pen4)
 0.0000 !not used
 5.7796 !p(tor2)
10.0000 !p(tor3)
 1.9487 !p(tor4)
 0.0000 !not used
 2.1645 !p(cot2)
 1.5591 !p(vdW1)
 0.1000 !Cutoff for bond order*100 (cutoff)
 2.1365 !p(coa4)
 0.6991 !p(ovun4)
50.0000 !p(ovun3)
 1.8512 !p(val8)
 0.0000 !not used
 0.0000 !not used
 0.0000 !not used
 0.0000 !not used
 2.6962 !p(coa3)
1      ! Nr of atoms; atomID;ro(sigma); Val;atom mass;Rvdw;Dij;gamma
      alfa;gamma(w);Val(angle);p(ovun5);n.u.;chiEEM;etaEEM;n.u.
      ro(pipi);p(lp2);Heat increment;p(boc4);p(boc3);p(boc5),n.u.;n.u.
      p(ovun2);p(val3);n.u.;Val(boc);p(val5);n.u.;n.u.;n.u.
C      1.3674  4.0000 12.0000  2.0453  0.1444  0.7920  1.1706  4.0000
      9.0000  1.5000  4.0000 27.5134 79.5548  6.7897  6.0000  0.0000
      1.1168  0.0000 181.0000 14.2732 24.4406  6.7313  0.8563  0.0000
      -4.1021  5.0000  1.0564  4.0000  2.9663  0.0000  0.0000  0.0000
1      ! Nr of bonds; at1;at2;De(sigma);De(pi);De(pipi);p(be1);p(b
      p(be2);p(bo3);p(bo4);n.u.;p(bo1);p(bo2)
1 1 80.8865 107.9944 52.0636  0.5218 -0.3636  1.0000 34.9876  0.7769
      6.1244 -0.1693  8.0804  1.0000 -0.0586  8.1850  1.0000  0.0000
0      ! Nr of off-diagonal terms. at1;at2;Dij;RvdW;alfa;ro(sigma);r
1      ! Nr of angles. at1;at2;at3;Thetao,o;p(val1);p(val2);p(coal);
1 1 1 74.9085 44.7514  0.9144  0.0000  0.0050  0.3556  2.5715
1      ! Nr of torsions. at1;at2;at3;at4;;V1;V2;V3;p(tor1);p(cot1);n
1 1 1 1 2.1207 26.8713  0.5160 -9.0000 -2.8394  0.0000  0.0000
0      ! Nr of hydrogen bonds. at1;at2;at3;r(hb);p(hb1);p(hb2);p(hb3

```