

# **Analysis 2**

yehelip

# Contents

<b>1</b>	<b>Getting Used to Integrals</b>	<b>4</b>
1.1	Suppose $f$ is Riemann integrable then $f^2$ is also Riemann integrable . . . . .	4
1.2	If $f$ is Continuous Then $f$ is Integrable . . . . .	4
<b>2</b>	<b>Intermediate Value Theorem for Integrals</b>	<b>5</b>
<b>3</b>	<b>Fundamental Theorem of Calculus</b>	<b>6</b>
3.1	Part One . . . . .	6
3.2	Part Two . . . . .	6
<b>4</b>	<b>Find Formula For Length of Continuous Graph</b>	<b>8</b>
<b>5</b>	<b>The Limit Comparison Test</b>	<b>9</b>
5.1	Proof . . . . .	9
<b>6</b>	<b>Some Practise</b>	<b>10</b>
6.1	Limits With Series: Find $\lim_{n \rightarrow \infty} \left( \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \right)$ . . . . .	10
6.2	Check convergion: $\int_{\frac{1}{2}}^1 \frac{1}{x\sqrt{1-x}}$ . . . . .	10
<b>7</b>	<b>Absolute Convergance Implies Conditional Convergance</b>	<b>11</b>
<b>8</b>	<b>Convergence Tests</b>	<b>12</b>
8.1	Dirichlet's Test . . . . .	12
8.1.1	Proof . . . . .	12
8.2	Abel's Test . . . . .	13
8.2.1	Proof . . . . .	13
8.3	Root And Quotient Test . . . . .	13
<b>9</b>	<b>Rabbe's Test</b>	<b>13</b>
<b>10</b>	<b>Integral Test for Series</b>	<b>14</b>
<b>11</b>	<b>Cauchy Condensation Test</b>	<b>15</b>
<b>12</b>	<b>Leibniz's Test</b>	<b>16</b>
<b>13</b>	<b>Riemann Series Theorem</b>	<b>17</b>
13.1	Dini's Theorem . . . . .	17
<b>14</b>	<b>Properties of Uniformly Converging Function Sequences</b>	<b>18</b>
14.1	Continuity . . . . .	18
14.2	Integrability . . . . .	18
14.3	Differentiability . . . . .	18
<b>15</b>	<b>Weierstrass M-Test</b>	<b>19</b>
15.1	Proof . . . . .	19
<b>16</b>	<b>Power Series Theorems</b>	<b>20</b>
16.1	Some exercises . . . . .	20
<b>17</b>	<b>Cauchy–Hadamard + D’alembert Theorem</b>	<b>21</b>
17.1	Proof the 1st . . . . .	21
17.2	Proof the 2nd . . . . .	21

<b>18 A Note on the Taylor Series</b>	<b>22</b>
<b>19 Continuous Partial Derivatives imply Differentiability</b>	<b>23</b>
19.1 Semi-Proof . . . . .	23
19.2 Note on Differentiability . . . . .	23
<b>20 Leibniz integral rule</b>	<b>24</b>
20.1 Lemma . . . . .	24
20.2 The Rule . . . . .	24
20.3 Generalization . . . . .	25
<b>21 Fubini's Theorem</b>	<b>26</b>
21.1 Proof . . . . .	26
<b>22 The Chain Rule</b>	<b>27</b>
22.1 Proof . . . . .	27
22.2 Corollary . . . . .	27
<b>23 Substitution For Multiple Variables</b>	<b>28</b>
<b>24 Calculate <math>\iint_{-\infty}^{\infty} e^{-x^2}</math></b>	<b>29</b>

# 1 Getting Used to Integrals

## 1.1 Suppose $f$ is Riemann integrable then $f^2$ is also Riemann integrable

First recall the definition of a Riemann integrable function. A function  $f$  is Riemann integrable on  $[a, b]$  iff exists  $I \in \mathbb{R}$  such that for all  $\varepsilon > 0$  exists  $\delta > 0$  such that for any partition  $X = (x_0, x_1, \dots, x_n)$  with  $\lambda(X) < \delta$  any sequence  $(c_1, c_2, \dots, c_n)$  such that  $c_i \in [x_{i-1}, x_i]$  also satisfies :

$$\left| \sum_{i=1}^n f(c_i) \Delta X_i - I \right| < \varepsilon$$

And we denote  $I = \int_a^b f(x) dx$ . Consider  $f^2$ :

$$\begin{aligned} U(f, P) - D(f, P) &< \varepsilon \\ \Rightarrow U(f, P) &< D(f, P) + \varepsilon \\ \Rightarrow U(f, P)^2 &< D(f, P)^2 + 2\varepsilon D(f, P) + \varepsilon^2 \\ \Rightarrow U(f, P)^2 - D(f, P)^2 &< \varepsilon(2D(f, P) + \varepsilon) \end{aligned}$$

Since  $f$  is integrable we know  $2D(f, P) + \varepsilon$  is bounded. Denote the bound  $M$ . Let  $\varepsilon > 0$ . Choosing the  $\delta$  that matches  $\varepsilon_\delta = \min(\frac{\varepsilon}{2M+1}, 1)$  under  $f$ 's integrability we get:

$$\begin{aligned} U(f, P) - D(f, P) &< \varepsilon_\delta \\ \Rightarrow U(f, P)^2 - D(f, P)^2 &< \varepsilon_\delta(2D(f, P) + \varepsilon_\delta) \\ \Rightarrow U(f, P)^2 - D(f, P)^2 &< \frac{\varepsilon}{2M+1}(2M+1) = \varepsilon \\ \Rightarrow U(f^2, P) - D(f^2, P) &< \varepsilon \end{aligned}$$

Since Darboux integrability implies Riemann integrability we finished.

## 1.2 If $f$ is Continuous Then $f$ is Integrable

Let  $f$  be continuous on  $[a, b]^*$ . By Cantor-Heine it is uniformly continuous.

$$U(f, P) - L(f, P) = \sum_{i=1}^n (M_i - m_i) \Delta X_i <^* \sum_{i=1}^n \varepsilon \Delta X_i = \varepsilon(b-a)$$

(\*) This is because by definition

$$\forall \varepsilon > 0: \exists \delta > 0: d(x, y) < \delta \rightarrow d(f(x), f(y)) < \varepsilon$$

And that delta is exactly what we wanted.

---

\*Think about the case  $(a, b)$

## 2 Intermediate Value Theorem for Integrals

Let  $f$  be a continuous function on  $[a, b]$  then exists  $x_0 \in [a, b]$  such that:

$$\int_a^b f(x)dx = f(x_0)(b-a)$$

We know  $f$  is continuous so it is Riemann integrable. We know by Weierstrass's theorem that since  $f$  is continuous it has a minimum and maximum in  $[a, b]$  which we'll denote  $m, M$ .

$$\begin{aligned} m(b-a) &\leq \int_a^b f(x) dx \leq M(b-a) \\ \Rightarrow m &\leq \frac{\int_a^b f(x) dx}{b-a} \leq M \end{aligned}$$

Let  $c = \frac{\int_a^b f(x) dx}{b-a}$ . By the intermediate value theorem we know that an  $x_0 \in (a, b)$  exists such that  $f(x_0) = c$  and thus:

$$\int_a^b f(x) dx = f(x_0)(b-a)$$

### 3 Fundamental Theorem of Calculus

Let  $f$  be a continuous real-valued function defined on a closed interval  $[a, b]$ . Let  $F$  be the function defined as:

$$F(x) = \int_a^x f(t) dt \quad \forall x \in [a, b]$$

Then  $F$  is uniformly continuous on  $[a, b]$  and differentiable on the open interval  $(a, b)$ , and

$$F'(x) = f(x) \quad \forall x \in (a, b)$$

If  $f$  is Riemann integrable on  $[a, b]$  then:

$$\int_a^b f(x) dx = F(b) - F(a)$$

#### 3.1 Part One

For any  $x_1, x_1 + \Delta x \in [a, b]$  we get:

$$F(x_1 + \Delta x) - F(x_1) = \int_a^{x_1 + \Delta x} f(t) dt - \int_a^{x_1} f(t) dt = \int_{x_1}^{x_1 + \Delta x} f(t) dt$$

According to the mean value theorem for integration we get that for  $c \in [x_1, x_1 + \Delta x]$ :

$$\begin{aligned} \int_{x_1}^{x_1 + \Delta x} f(t) dt &= f(c) \Delta x \\ \lim_{\Delta x \rightarrow 0} \frac{F(x_1 + \Delta x) - F(x_1)}{\Delta x} &= \lim_{\Delta x \rightarrow 0} f(c) = f(x_1) \end{aligned}$$

And thus according to the squeeze theorem and  $f$ 's continuity  $F'(x_1) = f(x_1)$

#### 3.2 Part Two

Let  $P = (x_0, x_1, \dots, x_n)$  be a partition of  $[a, b]$  such that  $(x_0, x_n) = (a, b)$

$$\begin{aligned} F(b) - F(a) &= F(x_n) - F(x_0) = [F(x_n) - F(x_{n-1})] + \dots + [F(x_1) - F(x_0)] \\ &= [F(x_n) - F(x_{n-1})] + [F(x_{n-1}) - F(x_{n-2})] + \dots + [F(x_1) - F(x_0)] \\ &= \sum_{i=1}^n [F(x_i) - F(x_{i-1})] \end{aligned}$$

Since  $F$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  according to Lagrange's theorem on  $[x_i, x_{i-1}]$  where  $c_i \in [x_i, x_{i-1}]$  we get

$$\sum_{i=1}^n [F(x_i) - F(x_{i-1})] = \sum_{i=1}^n [F'(c_i)(x_i - x_{i-1})]$$

According to part one we get that  $F'(c_i) = f(c_i)$  and so

$$\begin{aligned}
 F(b) - F(a) &= \sum_{i=1}^n [f(c_i)(\Delta x_i)] \\
 &\iff \\
 \lim_{\|\Delta x_i\| \rightarrow 0} (F(b) - F(a)) &= \lim_{\|\Delta x_i\| \rightarrow 0} \left( \sum_{i=1}^n [f(c_i)(\Delta x_i)] \right) \\
 &\iff \\
 F(b) - F(a) &= \int_a^b f(x) \, dx
 \end{aligned}$$

## 4 Find Formula For Length of Continuous Graph

Approximating the length of a graph using the pythagorean theorem for partition  $X = (x_0, x_1, \dots, x_n)$  we get:

$$\sum_{i=1}^n \sqrt{d(x_{i-1}, x_i)^2 + d(f(x_{i-1}), f(x_i))^2}$$

Assuming  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  by Lagrange's theorem we get:

$$\begin{aligned} \sum_{i=1}^n \sqrt{d(x_{i-1}, x_i)^2 + d(f(x_{i-1}), f(x_i))^2} &= \sum_{i=1}^n \sqrt{(x_i - x_{i-1})^2 + (f'(c_i)(x_i - x_{i-1}))^2} \\ &= \sum_{i=1}^n \sqrt{(x_i - x_{i-1})^2 (1 + (f'(c_i))^2)} \\ &= \sum_{i=1}^n \sqrt{1 + (f'(c_i))^2} \Delta X_i \end{aligned}$$

We can see that this summation is matching the integral

$$\lim_{\|\Delta x_i\| \rightarrow 0} \sum_{i=1}^n \sqrt{1 + (f'(c_i))^2} \Delta X_i = \int_a^b \sqrt{1 + (f'(x))^2} dx$$

The result of this integral will give us the length of any continuous graph.



## 5 The Limit Comparison Test

Let  $f, g$  be two integrable positive functions on  $[a, M]$  for any  $M \in \mathbb{R}$ . Suppose:

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = c$$

If  $c \in (0, \infty)$  then either both series converge or both series diverge. And if  $c = 0$  the convergence of  $f$  implies the convergence of  $g$ . If  $c = \infty$  the opposite is true.

### 5.1 Proof

Let  $\varepsilon > 0$  we know that  $\exists x_0 \in \mathbb{R}: \forall x > x_0$ :

$$g(x)(c - \varepsilon) < f(x) < g(x)(c + \varepsilon)$$

Then if  $g(x)$  converges then  $f(x)$  converges by the squeeze theorem. Similarly if  $g$  diverges we know that

$$g(x)(c - \varepsilon) < f(x)$$

So from a certain point onwards  $f$  will meet the requirements of the direct comparison test and so will diverge.

## 6 Some Practise

**6.1 Limits With Series: Find**  $\lim_{n \rightarrow \infty} \left( \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \right)$

$$\begin{aligned} \lim_{n \rightarrow \infty} \left( \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \right) &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{n+k} \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{1 + \frac{k}{n}} \frac{1}{n} \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^n f\left(\frac{k}{n}\right) \frac{1}{n} \\ &= \int_0^1 \frac{1}{1+x} \end{aligned}$$

**6.2 Check convergion:**  $\int_{\frac{1}{2}}^1 \frac{1}{x\sqrt{1-x}}$

This function seems to behave a lot like  $\frac{1}{\sqrt{1-x}}$  near 1 so let's compare them using the limit comparison test

$$\lim_{x \rightarrow 1^-} \frac{\frac{1}{x\sqrt{1-x}}}{\frac{1}{\sqrt{1-x}}} = \lim_{x \rightarrow 1^-} \frac{1}{x} = 1$$

So by the limit comparison test we know that the integral converges.

## 7 Absolute Convergence Implies Conditional Convergence

$$\int_a^\infty |f(x)| dx \text{ converges} \Rightarrow \int_a^\infty f(x) dx \text{ converges}$$

The first condition is called **"absolute convergence"**. We will prove it implies the second condition **"conditional convergence"**. Suppose  $f$  converges absolutely. Consider:

$$f^+ = \begin{cases} f(x) & f(x) > 0 \\ 0 & f(x) \leq 0 \end{cases}$$
$$f^- = \begin{cases} -f(x) & f(x) < 0 \\ 0 & f(x) \geq 0 \end{cases}$$

We know that  $|f| = f^+ + f^-$  converges and so by the direct comparison test we get that  $\int_a^\infty f^+, \int_a^\infty f^-$  converge and since  $f = f^+ - f^-$  we also get that  $\int_a^\infty f$  converges as well.

## 8 Convergence Tests

### 8.1 Dirichlet's Test

If  $a_n$  is a monotonic sequence and  $\lim_{n \rightarrow \infty} a_n = 0$  and  $\sum b_n$  is bounded by  $M$  then

$$\sum_{n=1}^{\infty} a_n b_n \text{ converges}$$

#### 8.1.1 Proof

Denote  $B_n = \sum_{k=1}^n b_k$  and by summation by parts we see that

$$\sum_{k=1}^n a_k b_k = a_n B_n - \sum_{k=1}^{n-1} (a_{k+1} - a_k) B_k$$

Since  $a_n$  converges to 0 and  $B_n$  is bounded  $\lim_{n \rightarrow \infty} a_n B_n$  exists. WLOG assume  $a_n$  is increasing, we can also see that

$$\sum_{k=1}^{n-1} |(a_{k+1} - a_k) B_k| \leq \sum_{k=1}^{n-1} M |a_{k+1} - a_k| \leq M \sum_{k=1}^{n-1} (a_{k+1} - a_k)$$

And

$$M \sum_{k=1}^{n-1} (a_{k+1} - a_k) = M(a_n - a_1)$$

Which converges to  $-Ma_1$  since  $\lim_{n \rightarrow \infty} a_n = 0$ . That means that this sequence is bounded. Which means that  $\sum_{k=1}^{n-1} |(a_{k+1} - a_k) B_k|$  is also bounded. It is also monotonic which means it converges. And if  $\sum_{k=1}^{n-1} (a_{k+1} - a_k) B_k$  converges absolutely it also converges conditionally which means  $\sum_{k=1}^{n-1} (a_{k+1} - a_k) B_k$  converges. And by additivity of limits we know  $\lim_{n \rightarrow \infty} \sum_{k=1}^n a_k b_k$  exists so  $\sum_{n=1}^{\infty} a_n b_n$  converges.

## 8.2 Abel's Test

Suppose  $\sum_1^\infty a_n$  converges, and  $b_n$  is monotone and bounded. Then  $\sum_1^\infty a_n b_n$  also converges.

### 8.2.1 Proof

We know  $b_n$  is monotone and bounded so it has a limit  $\lim_{n \rightarrow \infty} b_n = b$ . This implies  $\lim_{n \rightarrow \infty} b_n - b = 0$ . Since  $b_n - b$  is also monotonic we know by Dirichle's test that  $\sum_1^\infty a_n(b_n - b)$  converges. And by homogeneity of series we know that  $\sum_{n=1}^\infty a_n b$  converges as well. That means  $\sum_{n=1}^\infty (a_n b) + a_n(b_n - b)$  converges. So  $\sum_{n=1}^\infty (a_n b) + a_n(b_n - b) = \sum_{n=1}^\infty a_n b_n$  converges.

## 8.3 Root And Quotient Test

Let  $a_n$  be a sequence and suppose

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = q > 1$$

Then eventually  $\left| \frac{a_{n+1}}{a_n} \right|$  will be greater than  $q$ . which means  $a_n$  is diverging so  $\sum_{n=1}^\infty a_n$  diverges as well. If  $q < 1$  then eventually  $\left| \frac{a_{n+1}}{a_n} \right|$  will be smaller than  $q$ . That means  $a_n$  will converge to 0 and then we know that  $\sum_{n=1}^\infty (a_n)$  converges absolutely and that implies it converges in the usual sense as well. The Root test is very similar and stronger in general.

## 9 Rabbe's Test

In case the quotient test doesn't work - let  $a_n$  be a positive sequence then if

$$\lim_{n \rightarrow \infty} \left( n \left( 1 - \frac{a_{n+1}}{a_n} \right) \right) = q$$

Then for

$$\begin{cases} q > 1 & \text{the series converges} \\ q < 1 & \text{the series diverges} \\ q = 1 & \text{we must check using a better test...} \end{cases}$$

## 10 Integral Test for Series

Let  $f(x)$  be a positive monotone decreasing function on  $[1, \infty]$ . Define  $a_n = f(n)$  then:

$$\sum_{n=1}^{\infty} a_n \text{ converges} \iff \int_1^{\infty} f(x) dx \text{ converges}$$

Suppose the series converges then we get:

$$\left| \sum_{n=1}^{\infty} a_n < M \right|$$

But we also know that:

$$0 \leq \sum_{n=2}^{\infty} a_n \leq \int_1^{N+1} f(x) dx \leq \sum_{n=1}^{\infty} a_n$$

That means that the integral is increasing and bounded so it's converging. Suppose the integral was converging, to prove the series is also converging we could show similarly it's "bounded" by the integral's bound.

## 11 Cauchy Condensation Test

Let  $a_n$  be a non-increasing sequence of non-negative number.

$$\sum_{n=1}^{\infty} f(n) \leq \sum_{n=0}^{\infty} 2^n f(2^n) \leq 2 \sum_{n=1}^{\infty} f(n)$$

This is because of simple rearrangement of the numbers:

$$\begin{aligned} \sum_{n=1}^{\infty} f(n) &= f(1) + f(2) + f(3) + f(4) + f(5) + f(6) + f(7) + \cdots \\ &= f(1) + (f(2) + f(3)) + (f(4) + f(5) + f(6) + f(7)) + \cdots \\ &\leq f(1) + (f(2) + f(2)) + (f(4) + f(4) + f(4) + f(4)) + \cdots \\ &= \sum_{n=0}^{\infty} 2^n f(2^n) \\ &= f(1) + (f(2) + f(2)) + (f(4) + f(4) + f(4) + f(4)) + \cdots \\ &\leq (f(1) + f(1)) + (f(2) + f(2)) + (f(3) + f(3)) + (f(4) + f(4)) \cdots \\ &= 2 \sum_{n=1}^{\infty} f(n) \end{aligned}$$

## 12 Leibniz's Test

Let  $a_n$  be a monotone decreasing positive sequence such that

$$\lim_{n \rightarrow \infty} a_n = 0$$

Then

$$\sum_{n=1}^{\infty} (-1)^n a_n \quad \text{converges}$$

Since  $a_n$  is monotonically decreasing then we can say that

$$\begin{aligned} S_{2(m+1)} &= S_{2m} + (a_{2m+1} - a_{2m}) \geq S_{2m} \\ S_{2(m+1)+1} &= S_{2m+1} - (a_{2m+2} + a_{2m+3}) \leq S_{2m+1} \end{aligned}$$

Or in other words  $S_{2m}$  monotonically increases and  $S_{2m+1}$  monotonically decreases. But we also know that

$$S_{2m+1} - S_{2m} = a_{2m+1} \geq 0$$

And that means that

$$a_1 - a_2 = S_2 \leq S_{2m} \leq S_{2m+1} \leq S_1 = a_1$$

In other words our monotone sequences are bounded and so they converge. Recall as  $m \rightarrow \infty$

$$S_{2m+1} - S_{2m} = a_{2m+1} \rightarrow 0$$

So by Cantor's lemma  $S_{2m+1}$ ,  $S_{2m}$  converge to the same limit  $L$ . Moreover

$$S_{2m} \leq L \leq S_{2m+1}$$

And also

$$|S_k - L| \leq a_{k+1}$$



## 13 Riemann Series Theorem

Suppose that  $(a_1, a_2, a_3, \dots)$  is a sequence of real numbers, and that  $\sum_{n=1}^{\infty} a_n$  is conditionally convergent. Let  $M$  be a real number. Then there exists a permutation  $\sigma$  such that:

$$\sum_{n=1}^{\infty} a_{\sigma(n)} = M$$

This is also the case for  $M = \pm\infty$ . If  $a_n$  is absolutely converging then rearrangement of the elements is possible. If a series is converging then putting parentheses is valid. This can be shown by generating a new sequence such that each summation is an element, and showing that it converges.

### 13.1 Dini's Theorem

Let  $f_n(x) \rightarrow f(x)$  converge pointwise in  $D = [a, b]$  if  $\forall x \in D$  ( $f_n(x)$  is monotonic) and  $f, f_n$  are continuous then  $f_n(x) \rightarrow f(x)$  converges uniformly.

## 14 Properties of Uniformly Converging Function Sequences

### 14.1 Continuity

Suppose  $f_n \rightarrow f$  converges uniformly, and  $f_n$  is continuous for any  $n \in \mathbb{N}$ . Then  $f$  is continuous. The proof is based on the triangle inequality.

### 14.2 Integrability

Suppose  $f_n \rightarrow f$  converges uniformly on  $[a, b]$ , and  $f_n$  is integrable for any  $n \in \mathbb{N}$  then  $f$  is integrable and as  $n \rightarrow \infty$

$$\int_a^b \int f_n \rightarrow \int_a^b \int f$$

### 14.3 Differentiability

Suppose  $\forall n \in \mathbb{N}: f_n \in C^1$  on  $I$  such that:

- $f'_n$  uniformly converges on  $I$
- $\exists x_0 : f_n(x_0)$  converges

Then  $f_n$  uniformly converge on  $I$  to  $f$  and

$$f'_n \rightarrow f'$$

## 15 Weierstrass M-Test

Let  $\sum_{n=1}^{\infty} f_n(x)$  be a function series. Suppose exists a sequence  $M_n$  such that:

- $\forall n \in \mathbb{N} (|f_n(x)| \leq M_n)$
- $\sum_{n=1}^{\infty} M_n$  converges.

Then  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly.

### 15.1 Proof

Since  $M_n$  converges we can use an equivalent definition for the convergence of the series and so

$$\forall \varepsilon > 0: \exists N \in \mathbb{N} \text{ such that } (n > N) \wedge (p \in \mathbb{N}) \rightarrow \left( \left| \sum_{k=1}^{n+p} M_k(x) - \sum_{k=1}^n M_k(x) \right| < \varepsilon \right)$$

And since  $0 \leq M_n$  that implies

$$\sum_{k=n+1}^{n+p} M_k(x) < \varepsilon$$

And so we get that:

$$\left| \sum_{k=n+1}^{n+p} f_k(x) \right| \leq \sum_{k=n+1}^{n+p} |f_k(x)| \leq \sum_{k=n+1}^{n+p} M_n < \varepsilon$$

## 16 Power Series Theorems

Suppose  $\sum_{n=0}^{\infty} a_n x^n$  converges for some  $x_0$ , then it absolutely converges for any  $x$  such that  $|x| < |x_0|$ . Since the power series converges  $\lim_{n \rightarrow \infty} a_n x_0^n = 0$  and so the sequence is bounded and we denote that bound  $M$ .

$$0 \leq |a_n x^n| = |a_n x_0^n| \left| \frac{x^n}{x_0^n} \right| < M \left| \frac{x}{x_0} \right|^n$$

And this sequence's sum is a geometric series so it converges and so does  $\sum_{n=1}^{\infty} |a_n x^n|$ . We also know that  $|a_n x^n| < |a_n x_0^n|$  for all  $n \in \mathbb{N}$  so according to Weierstrass's M test  $\sum_{n=1}^{\infty} |a_n x^n|$  uniformly converges. Let

$$X = \{x \in \mathbb{R} : \sum_{n=1}^{\infty} a_n x^n \text{ converges}\}$$

We claim that exists  $R = \sup X$  - the radius of convergence - and that the series converges if  $|x| < |R|$  and diverges for  $|x| > |R|$ . For any  $x > R$  the series diverges by definition of  $R$ . If  $x < -R$  we know that exists  $x_1$  such that  $R < x_1 < |x|$  such that the series converges, in contradiction to  $R$ 's definition. if  $|x| < |R|$  then there exists  $x_2$  such that  $|x| < |x_2| < |R|$  for which the series converges and then it converges for  $x$  as well.

### 16.1 Some exercises

- We know the series converges uniformly for any close interval properly inside  $[-R, R]$ . If it converges uniformly on  $[0, R]$  then it is converging in  $R$  as well.
- Let a function series converge uniformly to  $f$ . Prove  $f$  is continuous on  $(a, b)$

## 17 Cauchy–Hadamard + D’alembert Theorem

Let  $\sum_{n=0}^{\infty} a_n x^n$  be a series and let  $R$  be the radius of convergence of the series - that is to say the series converges for any  $x \in (-R, R)$  then:

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}} = \frac{1}{\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|}$$

### 17.1 Proof the 1st

Let

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} &= L \\ \Rightarrow \sqrt[n]{|a_n x^n|} &= \sqrt[n]{|a_n|} |x| \rightarrow L|x| \\ \Rightarrow \begin{cases} |x| < \frac{1}{L} = R & \text{The series converges} \\ |x| > \frac{1}{L} = R & \text{The series diverges} \end{cases} \end{aligned}$$

This proves the series converges/diverges absolutely according to the root test. If it converges absolutely this implies it converges in the usual sense. Suppose it diverges - by the root test we know that if the series diverges the partial sums don’t converge to 0 so the series must diverge as well.

### 17.2 Proof the 2nd

Let

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= L \\ \Rightarrow \frac{|a_{n+1} x^{n+1}|}{|a_n x^n|} &= \frac{|a_{n+1}|}{|a_n|} |x| \rightarrow L|x| \\ \Rightarrow \begin{cases} |x| < \frac{1}{L} = R & \text{The series converges} \\ |x| > \frac{1}{L} = R & \text{The series diverges} \end{cases} \end{aligned}$$

If the series converges absolutely we can be sure yet again that it converges. If it diverges - then by the quotient test the partial sums diverge and so the series must also diverge, and the series will diverge as we claimed.

## 18 A Note on the Taylor Series

If  $f$  is smooth on  $(-R, R)$  then  $f$  can be the limit of a power series if and only if:

$$\forall x \in (-R, R)$$
$$\lim_{n \rightarrow \infty} R_n(x) = \lim_{n \rightarrow \infty} \sum_{n=N+1}^{\infty} a_n x^n = 0$$

This is because the following are equivalent:

$$f \text{ can be the limit of a power series}$$
$$\lim_{n \rightarrow \infty} S_n(x) = f(x), \forall x \in (-R, R)$$
$$\lim_{n \rightarrow \infty} R_n(x) = \lim_{n \rightarrow \infty} f(x) - S_n(x) = 0, \quad \forall x \in (-R, R)$$

## 19 Continuous Partial Derivatives imply Differentiability

### 19.1 Semi-Proof

We want to find the tangential plane to  $f$  for  $(x_0, y_0)$  assuming that the partial derivatives are continuous at that point. Let's denote

$$z_0 = f(x_0, y_0) \quad \text{and} \quad A = \frac{\partial f}{\partial x}(x_0, y_0) \quad \text{and} \quad B = \frac{\partial f}{\partial y}(x_0, y_0)$$

Now the tangential lines that intersect at  $z_0$  and are parallel to the axes (and in turn are perpendicular to one another) are (since the derivatives are continuous)

$$\begin{aligned} z &= B(y - y_0) + z_0 \\ z &= A(x - x_0) + z_0 \end{aligned}$$

Their directional vectors are in turn

$$\begin{aligned} \vec{\beta} &= (0, 1, B) \\ \vec{\alpha} &= (1, 0, A) \end{aligned}$$

And the normal vector to their spanning plane is

$$\vec{N} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & B \\ 1 & 0 & A \end{vmatrix} = (A, B, -1)$$

And so the plane equation is

$$\begin{aligned} A(x - x_0) + B(y - y_0) - (z - z_0) &= 0 \\ \Rightarrow z &= z_0 + \frac{\partial f}{\partial x}(x - x_0) + \frac{\partial f}{\partial y}(x_0, y_0)(y - y_0) \end{aligned}$$

We have shown that if continuous partial derivatives exists at  $(x_0, y_0)$  then  $f$  has a tangential plane at  $(x_0, y_0)$  which is equivalent to being differentiable at  $(x_0, y_0)$

### 19.2 Note on Differentiability

We say that  $f$  is differentiable at  $(x_0, y_0)$  if exist  $A, B$  such that

$$f(x_0 + h, y_0 + k) - f(x_0, y_0) = Ah + Bk + \alpha(h, k)\sqrt{h^2 + k^2} = Ah + Bk + \alpha(h, k)h + \beta(h, k)k$$

and  $\lim_{(h,k) \rightarrow (0,0)} \alpha(h, k) = 0$  and  $\lim_{(h,k) \rightarrow (0,0)} \beta(h, k) = 0$ . That's equivalent to

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{f(x, y) - f(x_0, y_0) - \frac{\partial f}{\partial x}(x_0, y_0)(x - x_0) - \frac{\partial f}{\partial y}(x_0, y_0)(y - y_0)}{\sqrt{(x - x_0)^2 + (y - y_0)^2}} = 0$$

## 20 Leibniz integral rule

Let  $f(x, y)$  be continuous on a rectangle  $[a, b] \times [c, d]$  and suppose  $\frac{\partial f}{\partial y}(x, y)$  exists and is continuous on  $[a, b] \times [c, d]$ . Define  $F(y) = \int_a^b f(x, y) dx$  then  $F$  is differentiable on  $[c, d]$  and

$$F'(y) = \frac{d}{dy} \int_a^b f(x, y) dx = \int_a^b \frac{\partial f}{\partial y}(x, y) dx$$

### 20.1 Lemma

Lemma: if  $f(x, y)$  is continuous on  $[a, b] \times [c, d]$  then  $F(y) = \int_a^b f(x, y) dx$  is uniformly continuous on  $[c, d]$ .

We know  $f$  is continuous on a compact space so it is uniformly continuous there:

$$\forall \varepsilon > 0 \exists \delta > 0 (\forall (p_1 = (x_1, y_1), p_2 = (x_2, y_2)) : d(p_1, p_2) < \delta \rightarrow |f(p_1) - f(p_2)| < \varepsilon)$$

Now consider  $y_1, y_2 \in [c, d]$ . such that  $d(y_1, y_2) < \delta$  we know that  $\forall x \in [a, b]$  that  $d((x, y_1), (x, y_2)) < \delta$  and then we can see that

$$\begin{aligned} |F(y_1) - F(y_2)| &= \left| \int_a^b f(x, y_1) dx - \int_a^b f(x, y_2) dx \right| = \left| \int_a^b (f(x, y_1) - f(x, y_2)) dx \right| \\ &\leq \int_a^b |(f(x, y_1) - f(x, y_2))| dx < \varepsilon(b - a) \end{aligned}$$

### 20.2 The Rule

Now denote  $G(y) = \int_a^b \frac{\partial f}{\partial y}(x, y) dx$ , by the lemma  $G$  is continuous.

$$\Delta F = F(y + \Delta y) - F(y) = \int_a^b f(x, y + \Delta y) dx - \int_a^b f(x, y) dx = \int_a^b (f(x, y + \Delta y) - f(x, y)) dx$$

We know by the Lagrange theorem that  $\exists t \in (0, 1)$  such that

$$\begin{aligned} \frac{f(x, y + \Delta y) - f(x, y)}{\Delta y} &= \frac{\partial f}{\partial y}(x, y + \Delta y) \\ \Rightarrow \int_a^b (f(x, y + \Delta y) - f(x, y)) dx &= \int_a^b \left( \frac{\partial f}{\partial y}(x, y + t\Delta y) \Delta y \right) dx \\ \Rightarrow \frac{\Delta F}{\Delta y} &= \int_a^b \left( \frac{\partial f}{\partial y}(x, y + t\Delta y) \right) dx = {}^\dagger G(y + t\Delta y) \rightarrow G(y) \end{aligned}$$



### 20.3 Generalization

Let  $f(x, y)$  be continuously differentiable on a rectangle  $[a, b] \times [c, d]$  and suppose  $\frac{\partial f}{\partial y}(x, y)$  exists and is continuous on  $[a, b] \times [c, d]$ , and  $\alpha(y), \beta(y)$  are differentiable on  $[c, d]$ . Define  $F(y) = \int_{\alpha(y)}^{\beta(y)} f(x, y) dx$  then  $F$  is differentiable on  $[c, d]$  and

$$F'(y) = \int_{\alpha(y)}^{\beta(y)} \frac{\partial f}{\partial y}(x, y) dx + f(\beta(y), y)\beta'(y) - f(\alpha(y), y)\alpha'(y)$$

Denote  $\Phi(s, t, y) = \int_s^t f(x, y) dx$  then:

$$F(y) = \Phi(\alpha(y), \beta(y), y) = \int_s^t f(x, y) dx$$

And now

$$F'(y) = \frac{\partial \Phi}{\partial s} \frac{ds}{dy} + \frac{\partial \Phi}{\partial t} \frac{dt}{dy} + \frac{\partial \Phi}{\partial y} \frac{dy}{dy}$$

So by the rule we proved earlier and the fundamental theorem

$$F'(y) = -f(\alpha(y), y)\alpha'(y) + f(\beta(y), y)\beta'(y) + \int_{\alpha(y)}^{\beta(y)} \frac{\partial f}{\partial y}(x, y) dx$$

## 21 Fubini's Theorem

Let  $f(x, y)$  be continuous on rectangle  $[a, b] \times [c, d]$  then

$$\int_a^b \left( \int_c^d f(x, y) dy \right) dx = \int_c^d \left( \int_a^b f(x, y) dx \right) dy$$

### 21.1 Proof

Denote

$$\begin{cases} \varphi(t) = \int_c^t \left( \int_a^b f(x, y) dx \right) dy \\ \Psi(t) = \int_a^b \left( \int_c^t f(x, y) dy \right) dx \end{cases}$$

Since  $f$  is continuous we know that  $F(y) = \int_a^b f(x, y) dx$  is continuous and so by the fundemetal theorem:

$$\varphi'(t) = \frac{d}{dt} \int_c^t F(y) dy = F(t) = \int_a^b f(x, t) dx$$

Denote  $G(x, t) = \int_c^t f(x, y) dy$ . Then by the fundamental theorem we get

$$\frac{\partial G}{\partial t} = f(x, t)$$

And thus by the Leibniz Integral Rule

$$\Psi'(t) = \frac{d}{dt} \int_a^b G(x, t) dx = \int_a^b f(x, t) dx$$

We concluded that  $\varphi, \Psi$  have the same derivative. That means that

$$\varphi = \Psi + \text{const.}$$

We know that  $\varphi(c) = \Psi(c) = 0$  and so  $\text{const.} = 0$  and so

$$\varphi = \Psi$$

and specifically

$$\begin{aligned} \varphi(d) &= \Psi(d) \\ \int_a^b \left( \int_c^d f(x, y) dy \right) dx &= \int_c^d \left( \int_a^b f(x, y) dx \right) dy \end{aligned}$$

## 22 The Chain Rule

Let  $f(x, y)$  have continuous partial derivatives on domain  $D$ . Let  $x(t), y(t)$  be differentiable on Interval  $I$  such that  $\forall t \in I : (x(t), y(t)) \in D$  and denote  $F(t) = f(x(t), y(t))$  then

$$F'(t) = \frac{\partial f}{\partial x} \bigg|_{(x(t), y(t))} \frac{dx}{dt} + \frac{\partial f}{\partial y} \bigg|_{(x(t), y(t))} \frac{dy}{dt}$$

### 22.1 Proof

$$\frac{dF}{dt} = \lim_{\Delta t \rightarrow 0} \frac{F(t + \Delta t) - F(t)}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{f(x(t + \Delta t), y(t + \Delta t)) - f(x(t), y(t))}{\Delta t}$$

Denote

$$\begin{aligned} & \begin{cases} \Delta x = x(t + \Delta t) - x(t) \\ \Delta y = y(t + \Delta t) - y(t) \end{cases} \\ & = \lim_{\Delta t \rightarrow 0} \frac{f(x(t) + \Delta x, y(t) + \Delta y) - f(x(t), y(t))}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{\Delta f}{\Delta t} \end{aligned}$$

Since  $f$  is differentiable

$$\Delta f = \frac{\partial f}{\partial x} \Delta x + \frac{\partial f}{\partial y} \Delta y + \alpha(\Delta x, \Delta y) \Delta x + \beta(\Delta x, \Delta y) \Delta y$$

Where  $\alpha, \beta \rightarrow 0$  So:

$$\begin{aligned} F'(t) &= \lim_{\Delta t \rightarrow 0} \frac{\Delta f}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{\frac{\partial f}{\partial x} \Delta x + \frac{\partial f}{\partial y} \Delta y + \alpha(\Delta x, \Delta y) \Delta x + \beta(\Delta x, \Delta y) \Delta y}{\Delta t} \\ &= \frac{\partial f}{\partial x} \bigg|_{(x(t), y(t))} \frac{dx}{dt} + \frac{\partial f}{\partial y} \bigg|_{(x(t), y(t))} \frac{dy}{dt} \end{aligned}$$

### 22.2 Corollary

suppose  $F(u, v) = f(x(u, v), y(u, v))$  then we see that

$$\begin{aligned} \frac{\partial f}{\partial u} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u} \\ \frac{\partial f}{\partial v} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v} \end{aligned}$$

## 23 Substitution For Multiple Variables

Let  $f$  be integrable over Domain  $D$ . Let  $x(u, v)$  and  $y(u, v)$  be in  $C_1^\dagger$  and let them define an invertible transformation  $\varphi : D \rightarrow E$  where  $D$  is defined on an  $xy$  plane and  $E$  on an  $uv$  plane. Now suppose

$$\mathbb{J} = \frac{\partial(x, y)}{\partial(u, v)} = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} \neq 0 \quad \forall (u, v) \in E$$

\*It can be equal to 0 in the domain if the measure of the set of those points is 0.

Then

$$\iint_D f(x, y) \, dx \, dy = \iint_E f(x(u, v), y(u, v)) |\mathbb{J}| \, du \, dv$$

---

<sup>†</sup>continuously differentiable

## 24 Calculate $\iint_{-\infty}^{\infty} e^{-x^2}$

$$\left(\int_{-\infty}^{\infty} e^{-x^2} dx\right)^2 = \left(\int_{-\infty}^{\infty} e^{-x^2} dx\right) \left(\int_{-\infty}^{\infty} e^{-y^2} dy\right) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)} dx dy$$

Now consider the integral in polar coordinates.

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)} dx dy &= \int_0^{\infty} \int_0^{2\pi} e^{-r^2} r^\dagger d\theta dr = \int_0^{\infty} \int_0^{2\pi} r e^{-r^2} d\theta dr \\ &= \int_0^{\infty} \theta r e^{-r^2} \Big|_{\theta=0}^{\theta=2\pi} dr = 2\pi \int_0^{\infty} r e^{-r^2} dr \\ &= 2\pi \int_0^{\infty} r e^{-r^2} dr \end{aligned}$$

And

$$\begin{aligned} 2\pi \int_0^{\infty} r e^{-r^2} dr &= 2\pi \lim_{M \rightarrow \infty} -\frac{1}{2} e^{-r^2} \Big|_0^M \\ &= 2\pi \lim_{M \rightarrow \infty} -\frac{1}{2} (e^{-M^2} - e^{-0^2}) = 2\pi \left(-\frac{1}{2}(0 - 1)\right) = \pi \\ &\Rightarrow \left(\int_{-\infty}^{\infty} e^{-x^2} dx\right)^2 = \pi \\ &\Rightarrow \int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi} \end{aligned}$$