



Design and Analysis of Algorithms

Divide-and-Conquer

Si Wu

School of CSE, SCUT

cswusi@scut.edu.cn

TA: Wenhao Wu (1565865638@qq.com)

Yi Liu (1337545838@qq.com)



Topics

- **Counting Inversions**
- **Matrix Multiplication**
- **Randomized Quick-Sort**



Counting Inversions

Music site tries to match your song preferences with others.

- You rank n songs.
- Music site consults database to find people with similar tastes.

Similarity metric: number of inversions between two rankings.

- My rank: $1, 2, \dots, n$.
- Your rank: a_1, a_2, \dots, a_n .
- Songs i and j are inverted if $i < j$, but $a_i > a_j$.

	A	B	C	D	E
me	1	2	3	4	5
you	1	3	4	2	5

2 inversions: 3-2, 4-2

Brute force: check all $\Theta(n^2)$ pairs.



Counting Inversions: divide-and-conquer

- Divide: separate list into two halves A and B.
- Conquer: recursively count inversions in each list.
- Combine: count inversions (a, b) with $a \in A$ and $b \in B$.
- Return sum of three counts.

input

1	5	4	8	10	2	6	9	3	7
---	---	---	---	----	---	---	---	---	---

count inversions in left half A

1	5	4	8	10
---	---	---	---	----

5-4

count inversions in right half B

2	6	9	3	7
---	---	---	---	---

6-3 9-3 9-7

count inversions (a, b) with $a \in A$ and $b \in B$

1	5	4	8	10
---	---	---	---	----

2	6	9	3	7
---	---	---	---	---

4-2 4-3 5-2 5-3 8-2 8-3 8-6 8-7 10-2 10-3 10-6 10-7 10-9

output $1 + 3 + 13 = 17$



Counting Inversions: how to combine two sub-problems?

Q. How to count inversions (a, b) with $a \in A$ and $b \in B$?

A. Easy if A and B are sorted!

Algorithm:

- Sort A and B.
- For each element $b \in B$,
 - Binary search in A to find the elements in A greater than b.

list A

7	10	18	3	14
---	----	----	---	----

list B

20	23	2	11	16
----	----	---	----	----

sort A

3	7	10	14	18
---	---	----	----	----

sort B

2	11	16	20	23
---	----	----	----	----

binary search to count inversions (a, b) with $a \in A$ and $b \in B$

3	7	10	14	18
---	---	----	----	----

2	11	16	20	23
---	----	----	----	----

5 2 1 0 0



Counting Inversions: how to combine two sub-problems?

Count inversions (a, b) with $a \in A$ and $b \in B$, assuming A and B are sorted.

- Scan A and B from left to right.
- Compare a_i and b_j .
- If $a_i < b_j$, then a_i is not inverted with any element left in B .
- If $a_i > b_j$, then b_j is inverted with every element left in A .
- Append smaller element to sorted list C .

count inversions (a, b) with $a \in A$ and $b \in B$

3	7	10	a_i	18
---	---	----	-------	----



2	11	b_j	20	23
---	----	-------	----	----

5

2



merge to form sorted list C

2	3	7	10	11					
---	---	---	----	----	--	--	--	--	--





Counting Inversions: Merge-and-Count

Merge-and-Count(A, B)

Maintain a *Current* pointer into each list, initialized to point to the front elements

Maintain a variable *Count* for the number of inversions, initialized to 0

While both lists are nonempty:

Let a_i and b_j be the elements pointed to by the *Current* pointer

Append the smaller of these two to the output list

If b_j is the smaller element then

Increment *Count* by the number of elements remaining in A

Endif

Advance the *Current* pointer in the list from which the smaller element was selected.

EndWhile

Once one list is empty, append the remainder of the other list to the output

Return *Count* and the merged list



Counting Inversions: algorithm implementation

Input. List L.

Output. Number of inversions in L and L in sorted order.

Sort-and-Count(L)

If (list L has one element)

Return (0, L).

Divide the list into two halves A and B.

$(r_A, A) \leftarrow \text{Sort-and-Count}(A).$

$(r_B, B) \leftarrow \text{Sort-and-Count}(B).$

$(r_{AB}, L) \leftarrow \text{Merge-and-Count}(A, B).$

Return $(r_A + r_B + r_{AB}, L).$



Counting Inversions: algorithm analysis

Proposition. The Sort-and-Count algorithm counts the number of inversions in a permutation of size n in $O(n \log n)$ time.

Pf. The worst-case running time $T(n)$ satisfies the recurrence:

$$T(n) = \begin{cases} \Theta(1), & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + T(\lfloor n/2 \rfloor) + \Theta(n), & \text{otherwise} \end{cases}$$



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----

sorted list B

2	11	16	20	23
---	----	----	----	----



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----



compare minimum entry in each list: copy 2 and add x to inversion count

sorted list C

--	--	--	--	--	--	--	--	--	--



$x = 5$ ← number of elements remaining in A
inversions = 0



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5



compare minimum entry in each list: copy 3 and decrement x

sorted list C

2									
---	--	--	--	--	--	--	--	--	--



$x = 5$

inversions = 5



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5



compare minimum entry in each list: copy 7 and decrement x

sorted list C

2	3								
---	---	--	--	--	--	--	--	--	--



$x = 4$

inversions = 5



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5



compare minimum entry in each list: copy 10 and decrement x

sorted list C

2	3	7							
---	---	---	--	--	--	--	--	--	--



x = 3

inversions = 5



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5



compare minimum entry in each list: copy 11 and add x to increment count

sorted list C

2	3	7	10						
---	---	---	----	--	--	--	--	--	--



$x = 2$

inversions = 5



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5

2



compare minimum entry in each list: copy 14 and decrement x

sorted list C

2	3	7	10	11					
---	---	---	----	----	--	--	--	--	--



$x = 2$

inversions = 7



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5

2



compare minimum entry in each list: copy 16 and add x to increment count

sorted list C

2	3	7	10	11	14				
---	---	---	----	----	----	--	--	--	--



$x = 1$

inversions = 7



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5

2

1



compare minimum entry in each list: copy 18 and decrement x

sorted list C

2	3	7	10	11	14	16			
---	---	---	----	----	----	----	--	--	--



$x = 1$

inversions = 8



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----

sorted list B

2	11	16	20	23
---	----	----	----	----



5

2

1



list A exhausted: copy 20

sorted list C

2	3	7	10	11	14	16	18		
---	---	---	----	----	----	----	----	--	--



$x = 0$

inversions = 8



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5

2

1

0



list A exhausted: copy 23

sorted list C

2	3	7	10	11	14	16	18	20	
---	---	---	----	----	----	----	----	----	--



$x = 0$

inversions = 8



Merge-and-Count Demo

Given two sorted lists A and B,

- Count number of inversions (a, b) with $a \in A$ and $b \in B$.
- Merge A and B into sorted list C.

sorted list A

3	7	10	14	18
---	---	----	----	----



sorted list B

2	11	16	20	23
---	----	----	----	----

5 2 1 0 0



done: return 8 inversions

sorted list C

2	3	7	10	11	14	16	18	20	23
---	---	---	----	----	----	----	----	----	----



$x = 0$

inversions = 8



Matrix Multiplication

Matrix multiplication. Given two n -by- n matrices A and B , compute $C = AB$.

$$\begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1n} \\ c_{21} & c_{22} & \cdots & c_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ c_{n1} & c_{n2} & \cdots & c_{nn} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \times \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{bmatrix}$$

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

$$\begin{bmatrix} .59 & .32 & .41 \\ .31 & .36 & .25 \\ .45 & .31 & .42 \end{bmatrix} = \begin{bmatrix} .70 & .20 & .10 \\ .30 & .60 & .10 \\ .50 & .10 & .40 \end{bmatrix} \times \begin{bmatrix} .80 & .30 & .50 \\ .10 & .40 & .10 \\ .10 & .30 & .40 \end{bmatrix}$$



Block Matrix Multiplication

$$\begin{array}{c} C_{11} \\ \swarrow \\ \left[\begin{array}{cc|cc} 152 & 158 & 164 & 170 \\ 504 & 526 & 548 & 570 \\ \hline 856 & 894 & 932 & 970 \\ 1208 & 1262 & 1316 & 1370 \end{array} \right] \\ \end{array} = \begin{array}{c} A_{11} \quad A_{12} \\ \swarrow \quad \swarrow \\ \left[\begin{array}{cc|cc} 0 & 1 & 2 & 3 \\ 4 & 5 & 6 & 7 \\ \hline 8 & 9 & 10 & 11 \\ 12 & 13 & 14 & 15 \end{array} \right] \times \left[\begin{array}{cc|cc} 16 & 17 & 18 & 19 \\ 20 & 21 & 22 & 23 \\ \hline 24 & 25 & 26 & 27 \\ 28 & 29 & 30 & 31 \end{array} \right] \\ \searrow \\ B_{21} \end{array}$$

$$C_{11} = A_{11} \times B_{11} + A_{12} \times B_{21} = \begin{bmatrix} 0 & 1 \\ 4 & 5 \end{bmatrix} \times \begin{bmatrix} 16 & 17 \\ 20 & 21 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 6 & 7 \end{bmatrix} \times \begin{bmatrix} 24 & 25 \\ 28 & 29 \end{bmatrix} = \begin{bmatrix} 152 & 158 \\ 504 & 526 \end{bmatrix}$$



Block Matrix Multiplication: Warmup

To multiply two n -by- n matrices A and B :

- Divide: partition A and B into $\frac{n}{2}$ -by- $\frac{n}{2}$ blocks.
- Conquer: multiply 8 pairs of $\frac{n}{2}$ -by- $\frac{n}{2}$ matrices, recursively.
- Combine: add appropriate products using 4 matrix additions.

$$\begin{array}{l} \text{Diagram illustrating the block matrix multiplication process.} \\ \text{Left side (Matrix Partitioning):} \\ \begin{array}{c} \text{Red arrows point from 'n-by-n matrices' to } A \text{ and } B \text{ in } C = A \times B. \\ \text{Red arrows point from 'n/2-by-n/2 matrices' to the } 2 \times 2 \text{ block matrices in the equation:} \\ \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} \end{array} \\ \text{Right side (Block Calculations):} \\ \begin{array}{c} \text{Red arrows point from '8 matrix multiplications' to the 8 products in the equations:} \\ \begin{aligned} C_{11} &= (A_{11} \times B_{11}) + (A_{12} \times B_{21}) \\ C_{12} &= (A_{11} \times B_{12}) + (A_{12} \times B_{22}) \\ C_{21} &= (A_{21} \times B_{11}) + (A_{22} \times B_{21}) \\ C_{22} &= (A_{21} \times B_{12}) + (A_{22} \times B_{22}) \end{aligned} \\ \text{Red arrows point from '4 matrix additions' to the 4 addition operations (+) in the equations.} \end{array} \end{array}$$

Running time. Apply Case 1 of the master theorem.

$$T(n) = 8T\left(\frac{n}{2}\right) + \Theta(n^2) \Rightarrow T(n) = \Theta(n^3)$$



Strassen's Trick

Key idea. Can multiply two 2-by-2 matrices via 7 scalar matrix multiplications (plus 11 additions and 7 subtractions).

scalars

$$\begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$C_{11} = P_5 + P_4 - P_2 + P_6$$

$$C_{12} = P_1 + P_2$$

$$C_{21} = P_3 + P_4$$

$$C_{22} = P_1 + P_5 - P_3 - P_7$$

$$P_1 \leftarrow A_{11} \times (B_{12} - B_{22})$$

$$P_2 \leftarrow (A_{11} + A_{12}) \times B_{22}$$

$$P_3 \leftarrow (A_{21} + A_{22}) \times B_{11}$$

$$P_4 \leftarrow A_{22} \times (B_{21} - B_{11})$$

$$P_5 \leftarrow (A_{11} + A_{22}) \times (B_{11} + B_{22})$$

$$P_6 \leftarrow (A_{12} - A_{22}) \times (B_{21} + B_{22})$$

$$P_7 \leftarrow (A_{11} - A_{21}) \times (B_{11} + B_{12})$$

7 scalar multiplications

Pf. $C_{12} = P_1 + P_2$

$$\begin{aligned} &= A_{11} \times (B_{12} - B_{22}) + (A_{11} + A_{12}) \times B_{22} \\ &= A_{11} \times B_{12} + A_{12} \times B_{22}. \quad \square \end{aligned}$$



Strassen's Trick

Key idea. Can multiply two **n-by-n** matrices via $7 \frac{n}{2}$ -by- $\frac{n}{2}$ multiplications (plus 11 additions and 7 subtractions).

$\frac{1}{2}n$ -by- $\frac{1}{2}n$ matrices

$$\begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$C_{11} = P_5 + P_4 - P_2 + P_6$$

$$C_{12} = P_1 + P_2$$

$$C_{21} = P_3 + P_4$$

$$C_{22} = P_1 + P_5 - P_3 - P_7$$

$$P_1 \leftarrow A_{11} \times (B_{12} - B_{22})$$

$$P_2 \leftarrow (A_{11} + A_{12}) \times B_{22}$$

$$P_3 \leftarrow (A_{21} + A_{22}) \times B_{11}$$

$$P_4 \leftarrow A_{22} \times (B_{21} - B_{11})$$

$$P_5 \leftarrow (A_{11} + A_{22}) \times (B_{11} + B_{22})$$

$$P_6 \leftarrow (A_{12} - A_{22}) \times (B_{21} + B_{22})$$

$$P_7 \leftarrow (A_{11} - A_{21}) \times (B_{11} + B_{12})$$

*7 matrix multiplications
(of $\frac{1}{2}n$ -by- $\frac{1}{2}n$ matrices)*

Pf. $C_{12} = P_1 + P_2$
 $= A_{11} \times (B_{12} - B_{22}) + (A_{11} + A_{12}) \times B_{22}$
 $= A_{11} \times B_{12} + A_{12} \times B_{22}.$



Strassen's Algorithm

Strassen (n, A, B)

If ($n = 1$) Return $A \times B$.

Partition A and B into $\frac{n}{2}$ -by- $\frac{n}{2}$ blocks.

$P_1 \leftarrow \text{Strassen}(n/2, A_{11}, B_{12} - B_{22})$.

$P_2 \leftarrow \text{Strassen}(n/2, A_{11} + A_{12}, B_{22})$.

$P_3 \leftarrow \text{Strassen}(n/2, A_{21} + A_{22}, B_{11})$.

$P_4 \leftarrow \text{Strassen}(n/2, A_{22}, B_{21} - B_{11})$.

$P_5 \leftarrow \text{Strassen}(n/2, A_{11} + A_{22}, B_{11} + B_{22})$.

$P_6 \leftarrow \text{Strassen}(n/2, A_{12} - A_{22}, B_{21} + B_{22})$.

$P_7 \leftarrow \text{Strassen}(n/2, A_{11} - A_{21}, B_{11} + B_{12})$.

$C_{11} = P_5 + P_4 - P_2 + P_6$.

$C_{12} = P_1 + P_1$.

$C_{21} = P_3 + P_4$.

$C_{22} = P_1 + P_5 - P_3 - P_7$.

Return C .



Analysis of Strassen's Algorithm

Theorem. Strassen's algorithm requires $O(n^{2.81})$ arithmetic operations to multiply two n -by- n matrices.

Pf.

Apply Case 1 of the master theorem to the recurrence:

$$\begin{aligned} T(n) &= 7T\left(\frac{n}{2}\right) + \Theta(n^2) \\ \Rightarrow T(n) &= \Theta(n^{\log_2 7}) \end{aligned}$$


If n is not a power of 2, could pad matrices with zeros.



Randomized Quick-Sort

3-way partition array so that:

- Pivot element p is in place.
- Smaller elements in left subarray L .
- Equal elements in middle subarray M .
- Larger elements in right subarray R .



Recur in both left and right subarrays.

Randomized-Quick-Sort (A)

if list A has zero or one element

Return.

Pick pivot $p \in A$ uniformly at random.

$(L, M, R) \leftarrow \text{Partition-3-Way}(A, p).$

Randomized-Quick-Sort (L).


Randomized-Quick-Sort (R).

3-way partitioning
can be done in-place
(using n compares)



Analysis of Randomized Quick-Sort

Proposition. The expected number of compares to Quick-Sort an array of n distinct elements is $O(n \log n)$.



Randomized-Quick-Sort (A)

if list A has zero or one element

Return.

Pick pivot $p \in A$ uniformly at random.

$(L, M, R) \leftarrow \text{Partition-3-Way}(A, p)$.

Randomized-Quick-Sort (L).


Randomized-Quick-Sort (R).



Analysis of Randomized Quick-Sort

Proposition. The expected number of compares to Quick-Sort an array of n distinct elements is $O(n \log n)$.

Pf. Consider BST representation of partitioning elements.






Analysis of Randomized Quick-Sort

Proposition. The expected number of compares to Quick-Sort an array of n distinct elements is $O(n \log n)$.

Pf. Consider BST representation of partitioning elements.

- An element is compared with only its ancestors and descendants.






Analysis of Randomized Quick-Sort

Proposition. The expected number of compares to Quick-Sort an array of n distinct elements is $O(n \log n)$.

Pf. Consider BST representation of partitioning elements.

- An element is compared with only its ancestors and descendants.






Analysis of Randomized Quick-Sort

Proposition. The expected number of compares to Quick-Sort an array of n distinct elements is $O(n \log n)$.

Pf. Consider BST representation of partitioning elements.

- An element is compared with only its ancestors and descendants.
- $\Pr[a_i \text{ and } a_j \text{ are compared}] = 2/(j-i+1)$, where $i < j$.





Analysis of Randomized Quick-Sort

Proposition. The expected number of compares to Quick-Sort an array of n distinct elements is $O(n \log n)$.

Pf. Consider BST representation of partitioning elements.

- An element is compared with only its ancestors and descendants.

- $\Pr[a_i \text{ and } a_j \text{ are compared}] = 2/(j-i+1)$, where $i < j$.

- Expected number of compares =
$$\begin{aligned} \sum_i^n \sum_{j=i+1}^n \frac{2}{j-i+1} \\ &= 2 \sum_i^n \sum_{j=2}^{n-i+1} \frac{1}{j} \\ &\leq 2n \sum_{j=1}^n \frac{1}{j} \\ &\sim 2n \int_{x=1}^n \frac{1}{x} dx = 2n \ln n \end{aligned}$$