

STAT211 Mandatory Homework 4

Yapi Donatien Achou

February 16, 2019

Contents

1	Problem 4.1	2
1.1	Part a: Invertibility	2
1.2	Part b: Linear filter π_j	2
2	Problem 4.2	3
2.1	Part a: Finding $\{\psi_j, j = 0, 1, 2\}$	3
2.2	Part b: Check causality and invertibility	4
2.3	Part c: Plot $\{\psi_j, j = 0, \dots, 50\}$	5
3	Problem 4.3	7
3.1	Part a: Finding $\{\gamma(h), h = 0, \dots, 4\}$ for ARMA(2,3)	7
3.2	Part b: Homogeneous difference equation $\phi(B)\gamma(h) = 0$	8
3.3	Part c: Plot of $\{\gamma(h), h = 0, \dots, 50\}$	8
4	Problem 4.4	10
5	All R code Code	10

1 Problem 4.1

Consider an ARMA(p,q) model

$$X_t - \sum_{k=1}^p \phi_k X_{t-k} = Z_t + \sum_{k=1}^p \theta_k Z_{t-k} \quad (1)$$

1.1 Part a: Invertibility

An ARMA(p,q) process $\{X_t\}$ is invertible if there exist constant $\{\pi_j\}$ such that

$$\sum_{j=0}^{\infty} |\pi_j| \leq \infty \quad (2)$$

and

$$Z_t = \sum_{j=0}^{\infty} \pi_j X_{t-j} \quad \text{for all } t. \quad (3)$$

In other word $\{X_t\}$ is invertible if Z_t can be written as a linear combination of X_{t-j} , $j = 0, 1, \dots, \infty, [1]$.

Invertibility is equivalent to

$$\theta(z) = 1 + \theta_1 z + \dots + \theta_q z^q \neq 0 \quad \text{for all } |z| \leq 1 \quad (4)$$

where $\theta(z)$ is the moving average polynomial.

The process X_t is invertible if and only if the zeros of the moving average polynomial $\theta(z)$ lie outside the unit circle.

1.2 Part b: Linear filter π_j

The sequence $\{\pi_j\}$ in (3) is determined by the relation

$$(1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q)(\pi_0 + \pi_1 z + \dots) = (1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p). \quad (5)$$

Multiplying the left hand side together gives

$$\begin{aligned} (1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q)(\pi_0 + \pi_1 z + \dots) &= \pi_0 + \pi_1 z + \pi_2 z^2 + \dots + \theta_1 \pi_0 z + \theta_1 \pi_1 z^2 + \dots + \theta_2 \pi_0 z^2 \\ &= \pi_0 + (\pi_1 + \theta_1 \pi_0)z + (\pi_2 + \theta_1 \pi_1 + \theta_2 \pi_0)z^2 + \dots \end{aligned}$$

and equation (5) can be rewritten as

$$\pi_0 + (\pi_1 + \theta_1 \pi_0)z + (\pi_2 + \theta_1 \pi_1 + \theta_2 \pi_0)z^2 + \dots = (1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p). \quad (6)$$

And equating the coefficients of $z^j, j = 0, 1, \dots$, we obtain

$$\begin{aligned}\pi_0 &= 1 \\ \pi_1 + \theta_1 \pi_0 &= -\phi_1 \\ \pi_2 + \theta_1 \pi_1 + \theta_2 \pi_0 &= -\phi_2 \\ &\vdots\end{aligned}$$

or equivalently

$$\pi_j + \sum_{k=1}^q \theta_k \pi_{j-k} = -\phi_j, \quad j = 0, 1, \dots \quad (7)$$

2 Problem 4.2

Consider a causal ARMA(2,3) given by

$$X_t - \sum_{k=1}^p \phi_k X_{t-k} = Z_t + \sum_{k=1}^p \theta_k Z_{t-k} \quad (8)$$

where the linear representation satisfies

$$\psi_j = \sum_{k=1}^p \phi_k \psi_{j-k} + \theta_j, \quad j \geq 0, \quad \theta_0 = 1 \quad (9)$$

2.1 Part a: Finding $\{\psi_j, j = 0, 1, 2\}$

From (9) we get

$$\begin{aligned}\psi_0 &= 1 \\ \psi_1 &= \theta_1 + \psi_0 \phi_1 = \theta_1 + \phi_1 \\ \psi_2 &= \theta_2 + \psi_1 \phi_1 + \psi_0 \phi_2 = \theta_2 + (\theta_1 + \phi_1) \phi_1 + \phi_2\end{aligned} \quad (10)$$

Expanding (9), for $p = 2$

$$\psi_j = \phi_1 \psi_{j-1} + \phi_2 \psi_{j-2} + \theta_j \quad (11)$$

or equivalently

$$\psi_{j+2} - \phi_1 \psi_{j+1} - \phi_2 \psi_j = \theta_{j+2}, \quad j = 0, 1, \dots \quad (12)$$

which is the second order difference equation with

$$\theta_j \equiv 0, \quad \text{for } j \notin [0, 3]. \quad (13)$$

The second order homogeneous difference equation is defined for $j = 2, 5, \dots$, because then the right hand side of (12) is zeros, and we have

$$\psi_{j+2} - \phi_1 \psi_{j+1} - \phi_2 \psi_j = 0, \quad j = 2, 3, \dots \quad (14)$$

2.2 Part b: Check causality and invertibility

The auto regressive polynomial $\phi(z)$ and the moving average polynomial $\theta(z)$ are given respectively by

$$\begin{aligned}\phi(z) &= 1 - \phi_1 z - \phi_2 z^2 \\ &= 1 - 1.7z + 0.9z^2\end{aligned}\tag{15}$$

$$\begin{aligned}\theta(z) &= 1 + \theta_1 z + \theta_2 z^2 + \theta_3 z^3 \\ &= 1 - 1.4z + 0.8z^2 + 0.1z^3\end{aligned}\tag{16}$$

The ARMA process is causal and invertible if the zeros of the auto regressive polynomial and the zeros of the moving average polynomial are located outside the unit circle respectively. A complex number $z = a + bi$ is located outside the unit circle if its magnitude is greater than 1, that is

$$|z| = |a + bi| = \sqrt{a^2 + b^2} > 1.$$

By solving

$$\phi(z) = 1 - 1.7z + 0.9z^2 = 0$$

we get

$$z_1 = 0.94 - 0.47i, \quad z_2 = 0.94 + 0.47i$$

The magnitude of z_1 and z_2 are

$$|z_i| = \sqrt{0.94^2 + 0.47^2} = 1.05 > 1, \quad i = 1, 2$$

Therefore we conclude that all the roots of the auto regressive polynomial are outside the unit circle, thus the ARMA(2,3) process is causal.

In the same fashion, by solving

$$\theta(z) = 1 - 1.4z + 0.8z^2 + 0.1z^3 = 0$$

we get

$$\begin{aligned}z_1 &= -9.57178 \\ z_2 &= 0.78589 + 0.65354i \\ z_3 &= 0.78589 - 0.65354i\end{aligned}$$

and

$$\begin{aligned}|z_1| &= \sqrt{(-9.57178)^2} = 9.57178 > 1 \\ |z_2| &= \sqrt{0.78589^2 + 0.65354^2} = 1.02 > 1 \\ |z_3| &= \sqrt{0.78589^2 + (-0.65354)^2} = 1.02 > 1\end{aligned}$$

and since all the roots of the moving average polynomial are located outside the unit circle, the ARMA(2,3) process is invertible

2.3 Part c: Plot $\{\psi_j, j = 0, \dots, 50\}$

Recall that

$$\psi_j = \sum_{k=1}^p \phi_k \psi_{j-k} + \theta_j, \quad j \geq 0, \quad \theta_0 = 1 \quad (17)$$

With

$$\boldsymbol{\phi} = (\phi_1, \phi_2) = (1.7, -0.9), \quad \boldsymbol{\theta} = (\theta_1, \theta_2, \theta_3) = (-1.4, 0.8, 0.1), \quad \sigma^2 = 1. \quad (18)$$

Expanding (17), for $p = 2$ and using $(\phi_1, \phi_2) = (1.7, -0.9)$ we get

$$\psi_j = 1.7\psi_{j-1} - 0.9\psi_{j-2} + \theta_j \quad (19)$$

or equivalently

$$\psi_{j+2} - 1.7\psi_{j+1} + 0.9\psi_j = \theta_{j+2}. \quad (20)$$

From part a) we know that

$$\begin{aligned} \psi_0 &= 1 \\ \psi_1 &= \theta_1 + \psi_0\phi_1 = \theta_1 + \phi_1 = -1.4 + 1.7 = 0.3 \\ \psi_2 &= \theta_2 + \psi_1\phi_1 + \psi_0\phi_2 \\ &= \theta_2 + (\theta_1 + \phi_1)\phi_1 + \phi_2 \\ &= 0.8 + (-1.4 + 1.7) \times 1.7 - 0.9 \\ &= 0.41 \end{aligned} \quad (21)$$

So we have the final difference equation

$$\psi_{j+2} - 1.7\psi_{j+1} + 0.9\psi_j = \theta_{j+2}, \quad j = 1, \dots, 50 \quad (22)$$

with initial conditions

$$\psi_0 = 1, \quad \psi_1 = 0.3, \quad \psi_2 = 0.41 \quad (23)$$

and

$$\theta_j \equiv 0, \quad \text{for } j \notin [0, 3] \quad (24)$$

```

#initialization
psi0 = 1
psi1 = 0.3
psi2 = 0.41
theta3 = 0.1
psi3 = 1.7*psi2 - 0.9*psi1 + theta3
psi <- c(psi0,psi1,psi2,psi3)
#compute the rest
for (j in 2:48)
  psi[j+2] = 1.7*psi[j+1] - 0.9*psi[j]

#plot
plot(psi)

```




Figure 1: Plot of ψ for $j = 0, 50$

3 Problem 4.3

Consider a causal ARMA(p,q) process. Then

$$\gamma(h) = \sum_{k=1}^p \phi_k \gamma(h-k) + \sigma^2 \sum_{j=0}^q \theta_{j+h} \psi_j, \quad h \geq 0 \quad (25)$$

3.1 Part a: Finding $\{\gamma(h), h = 0, \dots, 4\}$ for ARMA(2,3)

Expanding (25) for $p = 2, q = 3$, we get

$$\begin{aligned} \gamma(h) &= \sum_{k=1}^2 \phi_k \gamma(h-k) + \sigma^2 \sum_{j=0}^3 \theta_{j+h} \psi_j \\ &= \phi_1 \gamma(h-1) + \phi_2 \gamma(h-2) + \sigma^2 (\theta_h \psi_0 + \theta_{1+h} \psi_1 + \theta_{2+h} \psi_2 + \theta_{3+h} \psi_3) \end{aligned} \quad (26)$$

which can also be written as

$$\gamma(h+2) = \phi_1 \gamma(h+1) + \phi_2 \gamma(h) + \sigma^2 (\theta_{h+2} \psi_0 + \theta_{3+h} \psi_1 + \theta_{4+h} \psi_2 + \theta_{5+h} \psi_3). \quad (27)$$

From [1], page 88, equation (3.2.3) given by

$$\gamma(h) = \sigma^2 \sum_{j=0}^{\infty} \phi_j \psi_{j+|h|} \quad (28)$$

holds true for an ARMA(p,q) process. So we can compute $\gamma(0), \gamma(1)$ as

$$\gamma(0) = \sigma^2 \sum_{j=0}^3 \psi_j \psi_j \quad (29)$$

$$\gamma(1) = \sigma^2 \sum_{j=0}^3 \psi_j \psi_{j+1} \quad (30)$$

And from (27) we have

$$\begin{aligned} \gamma(0) &= 0 + \sigma^2 \sum_{j=0}^3 \psi_j \psi_j \\ \gamma(1) &= 0 + \sigma^2 \sum_{j=0}^3 \psi_j \psi_{j+1} \\ \gamma(2) &= \phi_1 \gamma(1) + \phi_2 \gamma(0) + \sigma^2 (\theta_2 \psi_0 + \theta_3 \psi_1 + \theta_4 \psi_2 + \theta_5 \psi_3) \\ \gamma(3) &= \phi_1 \gamma(2) + \phi_2 \gamma(1) + \sigma^2 (\theta_3 \psi_0 + \theta_4 \psi_1 + \theta_5 \psi_2 + \theta_6 \psi_3) \\ \gamma(4) &= \phi_1 \gamma(3) + \phi_2 \gamma(2) + \sigma^2 (\theta_4 \psi_0 + \theta_5 \psi_1 + \theta_6 \psi_2 + \theta_7 \psi_3) \end{aligned} \quad (31)$$

or in a matrix equation

$$\begin{bmatrix} \gamma(0) \\ \gamma(1) \\ \gamma(2) \\ \gamma(3) \\ \gamma(4) \end{bmatrix} = \begin{bmatrix} \phi_1 \\ \phi_2 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ \gamma(1) & \gamma(0) \\ \gamma(2) & \gamma(1) \\ \gamma(3) & \gamma(2) \end{bmatrix} + \sigma^2 \begin{bmatrix} \sum_{j=0}^3 \psi_j \psi_j \\ \sum_{j=0}^3 \psi_j \psi_{j+1} \\ \sum_{j=0}^3 \theta_{j+2} \psi_j \\ \sum_{j=0}^3 \theta_{j+3} \psi_j \\ \sum_{j=0}^3 \theta_{j+4} \psi_j \end{bmatrix} \quad (32)$$

3.2 Part b: Homogeneous difference equation $\phi(B)\gamma(h) = 0$

From part a) equation (27) we had

$$\gamma(h+2) = \phi_1 \gamma(h+1) + \phi_2 \gamma(h) + \sigma^2 (\theta_{h+2} \psi_0 + \theta_{3+h} \psi_1 + \theta_{4+h} \psi_2 + \theta_{5+h} \psi_3). \quad (33)$$

For an ARAM(p=2,q=3), for $h \geq 4$, the right hand side of (33) is 0, resulting in

$$\gamma(h+2) = \phi_1 \gamma(h+1) + \phi_2 \gamma(h) \quad (34)$$

or

$$\gamma(h+2) - \phi_1 \gamma(h+1) - \phi_2 \gamma(h) = 0 \quad (35)$$

3.3 Part c: Plot of $\{\gamma(h), h = 0, \dots, 50\}$

The parameter are given by

$$\boldsymbol{\phi} = (\phi_1, \phi_2) = (1.7, -0.9), \quad \boldsymbol{\theta} = (\theta_1, \theta_2, \theta_3) = (-1.4, 0.8, 0.1), \quad \sigma^2 = 1. \quad (36)$$

and

$$\gamma(h+2) = 1.7\gamma(h+1) - 0.9\gamma(h) \quad (37)$$

R code

```
#initialization for psi
psi0 = 1
psi1 = 0.3
psi2 = 0.41
theta3 = 0.1
psi3 = 1.7*psi2 - 0.9*psi1 + theta3
psi <- c(psi0,psi1,psi2,psi3)
#compute the rest
for (j in 2:48)
  psi[j+2] = 1.7*psi[j+1] - 0.9*psi[j]

#Initialize gamma with gamma0 and gamma1
gamma0 = 0
for (k in 1:5)
  gamma0 = gamma0 + psi[k]*psi[k]

gamma1 = 0
for (k in 1:5)
  gamma1 = gamma1 + psi[k]*psi[k+1]

gamma = c(gamma0,gamma1)
# compute the rest of the gamma's
for (k in 1:48)
  gamma[k+2] = 1.7*gamma[k+1] - 0.9*gamma[k]
print(length(gamma))

#plot
plot(gamma, col='blue')
```

We use the R function ARMAacf to compute γ with the following code

```
Rfunction<-ARMAacf( c(1.7,-0.9),c(-1.4,0.8,0.1),50)
plot(Rfunction, col='green')
```




Figure 2: Plot of γ for $j = 0, 50$. Computed in blue vs R function (green)

Figure 2 shows the computed γ versus the γ computed with the R function ARMAacf.

4 Problem 4.4

Let $\{X_t\}$ be a causal AR(2) process with white noise process $WN(0, \sigma^2)$,

$$X_t - \phi_1 X_{t-1} - \phi_2 X_{t-2} = Z_t, \quad X_t = \sum_{j=0}^{\infty} \psi_j Z_{t-j} \quad (38)$$

5 All R code Code

References

- [1] Petter J. Brockwell. Richard A. Davis *Introduction to Time Series and Forecasting*. Springer. Second edition. 2001