

Table 1. Clinical and pathological characteristics of the EPC cohort

Variable	Metastatic (n=26)	Non-metastatic (n=26)	p-value
Age at diagnosis, yr, median (IQR)	63.5 (61.2–69.0)	63.0 (61.2–67.8)	0.497
Ancestry†, n (%)			0.169
East Asian	11 (42.3%)	8 (30.8%)	
European	11 (42.3%)	17 (65.4%)	
Other	4 (15.4%)	1 (3.8%)	
PSA at diagnosis, ng/mL†, median (IQR)	7.6 (5.2–9.5)	5.8 (4.1–8.9)	0.328
Biopsy Gleason score (total)†, median (IQR)	8.0 (8.0–9.0)	8.0 (8.0–8.8)	0.155
Positive biopsy cores, %†, median (IQR)	66.7 (51.1–87.1)	50.0 (40.0–70.0)	0.017
Clinical T-stage, n (%)			0.764
T1/T2	17 (65.4%)	19 (73.1%)	
T3	9 (34.6%)	7 (26.9%)	
Surgical margin†, n (%)			1.000
NEGATIVE	15 (57.7%)	14 (53.8%)	
POSITIVE	11 (42.3%)	12 (46.2%)	
Lymph node involvement, n (%)			0.191
NEGATIVE	25 (96.2%)	21 (80.8%)	
POSITIVE	1 (3.8%)	5 (19.2%)	
Pathological T-stage, n (%)	[n=25]	[n=24]	0.396
pT2	10 (40.0%)	13 (54.2%)	
pT3	15 (60.0%)	11 (45.8%)	
Follow-up duration, months, median (IQR)	106.5 (89.8–129.1)	134.8 (111.6–143.0)	0.023
Alive at last follow-up, n (%)			<0.001
Alive	14 (53.8%)	26 (100.0%)	
Deceased	12 (46.2%)	0 (0.0%)	

† Matching criterion; p-values shown for completeness.
[n=X] after median (IQR): X patients had available data (when <26).
Fisher's exact: binary/2-category variables. Chi-square: >2 categories (* expected cell count <5 in ≥1 cell).
Abbreviations: EPC, extreme phenotype cohort; IQR, interquartile range.