

# How much Ice do You need?

Final Presentation

**Participants:**

Joyce Tan  
Yen Theng Tan

JHU AMS 2012 FALL

Last Compiled on November 27, 2012

# Outline

## Introduction

- Sponsor
- Problem Statement
- Deliverables
- Timeline

## Content

- Approach Assumptions
- Experimental Approach
- Physics-based Approach
- Results
- Analysis

## Conclusion

- Deliverables
- Advantages and Disadvantages
- Further Recommendations

## Sponsor: McDonald's Corporation

- McDonald's Corporation is the world's largest chain of hamburger fastfood restaurants, serving around 68 million customers daily in 119 countries.
- Mcdonald's primarily sells hamburgers, cheeseburgers, chicken, French fries, breakfast items, soft drinks, milkshakes and desserts.

## Sponsor: McDonald's Corporation

- In response to healthier consumer taste, the company has expanded its menu to include salads, wraps, smoothies and fruits.
- No meal is complete without a drink; and from Diet Coke to low-fat milk to fresh-brewed, hot coffee, McDonald's serves many different varieties of beverages

# Problem Statement

- Selling soft drinks is a complement to any meal that a customer purchases at McDonald's.
- However, the server is not accustomed to putting much thought in measuring the amount of ice put in the cup.
- This often results in a overly diluted, or overly cold drink for the customer. This is likely to lower overall customer satisfaction, since a drink is a significant complement to a meal.
- Thus, customers are likely to appreciate if the right amount of ice was added for optimal satisfaction.

# Problem Statement

- To further define this problem, the exogenous variables are the proportion of ice to put in a drink.
- The endogenous variable would be the resulting temperature and concentration of the drink, as we are assuming that a customer's satisfaction is affected only by the temperature and concentration of the drink.

## Deliverables - From Team to Sponsor

- A table of optimal ice proportions/ratios for each different type of soda (namely Coca Cola, Sprite, Fanta Orange, Diet Coke),
- Matlab code with complete set of documentations that resulting temperature and dilution based on specific heat capacities and ice proportions,
- Numerical experiment results reporting success rate of different ice proportions,
- Technical report and presentations summarizing the work.

# Deliverables - From Sponsor to Team

- Sufficient supply of the 4 different sodas we are concentrating on,
- Sufficient supply of cups used by McDonald's
- Computing resources,
- Timely responses to inquiries.



# Timeline

- Work Statement due date, Sep 28, 2012,
- Midterm Presentation due date, Oct 17, 2012,
- Progress Report due date, Oct 26, 2012,
- Final Presentation due date, Nov 28, 2012,
- Final Report due date, Dec 3, 2012.

Most of the experiments and coding have been done from mid-October to mid-November.

## Approach Assumptions

- Consumer's taste depends entirely on the dilution and temperature factors.
- Dilution and temperature of drink come hand-in-hand and rely entirely on the ice proportion.
- Sample group accurately represents the population's preferred combinations of temperature and dilution.
- The different time parameters which we perform the experiment is sufficient to represent the overall satisfaction the customer has with the drink.

# Approach 1: Experimental

- Experimenting with different types of soda - namely McDonald's Coca Cola, Sprite, Fanta Orange, and Diet Coke.
- By experiment, we will test out which ice proportion will yield the highest satisfaction from the test subjects.

## Approach 1: Experimental

- We will provide 3 different cups of the same soda (different ice proportions) for the test subject to drink and they will indicate their preference.
- The ice will be left in the drink for a time period of  $t$  ( $t=0.5$ mins, 2 mins, 5 mins, 30 mins). The different experiments for the time parameters will be spaced an hour apart.
- This will be repeated for 3 more days for the other 3 drinks.

## Approach 1: Experimental

- This will be a blind test and the subject will not know what ice proportions the cups A, B, C have.

Ice Proportion	A	B	C
t=0.5mins			
t=2mins			
t=5mins			
t=30mins			

**Table:** Sample form each test subject will need to fill out for each drink

## Approach 1: Experimental

- Subject will be required to rank preference of the labelled cups for each time parameter  $t$  (3 is most favorite).

Ice Proportion	A	B	C
$t=0.5\text{mins}$	3	2	1
$t=2\text{mins}$	1	3	2
$t=5\text{mins}$	2	3	1
$t=30\text{mins}$	1	2	3

Table: Example of a response

## Approach 2: Physics-based

- Utilizing the specific heat capacities of soda and ice (already found as specific values), we can calculate the different temperatures and dilution that the resulting drink will have.
- This will be used mainly as a support tool since it's just mathematical calculation, to see how much ice proportion actually affects dilution as well as resulting temperature

## Results - Experimental approach

	40%	60%	75%
t=0.5 mins	15	25	32
t=2 mins	14	24	34
t=5 mins	14	27	31
t=30 mins	18	36	18

Table: Experiment results for Coke



## Results - Experimental approach

	40%	60%	75%
t=0.5 mins	15	27	30
t=2 mins	20	19	33
t=5 mins	14	29	29
t=30 mins	17	30	25

Table: Experiment results for Sprite

## Results - Experimental approach

	40%	60%	75%
t=0.5 mins	15	23	34
t=2 mins	19	23	30
t=5 mins	18	27	27
t=30 mins	12	35	25

**Table:** Experiment results for Fanta Orange

## Results - Experimental approach

	40%	60%	75%
t=0.5 mins	15	24	33
t=2 mins	21	19	32
t=5 mins	16	24	32
t=30 mins	18	22	32

**Table:** Experiment results for Diet Coke

## Results - Physics-based approach

Volume of ice to volume of soda	Dilution	Temperature (Celsius)
1/10	0.09	16.2
1/8	0.11	14.3
1/6	0.15	11.2
1/5	0.18	8.8
1/4	0.23	5.5

**Table:** Calculated dilution and temperature for difference ice volumes

## Analysis - Experimental approach

	40%	60%	75%	p-value	significance?
t=0.5 mins	15	25	32	0.047	significant
t=2 mins	14	24	34	0.016	significant
t=5 mins	14	27	31	0.037	significant
t=30 mins	18	36	18	0.011	significant
Sum of significant rows	61	112	115		

Table: Experiment results for Coke

- 'Good' set of data, given that the data set are all considered significant by the Chi-Squared Test
- As time elapses, subjects tend to choose the cup with less ice, but not the least ice

## Analysis - Experimental approach

	40%	60%	75%	p-value	significance?
t=0.5 mins	15	27	30	0.072	not significant
t=2 mins	20	19	33	0.079	not significant
t=5 mins	14	29	29	0.044	significant
t=30 mins	17	30	25	0.011	significant
Sum of significant rows	31	59	54		

Table: Experiment results for Sprite

- The p-values for t=0.5 mins and t = 2 mins are marginally above 0.05, but is still considered insignificant
- Ignoring those row of values, we see that at t = 5 mins and t = 30 mins, there is a strong preference towards 60% and 75%

## Analysis - Experimental approach

	40%	60%	75%	p-value	significance?
t=0.5 mins	15	23	34	0.022	significant
t=2 mins	19	23	30	0.275	not significant
t=5 mins	18	27	27	0.325	not significant
t=30 mins	12	35	25	0.004	significant
Sum of significant rows	27	58	59		

**Table:** Experiment results for Fanta Orange

- P-values for  $t = 2$  mins and  $t = 5$  mins are quite significantly above our accepted significance levels
- $t = 30$  has a very low p-value, indicating a strong lack of randomness

## Analysis - Experimental approach

	40%	60%	75%	p-value	significance?
t=0.5 mins	15	24	33	0.034	significant
t=2 mins	21	19	32	0.130	not significant
t=5 mins	16	24	32	0.069	not significant
t=30 mins	18	22	32	0.115	not significant
Sum of significant rows	15	24	33		

Table: Experiment results for Diet Coke

- There is much more 'randomness' in this set of data
- Diet Coke's effect on ice/melting points?



## Analysis - Experimental approach

	40%	60%	75%
Coke	61	112	115
Sprite	31	59	54
Fanta Orange	27	58	59
Diet Coke	15	24	33
Total	134	253	261

Table: Experimental Totals

- Taking the significant sets of data into consideration, there is an overall tendency for our subjects to prefer the 60% and 75% choices

## Analysis - Physics-based approach


Volume of ice to volume of soda	Dilution	Temperature (Celsius)
1/10	0.09	16.2
1/8	0.11	14.3
1/6	0.15	11.2
1/5	0.18	8.8
1/4	0.23	5.5

**Table:** Calculated dilution and temperature for difference ice volumes

- Dilution / Temperature equilibrium

## Analysis - Physics-based approach

Graph of Dilution and Temperature against different ice volumes



## Deliverables - From Team to Sponsor

- A table of optimal ice proportions/ratios for each different type of soda (namely Coca Cola, Sprite, Fanta Orange, Diet Coke),
- Matlab code with a complete set of documentations that show the resulting temperature and dilution calculations based on specific heat capacities and ice proportions,
- Numerical experiment results (raw data) of subject's preferences,
- Technical report and presentations summarizing our work.

# Deliverables - From Sponsor to Team

- Sufficient supply of the 4 different sodas we are working with on,
- Sufficient supply of cups used by McDonald's
- Computing resources,
- Timely responses to inquiries.

# Advantages

- Comprehensive study of consumer preferences, factoring in time, as opposed to arbitrary ice-filling.
- Utilizing the specific heat capacities of soda and ice, we can calculate the different desired combinations of temperatures and dilution of the drink.
- Good foundation for further studies with larger populations and additional factors

# Disadvantages

- Different consumer tastes regarding temperature and dilution.
- Desired temperature of drink is likely to vary with location.
- Different types of Soda may have differing effects on ice and their melting points
- Physics-based calculation assumes no interference with the environment

## Further Recommendations

- Perform experiments on different days with different climates.
- Larger subject population
- Specificity in project objectives
- Split sample group based on gender and age.



# How much Ice do You need?

Final Presentation

**Participants:**

Joyce Tan  
Yen Theng Tan

JHU AMS 2012 FALL

Last Compiled on November 27, 2012