



6. EPINEPHRINE AUTO-INJECTOR

a) Indications

- (1) Moderate to severe allergic reaction with respiratory distress or mild allergic reaction with history of life-threatening allergic reaction
- (2) Pediatric patients with severe asthma

b) Adverse Effects

- (1) Tachycardia/Palpitations
- (2) Angina
- (3) Headache
- (4) Nausea/Vomiting
- (5) Dizziness
- (6) Hypertension
- (7) Nervousness/Anxiety
- (8) Tremors

c) Precautions



Unless in severe allergic reaction or severe asthma, medical consultation must be obtained before administering to pregnant, cardiac, or adult asthma patients.

d) Contraindications

None in the presence of anaphylaxis


e) Preparations

- Epinephrine Auto-injector (single or multi-dose) only
(Patient prescribed or EMS supplied)
- (1) Adult: 0.3 mg
 - (2) Pediatric: 0.15 mg



MEDICAL CONSULTATION IS REQUIRED FOR THE ADMINISTRATION OF EPINEPHRINE AUTO-INJECTOR TO ADULT ASTHMA PATIENTS.

f) Dosage

- (1) Less than 5 years of age: 0.15 mg IM in the lateral thigh via epinephrine auto-injector or manual administration 0.15 mg in 0.15 mL IM
- (2) 5 years and greater: administer 0.3 mg IM in the lateral thigh via epinephrine auto-injector or manual administration 0.5 mg in 0.5 mL IM
- (3)  Additional doses may be administered with medical consultation.