

D. ALS PHARMACOLOGY



1. ACETAMINOPHEN

a) Indications

Patients ages 2 years and above judged to be in mild to moderate discomfort (e.g., 2–5 on FACES scale)

b) Adverse Effects

Not clinically significant

c) Precautions

Administration of acetaminophen for mild to moderate pain does not eliminate the need for transport of the patient to the hospital to receive a comprehensive evaluation of the cause of the pain and appropriate definitive treatment.

d) Contraindications

- (1) Head Injury
- (2) Hypotension
- (3) Administration of acetaminophen or acetaminophen containing medications within the previous four hours



MANY COMMON COLD PREPARATIONS CONTAIN ACETAMINOPHEN.

- (4) Inability to swallow or take medications by mouth
- (5) Respiratory distress
- (6) Persistent vomiting
- (7) Known or suspected liver disease (including patients suspected of current alcohol ingestion)
- (8) Allergy to acetaminophen
- (9) Patients less than 2 years of age

e) Preparations Use Unit Dose Only

(DO NOT USE MULTIDOSE BOTTLE OF LIQUID)

Unit dose 160 mg/5 mL liquid

Unit dose 325 mg pill or tablet

f) Dosage

- (1) Less than 2 years of age: Not indicated
- (2) 2–4 years: Unit dose 160 mg/5 mL
- (3) 5–12 years: TWO unit doses of 160 mg/5 mL each for a total of 320 mg/10 mL
- (4) 13 years and above: FOUR unit doses of 160 mg/5 mL each for a total of 640 mg/20 mL OR in a form of 325 mg pill or tablet x2 for a total of 650 mg with sips of water as tolerated by the patient.