

# Interactive Features of Proteins Composing Eukaryotic Circadian Clocks

Brian R. Crane<sup>1</sup> and Michael W. Young<sup>2</sup>

<sup>1</sup>Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853; email: bc69@cornell.edu

<sup>2</sup>Laboratory of Genetics, The Rockefeller University, New York, New York 10065; email: young@rockefeller.edu

Annu. Rev. Biochem. 2014. 83:191–219

The *Annual Review of Biochemistry* is online at [biochem.annualreviews.org](http://biochem.annualreviews.org)

This article's doi:

10.1146/annurev-biochem-060713-035644

Copyright © 2014 by Annual Reviews.

All rights reserved

## Keywords

circadian rhythm, phosphorylation, glycosylation, metabolism, PAS domain, photoentrainment

## Abstract

Research into the molecular mechanisms of eukaryotic circadian clocks has proceeded at an electrifying pace. In this review, we discuss advances in our understanding of the structures of central molecular players in the timing oscillators of fungi, insects, and mammals. A series of clock protein structures demonstrate that the PAS (Per/Arnt/Sim) domain has been used with great variation to formulate the transcriptional activators and repressors of the clock. We discuss how posttranslational modifications and external cues, such as light, affect the conformation and function of core clock components. Recent breakthroughs have also revealed novel interactions among clock proteins and new partners that couple the clock to metabolic and developmental pathways. Overall, a picture of clock function has emerged wherein conserved motifs and structural platforms have been elaborated into a highly dynamic collection of interacting molecules that undergo orchestrated changes in chemical structure, conformational state, and partners.

## Contents

POSTTRANSLATIONAL	
REGULATION OF CIRCADIAN	
RHYTHMS IN ANIMAL	
CELLS .....	192
Phosphorylation and Glycosylation	
Determine Clock Protein	
Function .....	192
NOVEL PROTEIN	
PARTNERSHIPS LINK	
CIRCADIAN CLOCKS	
TO ENVIRONMENTAL	
CHANGES AND TO	
METABOLIC AND	
DEVELOPMENTAL	
PATHWAYS.....	196
Photoentrainment .....	196
Metabolic and Developmental	
Signaling.....	196
CORE OSCILLATOR	
STRUCTURAL	
CONSIDERATIONS.....	198
PAS Domains.....	199
Cofactor Binding to PAS.....	201
Structure Peripheral to PAS.....	202
Helical Repeats in TIM.....	202
The Many Faces of FRQ.....	202
FRH, the Partner of FRQ.....	203
MECHANISMS OF LIGHT	
SENSING .....	203
LOV Domain Proteins.....	203
Cryptochromes.....	205
Melanopsin.....	207

## POSTTRANSLATIONAL REGULATION OF CIRCADIAN RHYTHMS IN ANIMAL CELLS


Circadian clocks are often depicted as networks of transcription factors that regulate their own production at the level of RNA synthesis. For example, in *Drosophila*, two proteins, Period (PER) and Timeless (TIM), negatively regulate their own expression by suppressing two activators of *per* and *tim* transcription, dClock (dCLK) and Cycle (CYC)

(Figure 1) (1, 2). Nevertheless, genetic and biochemical studies of circadian clocks in both prokaryotic and eukaryotic organisms have identified posttranslational controls that are essential to the production of molecular oscillations. Posttranslational modifications can specify longevity, activity, and the subcellular localization of proteins composing circadian clocks. Genetic screens have demonstrated that evolutionarily conserved, posttranslational controls determine the period length of the circadian oscillation and provide a basis for entrainment to environmental cycles.

### Phosphorylation and Glycosylation Determine Clock Protein Function

*Doubletime* (*Dbt*), which encodes casein kinase 1 (CK1), was the first clock gene to be linked to posttranslational control of circadian rhythmicity. Discovered in *Drosophila*, *Dbt* phosphorylates the transcription factor PER (Figure 1*a*). Loss of *Dbt* (or sharply reduced DBT function) produces behavioral and molecular arrhythmicity that was associated with increased PER stability and altered PER subcellular localization (3–5). Regulation of PER stability ultimately involves Slimb, an F-box/WD40-repeat protein that interacts preferentially with DBT-phosphorylated PER. This association with Slimb moves PER to the ubiquitin/proteasome pathway for degradation (6, 7). Nevertheless, before Slimb is recruited, DBT appears to phosphorylate PER at several sites in a stepwise fashion that encompasses several hours in wild-type *Drosophila* (8). DBT is physically associated with PER and TIM (see later in this section) (Figure 1) throughout much of this multi-hour interval, and the subcellular distribution of DBT in many tissues is dictated by that of PER and TIM at times of day when the latter proteins are abundant (9). This finding suggests that PER may also rhythmically affect the activity of DBT with respect to DBT's many alternative substrates (9).

Although some phosphorylations produced by DBT have been associated with decreased PER stability, others appear to affect PER's activity as a transcriptional repressor (7, 8, 10).




**Figure 1**

Principal elements of the circadian clocks of *Drosophila* and mammals. (a) In *Drosophila*, in the absence of nuclear Period (PER, P) and Timeless (TIM, T) proteins, transcription factors dClock (dCLK, C) and Cycle (CYC, B) drive *per* and *tim* gene expression. PER and TIM proteins physically associate in the cytoplasm, and after a delay of several hours both proteins accumulate in the nucleus. PER is phosphorylated and destabilized by Doubletime (DBT, D) in the absence of TIM, slowing down cytoplasmic accumulation of PER and PER/TIM complexes. Nuclear PER and TIM interact with dCLK/CYC complexes and block further transcription of *per* and *tim*. In the presence of light, the photoreceptor Cryptochrome (Cr) binds TIM, and both proteins are rapidly degraded. Such daylight-induced loss of TIM reinforces molecular rhythmicity. DBT-dependent nuclear phosphorylations of PER eventually trigger PER turnover and resumption of the circadian cycle. Dashed arrows indicate RNA movements; solid arrows indicate movement of proteins. (b) In mammals, CLOCK (CLK) and BMAL1 (BMAL) activate *Per* and *Cry* expression. Mammalian PER proteins are phosphorylated by casein kinase (CK)1 $\delta$  and CK1 $\epsilon$  (both orthologs of *Drosophila* DBT), which affects PER function, subcellular localization, and stability. Physical association between PER and CRY promotes nuclear accumulation of both proteins (19), and nuclear PER/CRY represses *Per* and *Cry* expression by suppressing the activity of CLOCK/BMAL1. (c) In *Drosophila*, rhythmic PER/TIM interactions with dCLK/CYC produce circadian cycles of activity for hundreds of clock-controlled genes (CCGs) (66, 231). Broad CCG programs are also found in mammals and influence almost every aspect of biology (232). This regulation is achieved through direct, cycling associations between CCGs and PER and other clock proteins, by secondary cascades of such regulation (233), and by physical interactions that couple clock proteins to certain kinases, nuclear receptors, and other regulatory factors (also see Figure 3).

Because DBT-dependent PER phosphorylation influences the duration of PER function, altered DBT kinase activity should affect the period of the circadian clock. Consistent with this view, mutations that lower DBT activity increase the period length of the circadian cycle (11, 12). CK1 similarly regulates the circadian clocks of mammals, including humans, through interactions with PER orthologs (e.g., References 13–22).

In wild-type *Drosophila*, PER is stabilized in the presence of DBT by physical interaction with a second transcription factor, TIM (**Figure 1a**). This interaction between PER and TIM is also required for their nuclear accumulation and, thus, transcriptional autoregulation of *per* and *tim* gene activity (**Figure 1a**) (23, 24). In mutants that lack TIM, PER fails to accumulate in cells expressing *per* mRNA (23). Mutant PER proteins that are poor DBT substrates, however, are stable in the absence of TIM and accumulate to high levels in the cytoplasm (2, 23, 25). These mutant proteins are transported to nuclei in the presence of TIM, indicating that DBT normally acts as a brake to cytoplasmic accumulation of PER in the absence of TIM (23, 25). Because DBT suppresses accumulation of cytoplasmic PER proteins that are not bound to TIM, the timing of PER–TIM interaction and nuclear accumulation is preceded, and therefore delayed, by DBT–PER interaction (3, 9). DBT also contributes to this delay by blocking nuclear accumulation of PER proteins that are not associated with TIM (**Figure 2**) (5).

This regulation has been further studied in TIM mutants that carry an altered nuclear localization sequence (NLS) and in TIM mutants bearing another variation that impedes TIM nuclear accumulation [TIM P116L (25), TIM-NLS (26)]. Wild-type PER proteins accumulate to high levels in the cytoplasm of *tim*-NLS mutants, but the nuclear accumulation of PER and TIM-NLS proteins is severely delayed. Flies producing wild-type PER proteins and TIM-NLS or TIM P116L proteins have long-period molecular and behavioral rhythms that are consistent with the altered



**Figure 2**

Control of Period–Timeless (PER–TIM) nuclear accumulation. In *Drosophila*, PER and TIM coexpression is required for nuclear accumulation of either protein. Nuclear accumulation is further regulated by glycosylation and phosphorylation of one or both proteins. *O*-GlcNAc transferase (OGT) adds *N*-acetylglucosamine residues to PER, and *O*-GlcNAcase (OGA) removes them. Increasing the level of OGT delays the timing of nuclear accumulation of PER and TIM. Phosphorylation of PER and/or TIM by glycogen synthase kinase 3 (GSK) and casein kinase 2 $\alpha$  (CK2) promotes PER–TIM nuclear accumulation. Biochemical studies of PER in *Drosophila* and mammals indicate that glycosylation precedes phosphorylation and may block certain sites of PER phosphorylation, including sites that are targets of Doubletime/casein kinase 1 (DBT/CK1). In *Drosophila*, DBT inhibits nuclear accumulation of monomeric PER proteins (5).

accumulation profiles of PER and TIM-NLS (25, 26).

Just as PER requires TIM for nuclear accumulation, mutations that eliminate PER cause aberrant cytoplasmic accumulation of TIM proteins (**Figure 1**) (27, 28). Also, recent studies have shown that regulated translation of PER may contribute a further delay mechanism (29, 30).

In addition to delays that may reflect translational controls, or changing concentrations of cytoplasmic PER and TIM during each circadian cycle, cytoplasmic PER/TIM complexes must be modified to permit nuclear accumulation. These modifications appear to contribute to a so-called interval timer that promotes cytoplasmic retention of PER/TIM complexes for several hours following their formation. Such delays have been studied in

flies and in cultured *Drosophila* cells (S2 cells). Perhaps the clearest indication that physical interaction between PER and TIM is not sufficient to trigger nuclear accumulation comes from fluorescence resonance energy transfer (FRET) assays of PER–TIM interactions in live S2 cells (31). FRET emerges rapidly if high levels of PER-CFP and TIM-YFP are induced, suggesting rapid formation of PER/TIM complexes in S2 cells. Nevertheless, the onset of nuclear accumulation of the tagged proteins followed the appearance of FRET by several hours (31, 32). The alternative expression of mutant proteins PER-L and TIM-NLS, both of which yield long-period circadian rhythms associated with a delay in nuclear accumulation of PER and TIM in adult *Drosophila*, further separated phases of FRET and of nuclear accumulation in S2 cells. This finding suggests that the long-period mutations affect a process (or processes) that is (are) preceded by the physical association between PER and TIM (26, 31).

Three kinases in addition to DBT (CK1) may influence this interval timer in *Drosophila* (Figure 2). Studies of *Drosophila* with mutant forms of glycogen synthase kinase 3 (GSK-3), and of *Drosophila* in which the dosage of GSK-3 has been altered, indicate that both PER and TIM may be phosphorylated by this kinase in a fashion that can advance the timing of nuclear accumulation (25, 33–35). Certain GSK-3-dependent phosphorylations of PER depend on prior PER phosphorylation by an unidentified proline-directed kinase (35). Thus, a proline-dependent kinase and GSK-3 may cooperate to establish the correct phase of PER nuclear accumulation. Finally, hypomorphic mutations of casein kinase 2 $\alpha$  (CK2 $\alpha$ ) produce long-period circadian rhythms that are associated with delayed nuclear accumulation of PER and TIM (36–38). CK2 may play a related role in the mammalian circadian clock. CK2 overexpression and inhibition also shorten and lengthen circadian period, respectively, in cultured mammalian cells (39, 40).

Glycosylation (*O*-GlcNAcylation) regulates the timing of PER nuclear accumulation in *Drosophila* (Figure 2). *O*-GlcNAc transferase

(OGT) adds *N*-acetylglucosamine to serine and/or threonine residues on PER. This regulation occurs with a distinct circadian rhythm that precedes PER hyperphosphorylation by DBT (41, 42). *O*-GlcNAcylation and phosphorylation may affect the same or overlapping sites within PER. For example, DBT-mediated degradation of PER is delayed by elevating OGT levels in cultured *Drosophila* cells. OGT-mediated glycosylation may similarly delay PER phosphorylations by GSK-3 and CK2 that promote nuclear accumulation as described above. Consistent with this view, increased levels of PER glycosylation resulting from OGT overexpression delay the timing of PER nuclear accumulation and are associated with long-period circadian rhythms in adult *Drosophila*. Knockdown of OGT in adult flies produces short-period circadian rhythms that are accompanied by an advanced phase of PER nuclear accumulation (41).

OGT conditional knockout mice have short-period circadian behavioral rhythms (42). Thus, as in *Drosophila*, OGT activity appears to play an important role in setting the period length of the mammalian circadian clock. In mice, GSK-3 $\beta$  phosphorylates, and thereby activates, OGT, with a circadian rhythm (42). GSK-3 $\beta$  itself is phosphorylated on Ser9 with a robust circadian rhythm. Ser9 phosphorylation inhibits GSK-3 $\beta$  activity, which has been observed to cycle, as determined by its circadian phosphorylation of a great variety of substrates in several mouse tissues (42). One response to this regulation is rhythmic PER2 glycosylation in the mouse. In particular, *O*-GlcNAcylation of PER2 S662, S665, and S668 blocks CK1-dependent PER2 phosphorylation at these sites (42). The inability to phosphorylate human PER2 S662 has been linked to altered function of PER2 as a transcriptional repressor in a form of familial advanced sleep phase syndrome (reviewed in Reference 22).

In cultured mouse fibroblasts, levels of PER2 *O*-GlcNAcylation, and reciprocally phosphorylation, are altered in response to changing concentrations of glucose in the media; high levels of glucose promote

OGT-dependent PER2 O-GlcNAcylation (42). These observations indicate that PER2 O-GlcNAcylation may directly function as a metabolic sensor that can adjust the phase of circadian rhythms in response to the varying availability of certain nutrients.

## NOVEL PROTEIN PARTNERSHIPS LINK CIRCADIAN CLOCKS TO ENVIRONMENTAL CHANGES AND TO METABOLIC AND DEVELOPMENTAL PATHWAYS

Changing interactions among clock proteins are required to sustain cycles of gene activity. The transcription factors dClock (dCLK) and Cycle (CYC) in flies (CLOCK and BMAL1 in mammals) form a complex that activates expression of the *per* and *tim* genes. Subsequently, physical associations of PER and TIM proteins in *Drosophila* [PER and Cryptochrome (CRY) in mammals] allow accumulation, nuclear localization, and feedback repression (**Figure 1a,b**). PER/TIM and PER/CRY produce negative feedback by direct physical associations with the above activators. In mammals, CLOCK functions as a histone acetyltransferase (HAT). Thus, its ability to activate expression of genes such as *Per* and *Cry* may reflect, at least in part, its local modification of chromatin (43). Nevertheless, the focus of CLOCK's HAT activity is not limited to histones, because CLOCK acetylates its partner, BMAL1. This modification enhances recruitment of CRY so that the HAT activity is involved in both activation and repression of target gene activity (44).

## Photoentrainment

Biological clocks quickly adapt to changes in the environment. An especially potent environmental signal is light, which resets the *Drosophila* circadian clock by regulating the abundance of the TIM protein (**Figure 1a**). Early studies of the *Drosophila* clock revealed advances or delays in the phase of the circadian rhythm in response to short pulses of daylight. The amplitude and direction of the phase

shift reflected the time of day when the light pulse was delivered. We now know that these responses depend on an interaction between TIM and CRY. In flies, CRY functions as an intracellular photoreceptor related to the DNA photolyases (PLs; reviewed in more detail in the section below titled Cryptochromes and in Reference 45). All CRYs carry a flavin adenine dinucleotide (FAD) chromophore. The redox state of FAD is affected by blue light; in *Drosophila*, in the presence of blue light, CRY binds to TIM. This association recruits JETLAG (JET), an E3 ubiquitin ligase, leading to ubiquitination and rapid degradation of both TIM and CRY (46). Consequently, a pulse of light delivered in the early night delays the phase of the circadian clock by transiently depleting TIM. However, at that time of night TIM can quickly reaccumulate due to high levels of *tim* RNA. In contrast, a light pulse that occurs just prior to dawn advances the clock: TIM is depleted, but *tim* RNA levels are too low at that time of night for new TIM synthesis during the same molecular cycle (27, 28).

## Metabolic and Developmental Signaling

In addition to being responsible for programmed environmental responses, several clock proteins directly associate with proteins engaged in metabolic and developmental signaling. One such interaction involves AMP-activated protein kinase (AMPK). This kinase acts as a nutrient sensor by phosphorylating various protein substrates in response to activation by high intracellular concentrations of AMP versus ATP or by increasing levels of calcium, both of which are responses to limited availability of intracellular glucose. In mammals, CRY1 is an AMPK substrate, and phosphorylation by AMPK kinase destabilizes CRY1 (47). PER2 also appears to be a target for circadian regulation by AMPK in mammals; as reviewed elsewhere (48), PER2 is destabilized by CK1 $\epsilon$ -directed phosphorylation, and CK1 $\epsilon$  is a regulated substrate of AMPK. Both AMPK activity and subcellular localization vary with a circadian rhythm that has been linked to



the circadian expression of mRNA encoding a regulatory subunit of the kinase AMPK $\beta$ 2 (47). Therefore, nutrient availability influences the phase of the mammalian circadian clock through altered CRY1 and PER2 accumulation, which in turn affects rhythms of AMPK and nutrient sensing.


Sirtuins (SIRT1) and poly(ADP-ribose) polymerase (PARP) convey metabolic signals to the circadian clock in mammals. SIRT1 is a NAD<sup>+</sup>-dependent deacetylase that has been connected to the regulation of longevity, genome stability, and transcriptional silencing (49, 50). Schibler and colleagues (51) have found that SIRT1 accumulates with a circadian rhythm and physically associates with PER2 and CLOCK/BMAL1 complexes. PER2 is deacetylated by SIRT1, a modification of PER2 that causes its degradation. Sassone-Corsi and colleagues (52) have provided evidence that SIRT1 also deacetylates BMAL1 and histone targets of CLOCK's HAT activity.

PARP-1 is a NAD<sup>+</sup>-dependent ADP-ribosyltransferase whose activity cycles with a circadian rhythm. PARP-1 binds to and poly(ADP-ribose)-ylates CLOCK, altering its interaction with PER, CRY, and its DNA targets (53). Given that cellular NAD<sup>+</sup> levels must be coupled to metabolic activity, changing profiles of cellular NAD<sup>+</sup> should affect patterns of PER2 accumulation and loss, as well as BMAL1 activity through SIRT1, while affecting CLOCK activity through PARP-1. Both pathways can be expected to influence the phase of the circadian clock. As in the case of AMPK interactions with CRY1, circadian rhythms of the metabolic sensors SIRT1 and PARP-1 are also influenced by their own activity. Indeed, for SIRT1, this feedback may be observed at yet another level because SIRT1/CLOCK/BMAL1 complexes directly regulate transcription of *nicotinamide phosphoribosyltransferase* (*Nampt*), which encodes a rate-limiting regulator of the NAD<sup>+</sup> salvage pathway (54). Thus, SIRT1 appears to influence the circadian production of its coenzyme.

Nuclear receptors (NRs) are central regulators of reproductive function, development, metabolism, and immune responses. They are

deployed in a ligand-dependent fashion as regulators of gene expression. This large collection of ~50 receptors in mammals affects the expression of hundreds to thousands of genes in diverse tissues. More than half of these receptors accumulate with a circadian rhythm (55). Reciprocal regulation of circadian rhythms by NRs is also widespread, and certain NRs directly bind to and regulate expression of circadian clock genes, including *Per1*, *Per2*, and *Bmal1* (56). Cross talk connects AMPK signaling and NR function, given that the function of some NRs is affected by AMPK-directed phosphorylation (56).

Regulation of NR signaling is also influenced by physical interactions with circadian clock proteins (**Figure 3**). For example, the glucocorticoid receptor (GR), which is involved in stress responses, glucose homeostasis, and immune function, can physically associate with CRY1 or with CRY2 in a manner that is stimulated by the presence of glucocorticoids (57). These associations appear to alter receptor function by rhythmically repressing GR activity with respect to a subset of target genes: Cultured cells derived from *Cry*-deficient mice show elevated expression for hundreds of genes, indicating that in wild-type mice CRY interactions with GR substantially limit the number of targets induced by hormones.



**Figure 3**

Cryptochrome (CRY) and Period (PER) physically associate with certain nuclear receptors (NRs) and alter NR function. CRY1 and CRY2 interactions with glucocorticoid receptor (GR) lead to circadian rhythms of repression and activation for some GR target genes. PER2 similarly confers oscillating patterns of expression on genes regulated by peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) during adipocyte differentiation.

Because CRY proteins are recovered from promoters of GR-responsive genes following hormone administration, they appear to function as heterologous GR partners that bind to GR response elements on the affected target genes (**Figure 3**) (57). Thus, glucocorticoid signaling is tied to the circadian clock in mammals at multiple levels: Hormone levels themselves fluctuate with a circadian rhythm, receptors accumulate with a rhythm, and receptor function oscillates in response to rhythmically produced CRY1 and CRY2.


Some NRs are also regulated by physical associations with mammalian PER (mPER) proteins (**Figure 3**). For example, the NRs peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) and REV-ERB $\alpha$  physically interact with PER2 in hepatocytes to regulate *Bmal1* expression (58). PER2 appears to function at one time of day as a repressor of *Per2* expression (through PER2-BMAL1 or PER2-REV-ERB $\alpha$  protein interactions) and at other times within the circadian cycle as an activator of *Per2* expression (through PER2-PPAR $\alpha$  interactions). PER2 also regulates PPAR $\gamma$  during adipocyte differentiation (59). This control requires a specific physical interaction with PPAR $\gamma$ , impairs PPAR $\gamma$  recruitment to regulatory elements of PPAR $\gamma$  targets, and represses PPAR $\gamma$  transcriptional activity. In contrast to wild-type mice, *Per2*-deficient mice show a broad activation of adipogenic genes in their white adipose tissue, further indicating a significant role for PER2-PPAR $\gamma$  interactions in normal adipocyte development and function. *Per2* deletion alters both the circadian phase and the amplitude of rhythmically expressed PPAR $\gamma$  targets in white adipose tissue (59). PER3/PPAR $\gamma$  plays a related role in adipocyte differentiation (60).

## CORE OSCILLATOR STRUCTURAL CONSIDERATIONS

There are common structural features among the transcriptional activators and repressor proteins that compose the core oscillators

of the fly, fungal, and mammalian clocks (**Figure 4**). Plant oscillators are also composed of autoregulatory feedback loops; however, the components therein diverge considerably from those of the other systems, so we direct the reader elsewhere for a discussion about them (61, 62). A hallmark of clock oscillator proteins is the PAS (Per/Arnt/Sim) domain, whose signature sequence was first identified as a dimerization domain in *Drosophila* PER (dPER) (63). PAS domains are widespread modules of many signaling proteins in both eukaryotes and prokaryotes (64–67). They are typically composed of ~120–150 residues that fold into five antiparallel  $\beta$ -strands (**Figure 5a**). A helical cluster (E $\alpha$ -F $\alpha$ ) that often forms a cofactor binding pocket packs across one face of the  $\beta$ -sheet, and a more variable helical region formed from C- and/or N-terminal (Ncap) segments associates on the opposing side (**Figure 5a**). [A note on PAS nomenclature: Most fields alphabetically designate the secondary structural elements regardless of type, e.g., A $\beta$ -B $\beta$ -C $\alpha$ -D $\alpha$ -E $\alpha$ -F $\alpha$ -G $\beta$ -H $\beta$ -I $\beta$ ; others, as is common with PER, separate the  $\alpha$ -helices and  $\beta$ -strands into unique naming categories (**Figure 5a**).] Circadian clock proteins, such as CLK, CYC, PER, CLOCK, BMAL1, NPAS2, mPER1–3, WC-1, and WC-2, all contain PAS repeats, where a PAS-A domain and a PAS-B domain are tightly coupled in sequence and association (**Figure 4**). In all cases, the PAS-AB repeats compose only an ~300-residue segment of a much larger protein (>1,000 residues). The remaining polypeptide is composed of various DNA-binding motifs, chromatin modification domains, phosphorylation/glycosylation centers, coiled coils, poorly ordered regions, and stretches of low sequence complexity (**Figure 4**). Indeed, the repetitive stretches of PER are one of the first documented cases of intracellular protein sequences that did not match expectations for a globular structure (68, 69). The other major oscillator components of the clock system, TIM (*Drosophila*), FRQ/FRH (*Neurospora*), and CRY (*Drosophila*, mammals), belong to classes of entirely different proteins that do





**Figure 4**


Domain architectures of core circadian clock proteins. The major structural domains are as follows: PASA and PASB, Per/Arnt/Sim (PAS)-AB repeat; PTM, posttranslational modification by phosphorylation, glycosylation, and/or acetylation; ARM, armadillo helical repeats; CRY, Cryptochrome-binding; PHD, photolyase homology domain; CC, coiled-coil; PER, Period-binding; CK1, casein kinase 1-binding; FRH, FRH-binding; RecA, ATPase; KOW, RNA-binding; HTH, helix-turn-helix DNA-binding domain; HAT, histone acetyltransferase; Q, Gln-rich transcriptional activation; LOV, light-oxygen-voltage-sensing domain; GATA, zinc-finger DNA-binding; CTT, C-terminal tail. The protein size in residues is shown on the right. Other abbreviations: d, *Drosophila*; m, mammalian; TIM, Timeless; VVD, Vivid; WC, White-Collar.

not contain PAS domains, as discussed in the following sections.

## PAS Domains

Crystal structures have been determined for the isolated PAS-AB repeats of PER (70, 71), mPER1 (72), mPER2 (73), mPER3 (72), and the CLOCK/BMAL1 heterodimer (Figure 5b) (74). All these proteins are dimeric, and although the domains themselves adhere to the standard PAS architecture, they have surprisingly different arrangements within the dimers. PER forms a head-to-tail dimer in which PAS-A interacts with PAS-B' of the opposing subunit (70, 71). A highly conserved Trp residue in the loop connecting  $\beta$ D to  $\beta$ E

of PAS-B packs near the entrance of the  $\alpha$ A- $\alpha$ C pocket that often binds cofactors (Figure 5b) (70). A PAS-B C-terminal helix, termed  $\alpha$ F, wraps back around to latch against the  $\beta$ -sheet of PAS-A. The interaction between  $\alpha$ F and PAS-A appears highly dynamic because it can associate with varied helical registers (71), and some structures show it to be completely released from the PAS-A domain (Figure 5b) (70). The  $\alpha$ F-helix directly contacts the site of the *per<sup>L</sup>* mutation (V234D), which causes a lengthened clock period (75). Residue substitution on  $\alpha$ F (M560D) disrupts PER homodimerization both in vitro (73) and in flies (76). M560D slightly lengthens the period and retards nuclear



**Figure 5**

Structures of Per/Arnt/Sim (PAS)-AB repeats in clock proteins. (a) Archetypal PAS structure with elements labeled by standard nomenclature and that used with Period (PER) proteins (in parentheses). (b) (Top) Variation in PAS-AB organization displayed by *Drosophila* PER (dPER), mammalian PER (mPER), and CLOCK/BMAL1. (Bottom) Corresponding schematics. For dPER, the  $\alpha$ F-helix wraps the PAS-AB repeat but shows bound and released conformations in different structures. The structures of mPER1 and mPER3 (not shown) are very similar to that of mPER2. The model of CLOCK/BMAL1 combines the structures of helix-turn-helix (HTH)/PAS-AB and the HTH/DNA complex (DNA is in red). The conserved Trp residues (orange) on the H $\beta$ -I $\beta$  loop have very different environments in dPER, mPER, and CLOCK/BMAL1. Protein Data Bank identifiers: dPER, 1WA9 and 3RTY; mPER2, 3GDI; CLOCK/BMAL1, 4F3L and 4H1O.

accumulation of PER (76) but does not appear to affect interactions with TIM, which instead localize along the  $\beta$ -sheet of PAS-A (73, 75). Although M560D does not affect TIM interactions, the *per*<sup>L</sup> substitution does (75); this finding is curious because both changes perturb the same homodimer interface. Thus, nuclear entry of PER, and therefore repression of CLK/CYC, may depend on both PER homodimerization and TIM binding, which are coupled in a manner that is not yet understood at the molecular level.

Unlike the head-to-tail arrangement of dPER, the PAS repeats of all three mPER proteins show a much different association mode (Figure 5b), wherein the PAS-B  $\beta$ -sheets stack in an antiparallel fashion mediated by the

Trp in loop  $\beta$ D- $\beta$ E conserved with PER (72, 73). In the mPERs, the PAS-A domains curl around in opposite directions from PAS-B to form a compact structure; the juxtaposition of PAS-A to PAS-B is more expanded in mPER1 and mPER3 compared with that in mPER2. Mutation of the conserved Trp again affects mPER dimerization both in vitro and in vivo (72). An interaction motif for NRs, LXXLL (58), is found on the  $\beta$ E-strand of PAS-A in all mPERs, although the Ncap occludes this region to various extents in the respective structures. This position on the backside of the PAS-A  $\beta$ -sheet is similar to that of the TIM-interacting residues of PER PAS-A (73).

The structure of the mammalian CLOCK/BMAL1 heterodimer displays yet another

arrangement of the PAS-AB repeat (**Figure 5b**) (74). In this case, PAS-AB associate in a head-to-head parallel contact, dictated by the necessity of the N-terminal helix-turn-helix (HTH) DNA-binding motifs recognizing the palindromic sequence of the E-box (77, 78). The PAS-A domains form a heterodimer that is similar in structure to bacterial PAS dimers such as NifL (79). Surprisingly, the PAS-B domains are not related by similar twofold symmetry; instead, they align in a back-to-front association. This alignment engages the conserved Trp of BMAL1 with the  $\alpha$ B- $\alpha$ C pocket of CLOCK (as in PER), but because CLOCK PAS-B is rotated by 180°, the CLOCK Trp is exposed and perhaps made accessible for interaction with the mPER/mCRY (mammalian CRY) repressor complex (74). A structure of the HTH heterodimer of CLOCK/BMAL1 bound to DNA reveals the expected recognition of the palindromic E-box (**Figure 5b**). Each N-terminal helix inserts into the major groove and primarily contacts 3 bp (80). However, the structure also reveals that bases +1 and -1 of the palindrome are recognized by the protein. This breaking of symmetry explains the ability of the heterodimer to distinguish orientation and to recognize noncanonical E-box sequences such as those controlling *mper2* (77).

On the basis of sequence, the *Drosophila* dCLK/CYC activation complex is expected to be quite similar to that of CLOCK/BMAL1. However, the structure of the related White-Collar Complex [WCC (81–83)] of the fungal clock is more difficult to predict. Both WC-1 and WC-2, the two transcription factors that compose the WCC, also contain a PAS-AB repeat (81, 84), but in this case the PAS repeats are N-terminal to GATA-type zinc-finger DNA-binding domains (85–87). WC-1 also contains an additional PAS domain of the light-oxygen-voltage (LOV) subtype that binds a flavin nucleotide and is responsible for light-induced gene activation by the WCC (84, 88–90). In general, the sequences of the WC-1 and WC-2 PAS domains diverge substantially from those of CLOCK or BMAL1 and in fact are closer to those of PER. However, the head-to-tail

PER association would not allow each subunit of the WCC to recognize a short segment of DNA as do the head-to-head arrangements of CLOCK/BMAL1 and CLK/CYC. Indeed, the light-responsive elements (LREs) of the fungal clock contain imperfect palindromes of the GATA recognition motif that can occur in direct repeats separated by ~17 nt (85, 89). Thus, despite a similar domain construction, the architecture of the WCC may not parallel that of the animal transcriptional activators. By analogy to PER/NR associations, WC-1 interacts with a histone acetyl transferase via LXXLL motifs that localize to the PAS-AB repeat (91).

## Cofactor Binding to PAS

Many PAS domains bind cofactors or substrates to mediate the sensing of ligands, light, or redox (64–67). Cofactor binding by PAS-containing clock proteins has also attracted attention. NPAS2, a paralog of CLOCK, was identified as a gas-sensing protein (CO and NO) that depends on heme binding for activity (92). Since this discovery, several studies have characterized the *in vitro* heme-binding properties of the PAS domains from NPAS2, mPER1–3, and CLOCK (72, 93–100). However, an examination of heme binding by mPER2 found that the interaction is largely nonspecific, with properties characteristic of proteins that are not known to have functional interactions with heme (97). *In vivo* effects of heme on the clock are complicated by the involvement of the non-PAS NRs REV-ERB $\alpha$  and REV-ERB $\beta$ , which bind heme as their ligands and regulate the expression of *Clock*, *Bmal1*, and *Npas2* (101–105). Furthermore, the structures of CLOCK and mPER1–3 would have to undergo substantial changes in conformation and oligomeric state to accommodate heme in the PAS cofactor pocket. A novel heme-binding motif may also appear in the C-terminal domain of the mPERs (106). Mutants therein affect the association of exogenously added hemin and mPER stability. Recently, investigators used a fluorescent dye that is sensitive to by-products of heme-based oxidations to detect the presence of heme

in nuclear-localized NPAS2, but not CLOCK (107). In summary, the mammalian clock receives input from heme through the REV-ERB proteins; whether additional heme signals are integrated through the core oscillator proteins is an open question.

### Structure Peripheral to PAS

Outside their PAS domains, dPER and mPER contain dissimilar regions with characteristics of largely unstructured sequences and modification centers, with the exception of an ~30-residue conserved segment that directs phosphorylation by DBT and the casein kinases (108, 109). Immediately C-terminal to the PAS-AB repeat in dPER are the so-called per-short domain and the per-short downstream domain, which contain several phosphorylation sites, one of which is the position of the *perS* mutation [Ser589 (10, 110, 111)]. Missense mutations here (and even deletions limited to ~20 amino acids encompassing this region) dramatically shorten the period length of the fly clock. A hierarchical sequence of phosphorylation in these domains may gate PER degradation by controlling the conformational state of the protein. A more open phosphorylated state may favor access to DBT and the SLIMB ubiquitin ligase, the latter of which recognizes phosphorylation sites at the N terminus (7, 8). The mCRY interaction region localizes to the C termini of the mPERs, where it is presumed to assume a helical domain (112).

In CLOCK, the HAT domain C-terminal to the PAS-AB repeat has some sequence similarity to a conserved motif in a structurally defined histone acetyl transferase [yeast ESA1 (43)]. Several residues involved in binding acetyl-CoA are conserved between CLOCK and ESA, and mutations therein affect HAT activity of CLOCK (43). However, outside of this active site region there is little homology between the two proteins, and the entire ESA1 acetyltransferase domain could not be accommodated by the CLOCK sequence (113). Thus, the overall architecture of the CLOCK HAT domain could be quite different than those of known acetyltransferases.

### Helical Repeats in TIM

TIM is also a large protein that is predicted to have low order across its length. With some differing interpretations (114–118), bioinformatic analyses have identified TIM as a helical-repeat protein that may form two major domains of armadillo (ARM) repeats (118). In this scenario, each ARM repeat comprises three helices in a roughly triangular arrangement, and subsequent repeats layer on top of each other in a so-called  $\alpha$ -helical solenoid (114). The ARM domains are separated by a low-complexity region, which also contains repetitive elements. C-terminal to the second ARM domain is another low-complexity region followed by three two-helix structural motifs similar to those found in prenyltransferases (118). TIM residues 447–914, which localize within the second predicted ARM domain, are important for binding PER (75). Given the tendency of the PER PAS domains to associate helices across their respective  $\beta$ -sheets, a TIM ARM helix may associate with PER PAS-A in a similar fashion.

### The Many Faces of FRQ

Other than PER, the best-studied clock repressor protein is probably Frequency (FRQ), a key oscillator component of the fungal clock (for reviews, see References 119–121). FRQ plays the role of PER/TIM or mPER/mCRY in *Neurospora crassa* and is associated with the frequency-interacting RNA helicase (FRH). FRQ has two major structured domains: One binds FRH, the other casein kinase 1a (CK1a, analogous to DBT). FRQ also contains a coiled-coil domain for dimerization that is essential for clock function (122). In addition, two PEST degradation sequences are centers for phosphorylation and act to coordinate proteasome-mediated degradation of FRQ (123, 124). FRQ mRNA undergoes alternative splicing to produce long and short versions of the protein (lFRQ and sFRQ) that differ by 100 residues at the N terminus (125). An increase in temperature produces more lFRQ and is correlated with temperature compensation of the clock

(126–128). FRQ has been referred to as a phosphorylation platform because it contains a vast number of sites (>100) whose phosphorylation states vary over the circadian day and correlate with activity and degradation (129). Interestingly, the pattern of phosphorylation is spatially and temporally heterogeneous; the middle of the protein is rapidly phosphorylated after synthesis, followed by the PEST sequences toward the end of the day and then the N terminus of the IFRQ toward the end of the circadian cycle (129). Rather than single sites having specific functions, a cumulative effect of phosphorylation appears to drive conformational changes in the molecule that control associations of the ordered N- and C-terminal domains and possibly other factors (130). Two short stretches of amphipathic helices (residues 319–326 and 488–495) were identified as key elements of the FRQ–CK1a interaction domain (FCD1 and FCD2). Both are required for binding CK1a and inducing phosphorylation (130). Unphosphorylated FRQ adopts a closed conformation held by electrostatic interactions between the basic N-terminal domain (residues 1–355) and the acidic remainder. Progressive phosphorylation of N-terminal regions by CK1a negates this attraction to promote an open conformation that reveals the central PEST1 sequence for degradation.

The long unstructured regions of FRQ have special properties that place unexpected constraints on their production. These segments are encoded by relatively rare codons, which slow their rate of synthesis (131). If codon usage is switched to favor abundant tRNAs, FRQ levels are increased, phosphorylation patterns change, and circadian rhythmicity is lost in constant darkness. Codon optimizations at the N terminus of FRQ alter stability and phosphorylation in ways that differ from optimizations at the middle of the protein. Nevertheless, codon optimizations in both locations increase overall FRQ protein levels. Improper codon usage may affect the rate of translation for segments that rely on the influence of other factors to control folding and the modification state (131).

## FRH, the Partner of FRQ

FRH belongs to the Dob1p/Mtr4 family of RNA helicases that function in RNA surveillance as part of the TRAMP complex (132–134); the yeast versions have been crystallographically characterized (135, 136). Mtr4 has an ATPase core that is typical of other DExH helicases but also contains a KOW domain—as found in ribosomal proteins that bind 23S ribosomal RNA. FRQ always associates with FRH, which binds to FRQ at a specific interaction domain toward the C terminus of the protein. FRH stabilizes FRQ but also interacts with *frq* mRNA and influences the stability and length of its poly-A tail (137). FRH interacts with the *Neurospora* homologs of the TRAMP complex and with exosome components (137). FRH is essential for RNA metabolism, but there are viable mutants that leave these properties intact while still causing clock dysfunction (134).


## MECHANISMS OF LIGHT SENSING

### LOV Domain Proteins

In the fungal clock, light induces *frq* transcription in a fashion that depends on the WCC (reviewed in References 124, 138, and 139). The components of the WCC, WC-1 and WC-2, associate and bind at LREs in the regulatory regions of *frq* and other clock-control genes (89, 90, 140, 141). WC-1 contains a blue light-sensitive, LOV-sensing domain of the PAS superfamily that binds a flavin cofactor (84, 85, 89, 90). The WCC binds to LREs in both the presence and absence of light, but the size of the complex increases upon light exposure (89, 140) due to oligomerization of the LOV domains (142).

The WC-1 LOV domain is very similar in sequence (80% sequence similarity over 150 residues) to a second clock-associated light sensor in *N. crassa*, Vivid (VVD) (143). VVD opposes the action of the WCC to allow light adaptation (143–145). The WCC induces *vvd* expression, and VVD subsequently inhibits the





**Figure 6**

Flavin-binding clock proteins. (a) The fungal Vivid (VVD) light sensor undergoes light-induced dimerization. In the light-activated dimer (bottom), the N-terminal latch (orange) swings out to engage Y40 with the adjacent subunit. This structural change is driven by a cysteinyl-flavin adduct formation. (b) The Cryptochrome (CRY) flavin pocket is a hot spot for interactions. (Top left) The C-terminal tail (CTT) (red) binds adjacent to the flavin in *Drosophila* CRY (dCRY). (Bottom left) The C terminus of the F-box protein Fbx13 (cyan) inserts into the empty flavin pocket of mammalian CRY2 (mCRY2), and the leucine-rich repeats (LRRs) surround the last helix of the photolyase homology domain. (Top right) Flavin binds to mCRY2 in a similar mode as dCRY; apo-mCRY1 and mCRY2 have similar structures. (Bottom right) Close-up of a flavin pocket showing superposition and juxtapositions of dCRY flavin adenine dinucleotide (FAD) (yellow); dCRY FFW motif of the CTT (red); mCRY2 FAD (magenta); and F-box C-terminal Trp residue (cyan). Protein Data Bank identifiers: VVD monomer, 2PD7; VVD dimer, 3RH8; dCRY, 4GU5; mCRY2, 4I6G; mCRY1, 4K0R; Skp1/Fbx13/mCRY2, 4I6J.

WCC by binding directly to it in a light-dependent fashion (139, 142). VVD attenuates WCC function to a degree that depends on the level of light exposure (139). Consequently, the *Neurospora* clock receives a pulse of active WCC at dawn that is quickly attenuated. Domain-replacement studies demonstrate that the core PAS domain of VVD can functionally substitute for that of WC-1 (140). LOV domain proteins are broadly represented in photosensory roles throughout biology, and their structures and photocycles are well characterized (146–149). Although many features are conserved among all LOV domain proteins, the mechanisms by which downstream effectors are stimulated differ depending on the particular class of LOV proteins. In general, absorption of blue light

by the associated flavin (which can be FAD or FMN) produces a covalent adduct between a conserved cysteine residue on E $\alpha$  and the C4a position of the isoalloxazine ring of the flavin (Figure 6a) (146–149). Structures of VVD in the dark-adapted, light-adapted adduct, as well as signaling states (150–153), show that formation of the adduct propagates to larger structural rearrangements that culminate in protein dimerization (Figure 6a) (151, 154, 155). VVD has a typical LOV/PAS structure; FAD is bound in the PAS cofactor pocket between the  $\beta$ -sheet, E $\alpha$ , and F $\alpha$ . The Ncap of VVD is composed primarily of a  $\beta$ -strand, an  $\alpha$ -helix, and an N-terminal latch that reaches back around the  $\beta$ -sheet to the adenosine moiety of FAD. As shown in structures of illuminated crystals



grown from dark-state VVD, formation of the Cys adduct effectively reduces the flavin ring and results in protonation of flavin N5. Flavin protonation alters the conformation of conserved Gln182, whose perturbed interactions propagate to the  $\beta$ -strand and hinge leading to the Ncap. The net result is a modest shift of the  $\alpha$ -helix and  $\beta$  Ncap elements. Nonetheless, the crystalline environment prevents any larger-scale changes to the structure. An investigation of factors affecting the VVD photocycle led to the recovery of variants that stabilize the flavin adduct state and alter the lifetime of the light-adapted dimer (150). Subsequent crystallization of these species revealed subunit changes similar to those observed on illumination of dark-state crystals, but in addition, the N-terminal latch completely releases and reaches across to engage another VVD subunit (**Figure 6a**) (153). A conserved Tyr residue in the N-terminal latch inserts into the same hinge between the  $\beta$ -strand and the Ncap that promulgates latch release. Mutation of this Tyr and other interface residues in the dimer structure abrogates or diminishes the biological activity of VVD in vitro and in vivo (153). Molecular dynamics simulations of the VVD adduct state on the nanosecond timescale have suggested a different activation mechanism that involves similar regions of the protein but also much larger conformational changes of Gln182 than have been observed in crystal structures (156).

Notably, most of the structural features that mediate the VVD conformational response are conserved in the WC-1 sequence. Thus, a similar light-induced dimerization mechanism is very likely to be operative and responsible for recruitment of additional WC-1 subunits to the transcriptional activation complex (153). Indeed, the LOV domains of WC-1 also undergo light-dependent dimerization and VVD interferes with this process (142, 153). These findings gave rise to the idea that VVD may exert its inhibitory effect by producing a heterodimer with the WC-1 LOV domain that resembles the VVD light-state dimer (139, 142, 151, 153, 157). Such an interaction would break up the higher-order WC-1 complex formed with light

and diminish WC-1 transcriptional activation. Attractive as such a detailed mechanism for light adaptation is, a VVD/WC-1 heterodimer remains to be structurally characterized.

## Cryptochromes

CRYs are flavin-binding proteins that closely resemble PL DNA repair enzymes (158–160). CRYs play quite different roles in the circadian clocks of mammals, insects, and plants (158–160). *Drosophila* CRY (dCRY) is the light-entrainment sensor of the fly clock, although it also has non-clock-related functions that include magnetosensitivity (161–163). Mammals contain two CRY proteins (mCRY1 and mCRY2) that instead interact with mPERs as core elements of the circadian clock and do not mediate light responses (**Figure 1b**) (160, 164–167). PLs repair cross-linked pyrimidine dimers formed in DNA by UV radiation. PLs rely on two cofactors, FAD and a second antenna cofactor that is usually a pteridine moiety [methenyl-trihydrofolate (MTHF)] or a deazaflavin [hydroxy-ribo-deazaalloxazine (HDF)]. In PLs, light energy absorbed by the antenna cofactor is relayed to reduced FAD by energy transfer. The excited FAD then injects an electron into the DNA lesion and breaks the pyrimidine linkage by a cyclic electron-transfer reaction (158, 168). Although at least some CRYs bind FAD similar to PLs, the PL DNA repair mechanism is not thought to participate in CRY functions within the clock (158–160, 167).

The CRY/PL family can be grouped into several subfamilies on the basis of sequence and function: (a) cyclobutane pyrimidine dimer (CPD) PLs (classes I and II); (b) 6-4 PLs; (c) CRY-DASH proteins, which are single-stranded DNA PLs; (d) plant CRYs; and (e) animal CRYs (158, 169–171). All family members are composed of a common structural fold consisting of an N-terminal  $\alpha/\beta$  nucleotide-binding domain connected to a C-terminal  $\alpha$ -helical domain through a flexible linker. Animal CRYs can be further subdivided by sequence and function into type I, as found

in *Drosophila*, and type II, as found in mammals (172–176) and other insects (174). Different CRYs are distinguished by C-terminal extensions (CCEs) that vary considerably in sequence and length (158, 169–171).

As discussed above, dCRY regulates circadian function by targeting TIM for ubiquitin-mediated degradation in the presence of light through action of the E3 ubiquitin ligase JET (177–179). Light also causes dCRY degradation (177–180) through action of BRWD3 E3 ubiquitin ligase (181). dCRY engages TIM in the presence of light. Binding is necessary but not sufficient to degrade TIM (180, 182). The light-dependent recognition of TIM by dCRY involves changes at the unique CCE of dCRY (180). For example, dCRY with the 19 C-terminal residues removed (CRY<sub>M</sub>) binds TIM both in the dark and in the light and greatly reduces clock entrainment under normal levels of light (180). In vertebrates, CRY enters the nucleus and interacts directly with CLOCK/BMAL1 (164, 183–185). The mCRYs are targets of the E3 ubiquitin ligase complex SCF<sup>Fbxl3</sup>; the F-box protein Fbxl3 directly recognizes and marks CRY for degradation (186–188). AMPK phosphorylates mCRY1 to promote binding of Fbxl3 (47), whereas a homologous F-box protein can compete with Fbxl3 to prevent degradation and effectively lengthen clock period (188).

The structure of full-length dCRY reveals a typical PL-type fold in which the  $\alpha/\beta$  domain is devoid of cofactor and the  $\alpha$ -helical domain binds FAD in a deep pocket (**Figure 6b**) (189–191). Although there is some evidence that dCRY binds MTHF (192), key residues involved in this interaction are not conserved with PLs, and the structure of the antenna region differs considerably in CRYs compared with PLs. Currently, there is no known biological function for a CRY antenna cofactor, and no known structures contain antenna cofactors (189, 192, 193).

C-terminal to the photolyase homology domain (PHD), a small loop connects to the so-called C-terminal tail of dCRY (CTT) (residues 528–539). The CTT forms a helix containing

a conserved FFW motif that occupies a groove analogous to that where damaged DNA substrates bind in PLs (189–191). The sequence register of the CTT with respect to the PL-homologous groove in the first reported dCRY structure (189) was subsequently revised to yield an improved model (191). This update recognized a shift by two residues of the CTT sequence register within the PL-homologous groove (191). The revised structure was confirmed by an independent study of a dCRY variant that was slightly altered in the phosphate loop that helps sequester the flavin pocket (190). Nonetheless, in all cases the CTT is positioned to respond to light-induced redox changes at the FAD and appears to fulfill the role of a classic autoinhibitory motif (**Figure 6b**). Indeed, proteolytic protection assays indicate that the CTT becomes more exposed upon light exposure (190, 194).

The PHD structures of mCRY1 and mCRY2 closely resemble each other and are also quite similar to that of dCRY (**Figure 6b**) (190, 195). Differences include the phosphate-binding loops and CCEs, which are largely absent in the current mCRY structures. mCRYs mostly purify without FAD (160), although there are exceptions (196). Nonetheless, FAD binds to mCRY2 after purification, and the flavin-binding mode is analogous to that observed in dCRY (195). In the structure of the ternary complex of mCRY2 with its E3 ubiquitin ligase Fbxl3/Skp1, the highly conserved C-terminal projection of Fbxl3 inserts into what would be the flavin binding pocket of mCRY2 (**Figure 6b**) (195). The F-box protein then wraps 12 leucine-rich repeats (LRRs) around the  $\alpha$ -helical domain of mCRY2 to completely surround the C-terminal helix of the PHD. The flavin binding pocket of mCRY has also been revealed as the target for a set of synthetic carbazole compounds that compete with the E3 ligase and thereby inhibit mCRY degradation (197). These interactions were found during a small-molecule library screen for altered expression of a *Bmal1* reporter construct in human osteosarcoma cells. The crystallographic structure of mCRY2 in

complex with such a compound shows how it mimics interactions of both FAD and the C terminus of Fbxl3 with the mCRY pocket (198).

Thus, E3-ligases, autoinhibitory motifs, cofactors (e.g., FAD), and small molecules all compete for the analogous “receptor” pocket in the  $\alpha$ -helical domains of CRYs (**Figure 6b**). The competition among these interactions, which is modulated by flavin redox state (itself influenced by light), is probably the core property underlying CRY functionality. Nonetheless, there is still much to be learned about CRY interactions and the biological role of flavin. In mCRY, the C-terminal helix of the  $\alpha$ -helical domain bound by the LRRs of the F-box has also been predicted to form part of a coiled-coil domain. This presumed coiled coil, when contained in an isolated CCE, interacts with C-terminal fragments of BMAL1 (185). mCRY binds to multiple locations on BMAL1 (183–185, 190, 199) and with PAS domains of CLOCK (199). Indeed, the conserved Trp residue on PAS-B of CLOCK that has turned out to the solvent (see the previous section) may be recognized by mCRY (74). Yeast-2 hybrid studies of zebrafish CRY1 (zCRY) indicate that interactions with CLOCK and BMAL1 prevent their ability to dimerize (199). Another study found that zCRY binds to intact CLOCK/BMAL1 on DNA through its PHD but requires reductants for this activity (183, 184). The CCEs of mCRY also mediate interactions with the C terminus of mPER (200).

Regarding the mechanism of light sensing, the chemical states of dCRY that participate in the photoswitch remain unclear; both the anionic semiquinone and a further photoexcited state have been suggested to be the forms that elicit downstream signals (192, 194, 201–204). Photoreduction or chemical reduction to the anionic semiquinone correlates with conformational changes at the CTT that allow for binding to regions of TIM (203); however, dCRY will still promote TIM degradation when an internal Trp residue important for flavin photoreduction in vitro is removed (204). In the case of mCRY, as yet there is no strong evidence for flavin function in vivo, although structural and

biochemical evidence is consistent with flavin binding (167, 195, 196). Remarkably, a CPD PL from *Potorous* can rescue clock function in a mouse mCRY1/2 knockout strain (205). The CPD PL contains no CCE, so the requirement for these regions may stem from their ability to influence properties of the PHD, which must itself have properties that are capable of supporting the clock. Perhaps we have come full circle in generalizing the in vivo activities of CRY and PL.

## Melanopsin

Although investigators initially thought that CRY proteins might be responsible for light entrainment of the mammalian clock (167), research over the past decade has led to the realization that the principal light sensors are opsins that localize to specialized neurons that lie below the rods and cones of the eye (for reviews, see References 206–210). These cells, termed intrinsically photosensitive retinal ganglion cells (ipRGCs), project directly into the suprachiasmatic nucleus of the hypothalamus and, on their own, provide information about environmental light to this key coordinator of circadian rhythmicity within the brain (211–213). Although ipRGCs function independently of retinal rods and cones, they receive input from the major light-sensing cells. Thus, various light sensors can convey complex photoentrainment signals to the mammalian pacemaker. Photosensing by ipRGCs is distinct from spatial vision and has appropriate properties for monitoring background light-intensity levels, such as high thresholds for light responses, slow temporal responses, a large dynamic range, and enhanced sensitivity to blue wavelengths (212).

Photosensitivity of ipRGCs derives from a unique opsin molecule, melanopsin, that has higher sequence identity with opsins from cephalopods and arthropods (39%) than with vertebrate ocular rhodopsins (27%). Melanopsin was originally identified in the dermal melanophores of photosensitive frog skin (214); orthologs in mouse and human genomes

were then found to localize to ipRGCs by *in situ* hybridization techniques (212, 213, 215). The circadian clocks of melanopsin knockout mice have altered phase-resetting responses to light pulses (165, 216, 217), whereas mice also devoid of rods and cones show no response to light at all (165, 166). There are two types of melanopsins: *Opn4x*, which is found in all vertebrates except mammals, and *Opn4m*, which is found in mammals (218). Like other opsins, melanopsin belongs to the class of seven-transmembrane  $\alpha$ -helical proteins known as G protein-coupled receptors (GPCRs). Opsins bind a retinal cofactor through a Schiff base linkage to a conserved Lys residue (219). Photon absorption causes an isomerization of the chromophore from an 11-*cis* configuration to an all-*trans* configuration. Melanopsins differ from the rhodopsins of rods and cones in several important ways (207, 209, 210): Their action spectra is blueshifted [440–480 nm versus 510–560 nm (209, 220)], they are bleach resistant due to photoregeneration of the 11-*cis* configuration, and they couple to different downstream G protein targets ( $G_{q/11}$ ) than do vertebrate rhodopsins ( $G_{i/o}$ , transducin).

Sequence comparisons with bovine rhodopsin led to early structural predictions for melanopsin (221). Although the overall sequence identity among divergent opsins is not large (~27%), a 35-residue noncontiguous motif termed the ligand pocket vector (LPV) captures many of the distinguishing characteristics of the classes (222). Conservation of the LPV is high within classes of vertebrate rhodopsins (94–100% identical for rod rhodopsins, 89–100% for *Opn4x*, and 94–100% for *Opn4m*) and correlates well with differences in functionality. Most opsins bind the retinal cofactor among the seven-transmembrane helices in the center of the protein, closer to the extracellular side of the membrane (219). Transmembrane helix 3 and extracellular loop 2 (between helices 4 and 5) sandwich the cofactor. Nonetheless, distinguishing features of melanopsin include alternation of the Schiff base counterion from a Glu to a Tyr (contained on helix 3); charge compensation for the Schiff base by another Glu in extracellular

loop 2; an insertion in the third cytoplasmic loop, lengthening of the N and C termini; and alteration of various residues in helices 1–4, increasing the hydrophobic nature of the retinal binding site (221).

More recently, insights into the structure of melanopsin have come from structural studies of a squid rhodopsin [33% identical with melanopsin (223)], in which several distinguishing features of melanopsin are conserved, particularly in the chromophore binding pocket. Like melanopsin, the Schiff base environment of squid rhodopsin contains a Tyr hydroxyl group and an amide side chain supplied by a helix-2 residue. Also like melanopsin, the Schiff base counterion is supplied by a Glu residue on extracellular loop 2. The Glu carboxylate interacts with the chromophore indirectly through the aforementioned amide side chain. The increased hydrophobicity of the retinal binding pocket relaxes the retinal polyene-chain conformation and allows for tighter contact of the chromophore with helix 3 than in vertebrate rhodopsin. Such changes may contribute to the stability of the retinal all-*trans* form. Computational modeling studies of melanopsin based on the squid rhodopsin structure (220) indicate that the Glu counterion forms a hydrogen bonding network with the helix-3 Tyr, the helix-2 amide (Gln), and two water molecules in the vicinity of the Schiff base. This model also suggests that removal of bulky side chains in helices 3 and 7 allows for a more planar chromophore than even that found in squid rhodopsin. Finally, a key substitution of the helix-2 amide from an Asn to a Gln in melanopsin may alter hydrogen bonding around the Schiff base to reduce polarization of the cofactor. These hydrogen bonding perturbations combine with steric effects on the retinal to contract the polyene double bonds and ultimately blueshift the absorbance maximum of the chromophore.

For bovine rhodopsin, structures of intermediate states have helped reveal the mechanism of activation (219, 224). In the structure of metarhodopsin II, isomerization of retinal to the all-*trans* form tilts the  $\beta$ -ionone group of

the chromophore toward helices 5 and 6 (225–227). This tilting bends helix 6 at a conserved Pro residue, extends helix 5, and opens a pocket between these helices and the protein core on the cytoplasmic face. This opened pocket accommodates a peptide derived from transducin in which the C-terminal end of helix 5 binds surrounded by six of seven helices (227). The structure of the  $\beta_2$ -adrenergic receptor in complex with an intact heterotrimeric G protein reveals a similar interaction between helix  $\alpha 5$  of the G protein and a cytoplasmic pocket created in part by helices 5 and 6 of the GPCR (228).

In the  $\beta_2$ -adrenergic receptor, recognition of helix  $\alpha 5$  by the GPCR dramatically alters the conformation of the  $G_s\alpha$ -subunit, whose two domains crack open to expose the nucleotide binding pocket and facilitate exchange of GDP (228, 229). Studies of the initial stages of squid rhodopsin activation indicate that an early all-*trans* intermediate has a structure similar to that of vertebrate rhodopsin (230). Nonetheless, the later, larger-scale conformational changes necessary for G protein activation remain to be resolved for an invertebrate rhodopsin and certainly for the related melanopsin.

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

Our work was supported by the following grants from the National Institutes of Health: NS053087 and GM054339 (to M.W.Y.) and GM079679 (to B.R.C.).

## LITERATURE CITED

1. Allada R, Chung BY. 2010. Circadian organization of behavior and physiology in *Drosophila*. *Annu. Rev. Physiol.* 72:605–24
2. Zheng X, Sehgal A. 2012. Speed control: cogs and gears that drive the circadian clock. *Trends Neurosci.* 35:574–85
3. Price JL, Blau J, Rothenfluh A, Abodeely M, Kloss B, Young MW. 1998. *Double-time* is a novel *Drosophila* clock gene that regulates PERIOD protein accumulation. *Cell* 94:83–95
4. Kloss B, Price JL, Saez L, Blau J, Rothenfluh A, et al. 1998. The *Drosophila* clock gene *Double-time* encodes a protein closely related to human casein kinase I $\epsilon$ . *Cell* 94:97–107
5. Cyran SA, Yiannoulos G, Buchsbaum AM, Saez L, Young MW, Blau J. 2005. The Double-time protein kinase regulates the subcellular localization of the *Drosophila* clock protein Period. *J. Neurosci.* 25:5430–37
6. Ko HW, Jiang J, Edery I. 2002. Role for Slimb in the degradation of *Drosophila* Period protein phosphorylated by Doubletime. *Nature* 420:673–78
7. Chiu JC, Vanselow JT, Kramer A, Edery I. 2008. The phospho-occupancy of an atypical SLIMB binding site on PERIOD that is phosphorylated by DOUBLETIME controls the pace of the clock. *Genes Dev.* 22:1758–72
8. Chiu JC, Ko HW, Edery I. 2011. NEMO/NLK phosphorylates PERIOD to initiate a time-delay phosphorylation circuit that sets circadian clock speed. *Cell* 145:357–70
9. Kloss B, Rothenfluh A, Young MW, Saez L. 2001. Phosphorylation of period is influenced by cycling physical associations of Double-time, Period, and Timeless in the *Drosophila* clock. *Neuron* 30:699–706
10. Kivimäe S, Saez L, Young MW. 2008. Activating PER repressor through a DBT-directed phosphorylation switch. *PLoS Biol.* 6:e183
11. Muskus MJ, Preuss F, Fan JY, Bjes ES, Price JL. 2007. *Drosophila* DBT lacking protein kinase activity produces long-period and arrhythmic circadian behavioral and molecular rhythms. *Mol. Cell Biol.* 27:8049–64



12. Syed S, Saez L, Young MW. 2011. Kinetics of Doubletime kinase-dependent degradation of the *Drosophila* Period protein. *J. Biol. Chem.* 286:27654–62
13. Lowrey PL, Shimomura K, Antoch MP, Yamazaki S, Zemenides PD, et al. 2000. Positional syntenic cloning and functional characterization of the mammalian circadian mutation *tau*. *Science* 288:483–92
14. Lee C, Etchegaray JP, Cagampang FR, Loudon AS, Reppert SM. 2001. Posttranslational mechanisms regulate the mammalian circadian clock. *Cell* 107:855–67
15. Takano A, Isojima Y, Nagai K. 2004. Identification of mPer1 phosphorylation sites responsible for the nuclear entry. *J. Biol. Chem.* 279:32578–85
16. Eide EJ, Woolf MF, Kang H, Woolf P, Hurst W, et al. 2005. Control of mammalian circadian rhythm by CKIε-regulated proteasome-mediated PER2 degradation. *Mol. Cell Biol.* 25:2795–807
17. Xu Y, Toh KL, Jones CR, Shin JY, Fu YH, Ptáček LJ. 2007. Modeling of a human circadian mutation yields insights into clock regulation by PER2. *Cell* 128:59–70
18. Meng QJ, Maywood ES, Bechtold DA, Lu WQ, Li J, et al. 2010. Entrainment of disrupted circadian behavior through inhibition of casein kinase 1 (CK1) enzymes. *Proc. Natl. Acad. Sci. USA* 107:15240–45
19. Lee HM, Chen R, Kim H, Etchegaray JP, Weaver DR, Lee C. 2011. The period of the circadian oscillator is primarily determined by the balance between casein kinase 1 and protein phosphatase 1. *Proc. Natl. Acad. Sci. USA* 108:16451–56
20. Shanware NP, Hutchinson JA, Kim SH, Zhan L, Bowler MJ, Tibbetts RS. 2011. Casein kinase 1-dependent phosphorylation of familial advanced sleep phase syndrome-associated residues controls PERIOD 2 stability. *J. Biol. Chem.* 286:12766–74
21. Chen Z, Yoo SH, Park YS, Kim KH, Wei S, et al. 2012. Identification of diverse modulators of central and peripheral circadian clocks by high-throughput chemical screening. *Proc. Natl. Acad. Sci. USA* 109:101–6
22. Chong SY, Ptáček LJ, Fu YH. 2012. Genetic insights on sleep schedules: this time, it's PERsonal. *Trends Genet.* 28:598–605
23. Voshall LB, Price JL, Sehgal A, Saez L, Young MW. 1994. Block in nuclear localization of Period protein by a second clock mutation, *timeless*. *Science* 263:1606–9
24. Saez L, Young MW. 1996. Regulation of nuclear entry of the *Drosophila* clock proteins Period and Timeless. *Neuron* 17:911–20
25. Hara T, Koh K, Combs DJ, Sehgal A. 2011. Post-translational regulation and nuclear entry of TIMELESS and PERIOD are affected in new timeless mutant. *J. Neurosci.* 31:9982–90
26. Saez L, Derasmo M, Meyer P, Stieglitz J, Young MW. 2011. A key temporal delay in the circadian cycle of *Drosophila* is mediated by a nuclear localization signal in the Timeless protein. *Genetics* 188:591–600
27. Hunter-Ensor M, Ousley A, Sehgal A. 1996. Regulation of the *Drosophila* protein Timeless suggests a mechanism for resetting the circadian clock by light. *Cell* 84:677–85
28. Myers MP, Wager-Smith K, Rothenfluh-Hilfiker A, Young MW. 1996. Light-induced degradation of TIMELESS and entrainment of the *Drosophila* circadian clock. *Science* 271:1736–40
29. Lim C, Allada R. 2013. ATAXIN-2 activates PERIOD translation to sustain circadian rhythms in *Drosophila*. *Science* 340:875–79
30. Zhang Y, Ling J, Yuan C, Dubruille R, Emery P. 2013. A role for *Drosophila* ATAXIN-2 in the activation of PERIOD translation and circadian behavior. *Science* 340:879–82
31. Meyer P, Saez L, Young MW. 2006. PER-TIM interactions in living *Drosophila* cells: an interval timer for the circadian clock. *Science* 311:226–29
32. Saez L, Meyer P, Young MW. 2007. A PER/TIM/DBT interval timer for *Drosophila*'s circadian clock. *Cold Spring Harb. Symp. Quant. Biol.* 72:69–74
33. Martinek S, Inonog S, Manoukian AS, Young MW. 2001. A role for the segment polarity gene *shaggy*/GSK-3 in the *Drosophila* circadian clock. *Cell* 105:769–79
34. Fang Y, Sathyanarayanan S, Sehgal A. 2007. Post-translational regulation of the *Drosophila* circadian clock requires protein phosphatase 1 (PP1). *Genes Dev.* 21:1506–18
35. Ko HW, Kim EY, Chiu J, Vanselow JT, Kramer A, Edery I. 2010. A hierarchical phosphorylation cascade that regulates the timing of PERIOD nuclear entry reveals novel roles for proline-directed kinases and GSK-3β/SGG in circadian clocks. *J. Neurosci.* 30:12664–75



36. Lin JM, Kilman VL, Keegan K, Paddock B, Emery-Le M, et al. 2002. A role for casein kinase 2 $\alpha$  in the *Drosophila* circadian clock. *Nature* 420:816–20
37. Akten B, Jauch E, Genova GK, Kim EY, Edery I, et al. 2003. A role for CK2 in the *Drosophila* circadian oscillator. *Nat. Neurosci.* 6:251–57
38. Nawathean P, Rosbash M. 2004. The Doubletime and CKII kinases collaborate to potentiate *Drosophila* PER transcriptional repressor activity. *Mol. Cell* 13:213–23
39. Maier B, Wendt S, Vanselow JT, Wallach T, Reischl S, et al. 2009. A large-scale functional RNAi screen reveals a role for CK2 in the mammalian circadian clock. *Genes Dev.* 23:708–18
40. Tsuchiya Y, Akashi M, Matsuda M, Goto K, Miyata Y, et al. 2009. Involvement of the protein kinase CK2 in the regulation of mammalian circadian rhythms. *Sci. Signal.* 2:ra26
41. Kim EY, Jeong EH, Park S, Jeong HJ, Edery I, Cho JW. 2012. A role for O-GlcNAcylation in setting circadian clock speed. *Genes Dev.* 26:490–502
42. Kaasik K, Kivimäe S, Allen JJ, Chalkley RJ, Huang Y, et al. 2013. Glucose sensor O-GlcNAcylation coordinates with phosphorylation to regulate circadian clock. *Cell Metab.* 17:291–302
43. Doi M, Hirayama J, Sassone-Corsi P. 2006. Circadian regulator CLOCK is a histone acetyltransferase. *Cell* 125:497–508
44. Hirayama J, Sahar S, Grimaldi B, Tamaru T, Takamatsu K, et al. 2007. CLOCK-mediated acetylation of BMAL1 controls circadian function. *Nature* 450:1086–90
45. Young MW, Kay SA. 2001. Time zones: a comparative genetics of circadian clocks. *Nat. Rev. Genet.* 2:702–15
46. Sathyanarayanan S, Zheng X, Kumar S, Chen CH, Chen D, et al. 2008. *Genes Dev.* 22:1522–33
47. Lamia KA, Sachdeva UM, DiTacchio L, Williams EC, Alvarez JG, et al. 2009. AMPK regulates the circadian clock by cryptochrome phosphorylation and degradation. *Science* 326:437–40
48. Um JH, Yang S, Yamazaki S, Kang H, Viollet B, et al. 2007. Activation of 5'-AMP-activated kinase with diabetes drug metformin induces casein kinase I $\epsilon$  (CKI $\epsilon$ )-dependent degradation of clock protein mPer2. *J. Biol. Chem.* 282:20794–98
49. Guarente L. 2011. Sirtuins, aging, and metabolism. *Cold Spring Harb. Symp. Quant. Biol.* 76:81–90
50. Bellet MM, Orozco-Solis R, Sahar S, Eckel-Mahan K, Sassone-Corsi P. 2011. The time of metabolism: NAD<sup>+</sup>, SIRT1, and the circadian clock. *Cold Spring Harb. Symp. Quant. Biol.* 76:31–38
51. Asher G, Gatfield D, Stratmann M, Reinke H, Dibner C, et al. 2008. SIRT1 regulates circadian clock gene expression through PER2 deacetylation. *Cell* 134:317–28
52. Nakahata Y, Kaluzova M, Grimaldi B, Sahar S, Hirayama J, et al. 2008. The NAD<sup>+</sup>-dependent deacetylase SIRT1 modulates CLOCK-mediated chromatin remodeling and circadian control. *Cell* 134:329–40
53. Asher G, Reinke H, Altmeyer M, Gutierrez-Arcelus M, Hottiger MO, Schibler U. 2010. Poly(ADP-ribose) polymerase 1 participates in the phase entrainment of circadian clocks to feeding. *Cell* 142:943–53
54. Nakahata Y, Sahar S, Astarita G, Kaluzova M, Sassone-Corsi P. 2009. Circadian control of the NAD<sup>+</sup> salvage pathway by CLOCK-SIRT1. *Science* 324:654–57
55. Yang X, Downes M, Yu RT, Bookout AL, He W, et al. 2006. Nuclear receptor expression links the circadian clock to metabolism. *Cell* 126:801–10
56. Fan W, Downes M, Atkins A, Yu R, Evans RM. 2011. Nuclear receptors and AMPK: resetting metabolism. *Cold Spring Harb. Symp. Quant. Biol.* 76:17–22
57. Lamia KA, Papp SJ, Yu RT, Barish GD, Uhlenhaut NH, et al. 2011. Cryptochromes mediate rhythmic repression of the glucocorticoid receptor. *Nature* 480:552–56
58. Schmutz I, Ripperger JA, Baeriswyl-Aebischer S, Albrecht U. 2010. The mammalian clock component PERIOD2 coordinates circadian output by interaction with nuclear receptors. *Genes Dev.* 24:345–57
59. Grimaldi B, Bellet MM, Katada S, Astarita G, Hirayama J, et al. 2010. PER2 controls lipid metabolism by direct regulation of PPAR $\gamma$ . *Cell Metab.* 12:509–20
60. Costa MJ, So AY, Kaasik K, Krueger KC, Pillsbury ML, et al. 2011. Circadian rhythm gene period 3 is an inhibitor of the adipocyte cell fate. *J. Biol. Chem.* 286:9063–70
61. Kozma-Bognar L, Kaldi K. 2008. Synchronization of the fungal and the plant circadian clock by light. *ChemBioChem* 9:2565–73
62. Huang W, Pérez-García P, Pokhilko A, Millar AJ, Antoshechkin I, et al. 2012. Mapping the core of the *Arabidopsis* circadian clock defines the network structure of the oscillator. *Science* 336:75–79

63. Huang ZJ, Edery I, Rosbash M. 1993. PAS is a dimerization domain common to *Drosophila* period and several transcription factors. *Nature* 364:259–62
64. Pellequer JL, Wager-Smith KA, Kay SA, Getzoff ED. 1998. Photoactive yellow protein: a structural prototype for the three-dimensional fold of the PAS domain superfamily. *Proc. Natl. Acad. Sci. USA* 95:5884–90
65. Taylor BL, Zhulin IB. 1999. PAS domains: internal sensors of oxygen, redox potential, and light. *Microbiol. Mol. Biol. Rev.* 63:479–506
66. Claridge-Chang A, Wijnen H, Naef F, Boothroyd C, Rajewsky N, Young MW. 2001. Circadian regulation of gene expression systems in the *Drosophila* head. *Neuron* 32:657–71
67. Gu YZ, Hogenesch JB, Bradfield CA. 2000. The PAS superfamily: sensors of environmental and developmental signals. *Annu. Rev. Pharmacol. Toxicol.* 40:519–61
68. Jackson FR, Bargiello TA, Yun SH, Young MW. 1986. Product of *per* locus of *Drosophila* shares homology with proteoglycans. *Nature* 320:185–88
69. Reddy P, Jacquier AC, Abovich N, Petersen G, Rosbash M. 1986. The *period* clock locus of *D. melanogaster* codes for a proteoglycan. *Cell* 46:53–61
70. Yildiz O, Doi M, Yujnovsky I, Cardone L, Berndt A, et al. 2005. Crystal structure and interactions of the PAS repeat region of the *Drosophila* clock protein PERIOD. *Mol. Cell* 17:69–82
71. King HA, Hoelz A, Crane BR, Young MW. 2011. Structure of an enclosed dimer formed by the *Drosophila* period protein. *J. Mol. Biol.* 413:561–72
72. Kucera N, Schmalen I, Hennig S, Öllinger R, Strauss HM, et al. 2012. Unwinding the differences of the mammalian PERIOD clock proteins from crystal structure to cellular function. *Proc. Natl. Acad. Sci. USA* 109:3311–16
73. Hennig S, Strauss HM, Vanselow K, Yildiz O, Schulze S, et al. 2009. Structural and functional analyses of PAS domain interactions of the clock proteins *Drosophila* PERIOD and mouse PERIOD2. *PLoS Biol.* 7:e94
74. Huang N, Chelliah Y, Shan Y, Taylor CA, Yoo S-H, et al. 2012. Crystal structure of the heterodimeric CLOCK:BMAL1 transcriptional activator complex. *Science* 337:189–94
75. Gekakis N, Saez L, Delahaye-Brown AM, Myers MP, Sehgal A, et al. 1995. Isolation of timeless by PER protein interaction: defective interaction between timeless protein and long-period mutant PERL. *Science* 270:811–15
76. Landskron J, Chen KF, Wolf E, Stanewsky R. 2009. A role for the PERIOD:PERIOD homodimer in the *Drosophila* circadian clock. *PLoS Biol.* 7:e3
77. Yoo S-H, Ko CH, Lowrey PL, Buhr ED, Song E-J, et al. 2005. A noncanonical E-box enhancer drives mouse Period2 oscillations in vivo. *Proc. Natl. Acad. Sci. USA* 102:2608–13
78. Nakahata Y, Yoshida M, Takano A, Soma H, Yamamoto T, et al. 2008. A direct repeat of E-box-like elements is required for cell-autonomous circadian rhythm of clock genes. *BMC Mol. Biol.* 9:1
79. Ayers RA, Moffat K. 2008. Changes in quaternary structure in the signaling mechanisms of PAS domains. *Biochemistry* 47:12078–86
80. Wang Z, Wu Y, Li L, Su X-D. 2013. Intermolecular recognition revealed by the complex structure of human CLOCK–BMAL1 basic helix-loop-helix domains with E-box DNA. *Cell Res.* 23:213–24
81. Ballario P, Macino G. 1997. White collar proteins: PASSing the light signal in *Neurospora crassa*. *Trends Microbiol.* 5:458–62
82. Talora C, Franchi L, Linden H, Ballario P, Macino G. 1999. Role of a white collar-1–white collar-2 complex in blue-light signal transduction. *EMBO J.* 18:4961–68
83. Cheng P, Yang Y, Gardner KH, Liu Y. 2002. PAS domain-mediated WC-1/WC-2 interaction is essential for maintaining the steady-state level of WC-1 and the function of both proteins in circadian clock and light responses of *Neurospora*. *Mol. Cell. Biol.* 22:517–24
84. Ballario P, Talora C, Galli D, Linden H, Macino G. 1998. Roles in dimerization and blue light photore-sponse of the PAS and LOV domains of *Neurospora crassa* white collar proteins. *Mol. Microbiol.* 29:719–29
85. Ballario P, Vittorioso P, Magrelli A, Talora C, Cabibbo A, Macino G. 1996. White collar-1, a central regulator of blue light responses in *Neurospora*, is a zinc finger protein. *EMBO J.* 15:1650–57

86. Bates DL, Chen Y, Kim G, Guo L, Chen L. 2008. Crystal structures of multiple GATA zinc fingers bound to DNA reveal new insights into DNA recognition and self-association by GATA. *J. Mol. Biol.* 381:1292–306
87. Viger RS, Guittot SM, Anttonen M, Wilson DB, Heikinheimo M. 2008. Role of the GATA family of transcription factors in endocrine development, function, and disease. *Mol. Endocrinol.* 22:781–98
88. Crosthwaite SK, Dunlap JC, Loros JJ. 1997. *Neurospora* wc-1 and wc-2: transcription, photoresponses, and the origins of circadian rhythmicity. *Science* 276:763–69
89. Froehlich AC, Liu Y, Loros JJ, Dunlap JC. 2002. White collar-1, a circadian blue light photoreceptor, binding to the frequency promoter. *Science* 297:815–19
90. He Q, Cheng P, Yang Y, Wang L, Gardner KH, Liu Y. 2002. White collar-1, a DNA binding transcription factor and a light sensor. *Science* 297:840–43
91. Brenna A, Grimaldi B, Filetici P, Ballario P. 2012. Physical association of the WC-1 photoreceptor and the histone acetyltransferase NGF-1 is required for blue light signal transduction in *Neurospora crassa*. *Mol. Biol. Cell* 23:3863–72
92. Dıoum EM, Rutter J, Tuckerman JR, Gonzalez G, Gilles-Gonzalez M-A, McKnight SL. 2002. NPAS2: a gas-responsive transcription factor. *Science* 298:2385–87
93. Koudo R, Kurokawa H, Sato E, Igarashi J, Uchida T, et al. 2005. Spectroscopic characterization of the isolated heme-bound PAS-B domain of neuronal PAS domain protein 2 associated with circadian rhythms. *FEBS J.* 272:4153–62
94. Mukaiyama Y, Uchida T, Sato E, Sasaki A, Sato Y, et al. 2006. Spectroscopic and DNA-binding characterization of the isolated heme-bound basic helix–loop–helix–PAS-A domain of neuronal PAS protein 2 (NPAS2), a transcription activator protein associated with circadian rhythms. *FEBS J.* 273:2528–39
95. Ishida M, Ueha T, Sagami I. 2008. Effects of mutations in the heme domain on the transcriptional activity and DNA-binding activity of NPAS2. *Biochem. Biophys. Res. Commun.* 368:292–97
96. Kitanishi K, Igarashi J, Hayasaka K, Hikage N, Saiful I, et al. 2008. Heme-binding characteristics of the isolated PAS-A domain of mouse Per2, a transcriptional regulatory factor associated with circadian rhythms. *Biochemistry* 47:6157–68
97. Airola MV, Du J, Dawson JH, Crane BR. 2010. Heme binding to the mammalian circadian clock protein period 2 is nonspecific. *Biochemistry* 49:4327–38
98. Crumbley C, Wang Y, Kojetin DJ, Burris TP. 2010. Characterization of the core mammalian clock component, NPAS2, as a REV-ERB $\alpha$ /ROR $\alpha$  target gene. *J. Biol. Chem.* 285:35386–92
99. Lukat-Rodgers GS, Correia C, Botuyan MV, Mer G, Rodgers KR. 2010. Heme-based sensing by the mammalian circadian protein CLOCK. *Inorg. Chem.* 49:6349–65
100. Hayasaka K, Kitanishi K, Igarashi J, Shimizu T. 2011. Heme-binding characteristics of the isolated PAS-B domain of mouse Per2, a transcriptional regulatory factor associated with circadian rhythms. *Biochim. Biophys. Acta* 1814:326–33
101. Raghuram S, Stayrook KR, Huang P, Rogers PM, Nosie AK, et al. 2007. Identification of heme as the ligand for the orphan nuclear receptors REV-ERB $\alpha$  and REV-ERB $\beta$ . *Nat. Struct. Mol. Biol.* 14:1207–13
102. Yin L, Wu N, Curtin JC, Qatanani M, Szewergold NR, et al. 2007. Rev-erb $\alpha$ , a heme sensor that coordinates metabolic and circadian pathways. *Science* 318:1786–89
103. Wu N, Yin L, Hanniman EA, Joshi S, Lazar MA. 2009. Negative feedback maintenance of heme homeostasis by its receptor, Rev-erb $\alpha$ . *Genes Dev.* 23:2201–9
104. Phelan CA, Gampe RT, Lambert MH, Parks DJ, Montana V, et al. 2010. Structure of Rev-erb $\alpha$  bound to N-CoR reveals a unique mechanism of nuclear receptor–co-repressor interaction. *Nat. Struct. Mol. Biol.* 17:808–14
105. Solt LA, Kojetin DJ, Burris TP. 2011. The REV-ERBs and RORs: molecular links between circadian rhythms and lipid homeostasis. *Future Med. Chem.* 3:623–38
106. Yang J, Kim KD, Lucas A, Drahos KE, Santos CS, et al. 2008. A novel heme-regulatory motif mediates heme-dependent degradation of the circadian factor period 2. *Mol. Cell. Biol.* 28:4697–711
107. Itoh R, Fujita K-I, Mu A, Kim DHT, Tai TT, et al. 2013. Imaging of heme/hemeproteins in nucleus of the living cells expressing heme-binding nuclear receptors. *FEBS Lett.* 587:2131–36

108. Kim EY, Ko HW, Yu W, Hardin PE, Edery I. 2007. A DOUBLETIME kinase binding domain on the *Drosophila* PERIOD protein is essential for its hyperphosphorylation, transcriptional repression, and circadian clock function. *Mol. Cell. Biol.* 27:5014–28
109. Nawatheatan P, Stoleru D, Rosbash M. 2007. A small conserved domain of *Drosophila* PERIOD is important for circadian phosphorylation, nuclear localization, and transcriptional repressor activity. *Mol. Cell. Biol.* 27:5002–13
110. Baylies MK, Vossahl LB, Sehgal A, Young MW. 1992. New short period mutations of the *Drosophila* clock gene *per*. *Neuron* 9:575–81
111. Rutla JE, Edery I, Hall JC, Rosbash M. 1992. The analysis of new short-period circadian rhythm mutants suggests features of *D. melanogaster* period gene function. *J. Neurogenet.* 8:101–13
112. Tomita T, Miyazaki K, Onishi Y, Honda S, Ishida N, Oishi K. 2010. Conserved amino acid residues in C-terminus of PERIOD 2 are involved in interaction with CRYPTOCHROME 1. *Biochim. Biophys. Acta* 1803:492–98
113. Yan Y, Barlev NA, Haley RH, Berger SL, Marmorstein R. 2000. Crystal structure of yeast Esa1 suggests a unified mechanism for catalysis and substrate binding by histone acetyltransferases. *Mol. Cell* 6:1195–205
114. Andrade MA, Petosa C, O'Donoghue SI, Müller CW, Bork P. 2001. Comparison of ARM and HEAT protein repeats. *J. Mol. Biol.* 309:1–18
115. Vodovar N, Clayton JD, Costa R, Odell M, Kyriacou CP. 2002. The *Drosophila* clock protein Timeless is a member of the Arm/HEAT family. *Curr. Biol.* 12:R610–11
116. Kippert F, Gerloff DL. 2004. Timeless and Armadillo: a link too far (comment on Vodovar et al. *Curr. Biol.* 2002. 12:R610–11). *Curr. Biol.* 14:R650–51
117. Kyriacou CP, Odell M. 2004. No ARM in it? (Reply to Kippert and Gerloff.) *Curr. Biol.* 14:R652–53
118. Perry J. 2005. Weighing in on a Timeless controversy. *Proteins* 61:699–703
119. Dunlap JC, Loros JJ. 2006. How fungi keep time: circadian system in *Neurospora* and other fungi. *Curr. Opin. Microbiol.* 9:579–87
120. Vitalini MW, de Paula RM, Park WD, Bell-Pedersen D. 2006. The rhythms of life: circadian output pathways in *Neurospora*. *J. Biol. Rhythms* 21:432–44
121. Baker CL, Loros JJ, Dunlap JC. 2012. The circadian clock of *Neurospora crassa*. *FEMS Microbiol. Rev.* 36:95–110
122. Cheng P, Yang Y, Heintzen C, Liu Y. 2001. Coiled-coil domain-mediated FRQ–FRQ interaction is essential for its circadian clock function in *Neurospora*. *EMBO J.* 20:101–8
123. Gorl M, Mellow M, Huttner B, Johnson J, Roenneberg T, Brunner M. 2001. A PEST-like element in FREQUENCY determines the length of the circadian period in *Neurospora crassa*. *EMBO J.* 20:7074–84
124. Schafmeier T, Káldi K, Diernfellner A, Mohr C, Brunner M. 2006. Phosphorylation-dependent maturation of *Neurospora* circadian clock protein from a nuclear repressor toward a cytoplasmic activator. *Genes Dev.* 20:297–306
125. Garceau NY, Liu Y, Loros JJ, Dunlap JC. 1997. Alternative initiation of translation and time-specific phosphorylation yield multiple forms of the essential clock protein FREQUENCY. *Cell* 89:469–76
126. Colot HV, Loros JJ, Dunlap JC. 2005. Temperature-modulated alternative splicing and promoter use in the circadian clock gene frequency. *Mol. Biol. Cell* 16:5563–71
127. Ruoff P, Loros JJ, Dunlap JC. 2005. The relationship between FRQ-protein stability and temperature compensation in the *Neurospora* circadian clock. *Proc. Natl. Acad. Sci. USA* 102:17681–86
128. Diernfellner A, Colot HV, Dintsis O, Loros JJ, Dunlap JC, Brunner M. 2007. Long and short isoforms of *Neurospora* clock protein FRQ support temperature-compensated circadian rhythms. *FEBS Lett.* 581:5759–64
129. Baker CL, Kettenbach AN, Loros JJ, Gerber SA, Dunlap JC. 2009. Quantitative proteomics reveals a dynamic interactome and phase-specific phosphorylation in the *Neurospora* circadian clock. *Mol. Cell* 34:354–63

130. Querfurth C, Diernfellner ACR, Gin E, Malzahn E, Höfer T, Brunner M. 2011. Circadian conformational change of the *Neurospora* clock protein FREQUENCY triggered by clustered hyperphosphorylation of a basic domain. *Mol. Cell* 43:713–22
131. Zhou M, Guo J, Cha J, Chae M, Chen S, et al. 2013. Non-optimal codon usage affects expression, structure and function of clock protein FRQ. *Nature* 495:111–15
132. Cheng P, He Q, He Q, Wang L, Liu Y. 2005. Regulation of the *Neurospora* circadian clock by an RNA helicase. *Genes Dev.* 19:234–41
133. Guo J, Cheng P, Liu Y. 2010. Functional significance of FRH in regulating the phosphorylation and stability of *Neurospora* circadian clock protein FRQ. *J. Biol. Chem.* 285:11508–15
134. Shi M, Collett M, Loros JJ, Dunlap JC. 2010. FRQ-interacting RNA helicase mediates negative and positive feedback in the *Neurospora* circadian clock. *Genetics* 184:351–61
135. Jackson RN, Klauer AA, Hintze BJ, Robinson H, van Hoof A, Johnson SJ. 2010. The crystal structure of Mtr4 reveals a novel arch domain required for rRNA processing. *EMBO J.* 29:2205–16
136. Weir JR, Bonneau F, Hentschel J, Conti E. 2010. Structural analysis reveals the characteristic features of Mtr4, a DExH helicase involved in nuclear RNA processing and surveillance. *Proc. Natl. Acad. Sci. USA* 107:12139–44
137. Guo J, Cheng P, Yuan H, Liu Y. 2009. The exosome regulates circadian gene expression in a posttranscriptional negative feedback loop. *Cell* 138:1236–46
138. Belden WJ, Loros JJ, Dunlap JC. 2007. Execution of the circadian negative feedback loop in *Neurospora* requires the ATP-dependent chromatin-remodeling enzyme CLOCKSWITCH. *Mol. Cell* 25:587–600
139. Chen C-H, Dunlap JC, Loros JJ. 2010. *Neurospora* illuminates fungal photoreception. *Fungal Genet. Biol.* 47:922–29
140. Cheng P, He Q, Yang Y, Wang L, Liu Y. 2003. Functional conservation of light, oxygen, or voltage domains in light sensing. *Proc. Natl. Acad. Sci. USA* 100:5938–43
141. Chen C-H, Ringelberg CS, Gross RH, Dunlap JC, Loros JJ. 2009. Genome-wide analysis of light-inducible responses reveals hierarchical light signalling in *Neurospora*. *EMBO J.* 28:1029–42
142. Malzahn E, Ciprianidis S, Káldi K, Schafmeier T, Brunner M. 2010. Photoadaptation in *Neurospora* by competitive interaction of activating and inhibitory LOV domains. *Cell* 142:762–72
143. Heintzen C, Loros JJ, Dunlap JC. 2001. The PAS protein VIVID defines a clock-associated feedback loop that represses light input, modulates gating, and regulates clock resetting. *Cell* 104:453–64
144. Schwerdtfeger C, Linden H. 2003. VIVID is a flavoprotein and serves as a fungal blue light photoreceptor for photoadaptation. *EMBO J.* 22:4846–55
145. Shrode LB, Lewis ZA, White LD, Bell-Pedersen D, Ebbole DJ. 2001. vvd is required for light adaptation of conidiation-specific genes of *Neurospora crassa*, but not circadian conidiation. *Fungal Genet. Biol.* 32:169–81
146. Crosson S, Rajagopal S, Moffat K. 2003. The LOV domain family: photoresponsive signaling modules coupled to diverse output domains. *Biochemistry* 42:2–10
147. Briggs WR. 2007. Phototropins and their LOV domains: versatile plant blue-light receptors. *J. Integr. Plant Biol.* 49:4–10
148. Herrou J, Crosson S. 2011. Function, structure and mechanism of bacterial photosensory LOV proteins. *Nat. Rev. Microbiol.* 9:713–23
149. Zoltowski BD, Gardner KH. 2011. Tripping the light fantastic: blue-light photoreceptors as examples of environmentally modulated protein-protein interactions. *Biochemistry* 50:4–16
150. Zoltowski BD, Schwerdtfeger C, Widom J, Loros JJ, Bilwes AM, et al. 2007. Conformational switching in the fungal light sensor Vivid. *Science* 316:1054–57
151. Zoltowski BD, Crane BR. 2008. Light activation of the LOV protein vivid generates a rapidly exchanging dimer. *Biochemistry* 47:7012–19
152. Zoltowski BD, Vaccaro B, Crane BR. 2009. Mechanism-based tuning of a LOV domain photoreceptor. *Nat. Chem. Biol.* 5:827–34
153. Vaidya AT, Chen C-H, Dunlap JC, Loros JJ, Crane BR. 2011. Structure of a light-activated LOV protein dimer that regulates transcription. *Sci. Signal.* 4:ra50



154. Lamb JS, Zoltowski BD, Pabit SA, Crane BR, Pollack L. 2008. Time-resolved dimerization of a PAS-LOV protein measured with photocoupled small angle X-ray scattering. *J. Am. Chem. Soc.* 130:12226–27
155. Lamb JS, Zoltowski BD, Pabit SA, Li L, Crane BR, Pollack L. 2009. Illuminating solution responses of a LOV domain protein with photocoupled small-angle X-ray scattering. *J. Mol. Biol.* 393:909–19
156. Peter E, Dick B, Baeurle SA. 2012. Illuminating the early signaling pathway of a fungal light-oxygen-voltage photoreceptor. *Proteins* 80:471–81
157. Hunt SM, Thompson S, Elvin M, Heintzen C. 2010. VIVID interacts with the WHITE COLLAR complex and FREQUENCY-interacting RNA helicase to alter light and clock responses in *Neurospora*. *Proc. Natl. Acad. Sci. USA* 107:16709–14
158. Sancar A. 2003. Structure and function of DNA photolyase and cryptochrome blue-light photoreceptors. *Chem. Rev.* 103:2203–37
159. Parth CL, Sancar A. 2005. Photochemistry and photobiology of cryptochrome blue-light photopigments: the search for a photocycle. *Photochem. Photobiol.* 81:1291–304
160. Chaves I, Pokorny R, Byrdin M, Hoang N, Ritz T, et al. 2011. The cryptochromes: blue light photoreceptors in plants and animals. *Annu. Rev. Plant Biol.* 62:335–64
161. Gegear RJ, Casselman A, Waddell S, Reppert SM. 2008. Cryptochrome mediates light-dependent magnetosensitivity in *Drosophila*. *Nature* 454:1014–18
162. Tang C-HA, Hinteregger E, Shang Y, Rosbash M. 2010. Light-mediated TIM degradation within *Drosophila* pacemaker neurons (s-LNvs) is neither necessary nor sufficient for delay zone phase shifts. *Neuron* 66:378–85
163. Fogle KJ, Parson KG, Dahm NA, Holmes TC. 2011. CRYPTOCHROME is a blue-light sensor that regulates neuronal firing rate. *Science* 331:1409–13
164. Kume K, Zylka MJ, Sriram S, Shearman LP, Weaver DR, et al. 1999. mCRY1 and mCRY2 are essential components of the negative limb of the circadian clock feedback loop. *Cell* 98:193–205
165. Lucas RJ, Hattar S, Takao M, Berson DM, Foster RG, Yau KW. 2003. Diminished pupillary light reflex at high irradiances in melanopsin-knockout mice. *Science* 299:245–47
166. Panda S, Provencio I, Tu DC, Pires SS, Rollag MD, et al. 2003. Melanopsin is required for non-image-forming photic responses in blind mice. *Science* 301:525–27
167. Sancar A. 2004. Regulation of the mammalian circadian clock by cryptochrome. *J. Biol. Chem.* 279:34079–82
168. Carell T, Burgdorf LT, Kundu LM, Cichon M. 2001. The mechanism of action of DNA photolyases. *Curr. Opin. Chem. Biol.* 5:491–98
169. Kanai S, Kikuno R, Toh H, Ryo H, Todo T. 1997. Molecular evolution of the photolyase-blue-light photoreceptor family. *J. Mol. Evol.* 45:535–48
170. Brudler R, Hitomi K, Daiyasu H, Toh H, Kucho K-I, et al. 2003. Identification of a new cryptochrome class. Structure, function, and evolution. *Mol. Cell* 11:59–67
171. Lin C, Todo T. 2005. The cryptochromes. *Genome Biol.* 6:220
172. Froy O, Chang DC, Reppert SM. 2002. Redox potential: differential roles in dCRY and mCRY1 functions. *Curr. Biol.* 12:147–52
173. Oztürk N, Song SH, Ozgür S, Selby CP, Morrison L, et al. 2007. Structure and function of animal cryptochromes. *Cold Spring Harb. Symp. Quant. Biol.* 72:119–31
174. Yuan Q, Metterville D, Briscoe AD, Reppert SM. 2007. Insect cryptochromes: Gene duplication and loss define diverse ways to construct insect circadian clocks. *Mol. Biol. Evol.* 24:948–55
175. Oztürk N, Song SH, Selby CP, Sancar A. 2008. Animal type 1 cryptochromes. Analysis of the redox state of the flavin cofactor by site-directed mutagenesis. *J. Biol. Chem.* 283:3256–63
176. Oztürk N, Selby CP, Song SH, Ye R, Tan C, et al. 2009. Comparative photochemistry of animal type 1 and type 4 cryptochromes. *Biochemistry* 48:8585–93
177. Naidoo N, Song W, Hunter-Ensor M, Sehgal A. 1999. A role for the proteasome in the light response of the timeless clock protein. *Science* 285:1737–41
178. Koh K, Zheng X, Sehgal A. 2006. JETLAG resets the *Drosophila* circadian clock by promoting light-induced degradation of TIMELESS. *Science* 312:1809–12



179. Peschel N, Chen KF, Szabo G, Stanewsky R. 2009. Light-dependent interactions between the *Drosophila* circadian clock factors cryptochrome, jetlag, and timeless. *Curr. Biol.* 19:241–47
180. Busza A, Emery-Le M, Rosbash M, Emery P. 2004. Roles of the two *Drosophila* CRYPTOCHROME structural domains in circadian photoreception. *Science* 304:1503–6
181. Oztürk N, VanVickle–Chavez SJ, Akileswaran L, Van Gelder RN, Sancar A. 2013. Ramshackle (Brwd3) promotes light-induced ubiquitylation of *Drosophila* cryptochrome by DDB1–CUL4–ROC1 E3 ligase complex. *Proc. Natl. Acad. Sci. USA* 110:4980–85
182. Ceriani MF, Darlington TK, Staknis D, Más P, Petti AA, et al. 1999. Light-dependent sequestration of TIMELESS by CRYPTOCHROME. *Science* 285:553–56
183. Ishikawa T, Hirayama J, Kobayashi Y, Todo T. 2002. Zebrafish CRY represses transcription mediated by CLOCK–BMAL heterodimer without inhibiting its binding to DNA. *Genes Cells* 7:1073–86
184. Hirayama J, Nakamura H, Ishikawa T, Kobayashi Y, Todo T. 2003. Functional and structural analyses of cryptochrome. Vertebrate CRY regions responsible for interaction with the CLOCK:BMAL1 heterodimer and its nuclear localization. *J. Biol. Chem.* 278:35620–28
185. Czarna A, Breitzkreuz H, Mahrenholz CC, Arens J, Strauss HM, Wolf E. 2011. Quantitative analyses of cryptochrome–mBMAL1 interactions: mechanistic insights into the transcriptional regulation of the mammalian circadian clock. *J. Biol. Chem.* 286:22414–25
186. Busino L, Bassermann F, Maiolica A, Lee C, Nolan PM, et al. 2007. SCF<sup>FBXL3</sup> controls the oscillation of the circadian clock by directing the degradation of cryptochrome proteins. *Science* 316:900–4
187. Siepka SM, Yoo S-H, Park J, Song W, Kumar V, et al. 2007. Circadian mutant Overtime reveals F-box protein FBXL3 regulation of cryptochrome and period gene expression. *Cell* 129:1011–23
188. Yoo S-H, Mohawk JA, Siepka SM, Shan Y, Huh SK, et al. 2013. Competing E3 ubiquitin ligases govern circadian periodicity by degradation of CRY in nucleus and cytoplasm. *Cell* 152:1091–105
189. Zoltowski BD, Vaidya AT, Top D, Widom J, Young MW, Crane BR. 2011. Structure of full-length *Drosophila* cryptochrome. *Nature* 480:396–99
190. Czarna A, Berndt A, Singh HR, Grudziecki A, Ladurner AG, et al. 2013. Structures of *Drosophila* cryptochrome and mouse cryptochrome 1 provide insight into circadian function. *Cell* 153:1394–405
191. Levy C, Zoltowski BD, Jones AR, Vaidya AT, Top D, et al. 2013. Updated structure of *Drosophila* cryptochrome. *Nature* 495:E3–4
192. Berndt A, Kottke T, Breitzkreuz H, Dvorsky R, Hennig S, et al. 2007. A novel photoreaction mechanism for the circadian blue light photoreceptor *Drosophila* cryptochrome. *J. Biol. Chem.* 282:13011–21
193. Selby CP, Sancar A. 2012. The second chromophore in *Drosophila* photolyase/cryptochrome family photoreceptors. *Biochemistry* 51:167–71
194. Oztürk N, Selby CP, Annayev Y, Zhong D, Sancar A. 2011. Reaction mechanism of *Drosophila* cryptochrome. *Proc. Natl. Acad. Sci. USA* 108:516–21
195. Xing W, Busino L, Hinds TR, Marionni ST, Saifee NH, et al. 2013. SCF<sup>FBXL3</sup> ubiquitin ligase targets cryptochromes at their cofactor pocket. *Nature* 496:64–68
196. Vieira J, Jones AR, Danon A, Sakuma M, Hoang N, et al. 2012. Human cryptochrome-1 confers light independent biological activity in transgenic *Drosophila* correlated with flavin radical stability. *PLoS ONE* 7:e31867
197. Hirota T, Lee JW, St John PC, Sawa M, Iwaisako K, et al. 2012. Identification of small molecule activators of cryptochrome. *Science* 337:1094–97
198. Nangle S, Zing W, Zheng N. 2013. Crystal structure of mammalian cryptochrome in complex with a small molecule competitor of its ubiquitin ligase. *Cell Res.* 23:1417–19
199. Tamai TK, Young LC, Whitmore D. 2007. Light signaling to the zebrafish circadian clock by cryptochrome 1a. *Proc. Natl. Acad. Sci. USA* 104:14712–17
200. Rosato E, Codd V, Mazzotta G, Piccin A, Zordan M, et al. 2001. Light-dependent interaction between *Drosophila* CRY and the clock protein PER mediated by the carboxy terminus of CRY. *Curr. Biol.* 11:909–17
201. Hoang H, Huang X, Skibo EB. 2008. Synthesis and in vitro evaluation of imidazole-based wakayin analogues. *Org. Biomol. Chem.* 6:3059–64

202. Li X, Wang Q, Yu X, Liu H, Yang H, et al. 2011. *Arabidopsis* cryptochrome 2 (CRY2) functions by the photoactivation mechanism distinct from the tryptophan (trp) triad-dependent photoreduction. *Proc. Natl. Acad. Sci. USA* 108:20844–49
203. Vaidya AT, Top D, Manahan CC, Tokuda JM, Zhang S, et al. 2013. Flavin reduction activates *Drosophila* cryptochrome. *Proc. Natl. Acad. Sci. USA* 110:20455–60
204. Ozturk N, Selby CP, Zhong D, Sancar A. 2013. Mechanism of photoreduction by *Drosophila* cryptochrome: role of the redox status of the flavin cofactor. *J. Biol. Chem.* 289:4634–42
205. Chaves I, Nijman RM, Biernat MA, Bajek MI, Brand K, et al. 2011. The Potorous CPD photolyase rescues a cryptochrome-deficient mammalian circadian clock. *PLoS ONE* 6:e23447
206. Berson DM. 2003. Strange vision: ganglion cells as circadian photoreceptors. *Trends Neurosci.* 26:314–20
207. Hankins MW, Peirson SN, Foster RG. 2008. Melanopsin: an exciting photopigment. *Trends Neurosci.* 31:27–36
208. Schmidt TM, Chen S-K, Hattar S. 2011. Intrinsically photosensitive retinal ganglion cells: many subtypes, diverse functions. *Trends Neurosci.* 34:572–80
209. Sexton T, Buhr E, Van Gelder RN. 2012. Melanopsin and mechanisms of non-visual ocular photoreception. *J. Biol. Chem.* 287:1649–56
210. Lucas RJ. 2013. Mammalian inner retinal photoreception. *Curr. Biol.* 23:R125–33
211. Gooley JJ, Lu J, Chou TC, Scammell TE, Saper CB. 2001. Melanopsin in cells of origin of the retino-hypothalamic tract. *Nat. Neurosci.* 4:1165
212. Berson DM, Dunn FA, Takao M. 2002. Phototransduction by retinal ganglion cells that set the circadian clock. *Science* 295:1070–73
213. Hattar S, Liao HW, Takao M, Berson DM, Yau KW. 2002. Melanopsin-containing retinal ganglion cells: architecture, projections, and intrinsic photosensitivity. *Science* 295:1065–70
214. Provencio I, Jiang G, De Grip WJ, Hayes WP, Rollag MD. 1998. Melanopsin: an opsin in melanophores, brain, and eye. *Proc. Natl. Acad. Sci. USA* 95:340–45
215. Provencio I, Rodriguez IR, Jiang G, Hayes WP, Moreira EF, Rollag MD. 2000. A novel human opsin in the inner retina. *J. Neurosci.* 20:600–5
216. Panda S, Antoch MP, Miller BH, Su AI, Schook AB, et al. 2002. Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell* 109:307–20
217. Ruby NF, Brennan TJ, Xie X, Cao V, Franken P, et al. 2002. Role of melanopsin in circadian responses to light. *Science* 298:2211–13
218. Bellingham J, Chaurasia SS, Melyan Z, Liu C, Cameron MA, et al. 2006. Evolution of melanopsin photoreceptors: discovery and characterization of a new melanopsin in nonmammalian vertebrates. *PLoS Biol.* 4:e254
219. Kouyama T, Murakami M. 2010. Structural divergence and functional versatility of the rhodopsin superfamily. *Photochem. Photobiol. Sci.* 9:1458–65
220. Sekharan S, Wei JN, Batista VS. 2012. The active site of melanopsin: the biological clock photoreceptor. *J. Am. Chem. Soc.* 134:19536–39
221. Hermann R, Poppe L, Pilbák S, Boden C, Maurer J, et al. 2005. Predicted 3D structure of melanopsin, the non-rod, non-cone photopigment of the mammalian circadian clock, from Djungarian hamsters (*Phodopus sungorus*). *Neurosci. Lett.* 376:76–80
222. Kratochwil NA, Gatti-McArthur S, Hoener MC, Lindemann L, Christ AD, et al. 2011. G protein-coupled receptor transmembrane binding pockets and their applications in GPCR research and drug discovery: a survey. *Curr. Topics Med. Chem.* 11:1902–24
223. Murakami M, Kouyama T. 2008. Crystal structure of squid rhodopsin. *Nature* 453:363–67
224. Zhou XE, Melcher K, Xu HE. 2012. Structure and activation of rhodopsin. *Acta Pharmacol. Sin.* 33:291–99
225. Scheerer P, Park JH, Hildebrand PW, Kim YJ, Krauss N, et al. 2008. Crystal structure of opsin in its G-protein-interacting conformation. *Nature* 455:497–502
226. Choe H-W, Kim YJ, Park JH, Morizumi T, Pai EF, et al. 2011. Crystal structure of metarhodopsin II. *Nature* 471:651–55

227. Standfuss J, Edwards PC, D'Antona A, Fransen M, Xie G, et al. 2011. The structural basis of agonist-induced activation in constitutively active rhodopsin. *Nature* 471:656–60
228. Rasmussen SGF, DeVree BT, Zou Y, Kruse AC, Chung KY, et al. 2011. Crystal structure of the  $\beta_2$  adrenergic receptor–G<sub>s</sub> protein complex. *Nature* 477:549–55
229. Chung KY, Rasmussen SGF, Liu T, Li S, DeVree BT, et al. 2011. Conformational changes in the G protein G<sub>s</sub> induced by the  $\beta_2$  adrenergic receptor. *Nature* 477:611–15
230. Murakami M, Kouyama T. 2011. Crystallographic analysis of the primary photochemical reaction of squid rhodopsin. *J. Mol. Biol.* 413:615–27
231. McDonald MJ, Rosbash M. 2001. Microarray analysis and organization of circadian gene expression in *Drosophila*. *Cell* 107:567–78
232. Koike N, Yoo SH, Huang HC, Kumar V, Lee C, et al. 2012. Transcriptional architecture and chromatin landscape of the core circadian clock in mammals. *Science* 338:349–54
233. Rodriguez J, Tang CH, Khodor YL, Vodala S, Menet JS, Rosbash M. 2013. Nascent-Seq analysis of *Drosophila* cycling gene expression. *Proc. Natl. Acad. Sci. USA* 110:E275–284



# Contents

Journeys in Science: Glycobiology and Other Paths <i>Raymond A. Dwek</i> .....	1
Lipids and Extracellular Materials <i>William Dowhan</i> .....	45
Topological Regulation of Lipid Balance in Cells <i>Guillaume Drin</i> .....	51
Lipidomics: Analysis of the Lipid Composition of Cells and Subcellular Organelles by Electrospray Ionization Mass Spectrometry <i>Britta Brügger</i> .....	79
Biosynthesis and Export of Bacterial Lipopolysaccharides <i>Chris Whitfield and M. Stephen Trent</i> .....	99
Demystifying Heparan Sulfate-Protein Interactions <i>Ding Xu and Jeffrey D. Esko</i> .....	129
Dynamics and Timekeeping in Biological Systems <i>Christopher M. Dobson</i> .....	159
Metabolic and Nontranscriptional Circadian Clocks: Eukaryotes <i>Akhilesh B. Reddy and Guillaume Rey</i> .....	165
Interactive Features of Proteins Composing Eukaryotic Circadian Clocks <i>Brian R. Crane and Michael W. Young</i> .....	191
Metabolic Compensation and Circadian Resilience in Prokaryotic Cyanobacteria <i>Carl Hirschbie Johnson and Martin Egli</i> .....	221
Activity-Based Profiling of Proteases <i>Laura E. Sanman and Matthew Bogoy</i> .....	249
Asymmetry of Single Cells and Where That Leads <i>Mark S. Bretscher</i> .....	275
Bringing Dynamic Molecular Machines into Focus by Methyl-TROSY NMR <i>Rina Rosenzweig and Lewis E. Kay</i> .....	291

Chlorophyll Modifications and Their Spectral Extension in Oxygenic Photosynthesis <i>Min Chen</i> .....	317
Enzyme Inhibitor Discovery by Activity-Based Protein Profiling <i>Micah J. Niphakis and Benjamin F. Cravatt</i> .....	341
Expanding and Reprogramming the Genetic Code of Cells and Animals <i>Jason W. Chin</i> .....	379
Genome Engineering with Targetable Nucleases <i>Dana Carroll</i> .....	409
Hierarchy of RNA Functional Dynamics <i>Anthony M. Mustoe, Charles L. Brooks, and Hashim M. Al-Hashimi</i> .....	441
High-Resolution Structure of the Eukaryotic 80S Ribosome <i>Gulnara Yusupova and Marat Yusupov</i> .....	467
Histone Chaperones: Assisting Histone Traffic and Nucleosome Dynamics <i>Zachary A. Gurard-Levin, Jean-Pierre Quivy, and Geneviève Almouzni</i> .....	487
Human RecQ Helicases in DNA Repair, Recombination, and Replication <i>Deborah L. Croteau, Venkateswarlu Popuri, Patricia L. Opresko, and Vilhelm A. Bohr</i> .....	519
Intrinsically Disordered Proteins and Intrinsically Disordered Protein Regions <i>Christopher J. Oldfield and A. Keith Dunker</i> .....	553
Mechanism and Function of Oxidative Reversal of DNA and RNA Methylation <i>Li Shen, Chun-Xiao Song, Chuan He, and Yi Zhang</i> .....	585
Progress Toward Synthetic Cells <i>J. Craig Blain and Jack W. Szostak</i> .....	615
PTEN <i>Carolyn A. Worby and Jack E. Dixon</i> .....	641
Regulating the Chromatin Landscape: Structural and Mechanistic Perspectives <i>Blaine Bartholomew</i> .....	671
RNA Helicase Proteins as Chaperones and Remodelers <i>Inga Jarmoskaite and Rick Russell</i> .....	697

Selection-Based Discovery of Druglike Macrocyclic Peptides <i>Toby Passioura, Takayuki Katoh, Yuki Goto, and Hiroaki Suga</i> .....	727
Small Proteins Can No Longer Be Ignored <i>Gisela Storz, Yuri I. Wolf, and Kumaran S. Ramamurthi</i> .....	753
The Scanning Mechanism of Eukaryotic Translation Initiation <i>Alan G. Hinnebusch</i> .....	779
Understanding Nucleic Acid–Ion Interactions <i>Jan Lipfert, Sebastian Doniach, Rhiju Das, and Daniel Herschlag</i> .....	813

## Indexes

Cumulative Index of Contributing Authors, Volumes 79–83 .....	843
Cumulative Index of Article Titles, Volumes 79–83 .....	847

## Errata

An online log of corrections to *Annual Review of Biochemistry* articles may be found at <http://www.annualreviews.org/errata/biochem>





# ANNUAL REVIEWS

It's about time. Your time. It's time well spent.

## New From Annual Reviews:

### ***Annual Review of Statistics and Its Application***

Volume 1 • Online January 2014 • <http://statistics.annualreviews.org>

Editor: **Stephen E. Fienberg**, *Carnegie Mellon University*

Associate Editors: **Nancy Reid**, *University of Toronto*

**Stephen M. Stigler**, *University of Chicago*

The *Annual Review of Statistics and Its Application* aims to inform statisticians and quantitative methodologists, as well as all scientists and users of statistics about major methodological advances and the computational tools that allow for their implementation. It will include developments in the field of statistics, including theoretical statistical underpinnings of new methodology, as well as developments in specific application domains such as biostatistics and bioinformatics, economics, machine learning, psychology, sociology, and aspects of the physical sciences.

**Complimentary online access to the first volume will be available until January 2015.**

#### TABLE OF CONTENTS:

- *What Is Statistics?* Stephen E. Fienberg
- *A Systematic Statistical Approach to Evaluating Evidence from Observational Studies*, David Madigan, Paul E. Stang, Jesse A. Berlin, Martijn Schuemie, J. Marc Overhage, Marc A. Suchard, Bill Dumouchel, Abraham G. Hartzema, Patrick B. Ryan
- *The Role of Statistics in the Discovery of a Higgs Boson*, David A. van Dyk
- *Brain Imaging Analysis*, F. DuBois Bowman
- *Statistics and Climate*, Peter Guttorp
- *Climate Simulators and Climate Projections*, Jonathan Rougier, Michael Goldstein
- *Probabilistic Forecasting*, Tilmann Gneiting, Matthias Katzfuss
- *Bayesian Computational Tools*, Christian P. Robert
- *Bayesian Computation Via Markov Chain Monte Carlo*, Radu V. Craiu, Jeffrey S. Rosenthal
- *Build, Compute, Critique, Repeat: Data Analysis with Latent Variable Models*, David M. Blei
- *Structured Regularizers for High-Dimensional Problems: Statistical and Computational Issues*, Martin J. Wainwright
- *High-Dimensional Statistics with a View Toward Applications in Biology*, Peter Bühlmann, Markus Kalisch, Lukas Meier
- *Next-Generation Statistical Genetics: Modeling, Penalization, and Optimization in High-Dimensional Data*, Kenneth Lange, Jeanette C. Papp, Janet S. Sinsheimer, Eric M. Sobel
- *Breaking Bad: Two Decades of Life-Course Data Analysis in Criminology, Developmental Psychology, and Beyond*, Elena A. Erosheva, Ross L. Matsueda, Donatello Telesca
- *Event History Analysis*, Niels Keiding
- *Statistical Evaluation of Forensic DNA Profile Evidence*, Christopher D. Steele, David J. Balding
- *Using League Table Rankings in Public Policy Formation: Statistical Issues*, Harvey Goldstein
- *Statistical Ecology*, Ruth King
- *Estimating the Number of Species in Microbial Diversity Studies*, John Bunge, Amy Willis, Fiona Walsh
- *Dynamic Treatment Regimes*, Bibhas Chakraborty, Susan A. Murphy
- *Statistics and Related Topics in Single-Molecule Biophysics*, Hong Qian, S.C. Kou
- *Statistics and Quantitative Risk Management for Banking and Insurance*, Paul Embrechts, Marius Hofert

Access this and all other Annual Reviews journals via your institution at [www.annualreviews.org](http://www.annualreviews.org).

## ANNUAL REVIEWS | Connect With Our Experts

Tel: 800.523.8635 (US/CAN) | Tel: 650.493.4400 | Fax: 650.424.0910 | Email: [service@annualreviews.org](mailto:service@annualreviews.org)

