

index

April 21, 2018

0.1 Introduction to Finite Element Methods

0.1.1 Gauss quadrature rules

0.1.2 Yerlan Amanbek

yerlan@utexas.edu

1 Objective

- Revisit numerical integration methods
- Gauss integration(optimal)

1.1 Set-up

The notebook relies on Python code. To initialize the notebook, select **Cell->Run All Below**

1.2 Introduction

In general, case a mapping between elements and reference(master) elements is used to simplify the integration domain.

We need to compute the definite integral a lot in the FEM computations.

$$I := \int_a^b f(x) dx$$

The integral can be viewed graphically as the area between the x-axis and the curve $y = f(x)$ in the region of the limits of integration. Thus, we can interpret numerical integration as an approximation of that area.

```
In [1]: from ipywidgets import interact, fixed, IntSlider, FloatSlider, Text, Dropdown
        %matplotlib inline
        import areaTools as AT
        import ipywidgets as widgets
```

```
In [2]: %%javascript
        IPython.OutputArea.prototype._should_scroll = function(lines) {
            return false;
        }
```

<IPython.core.display.Javascript object>

In [3]: *## Do not change the text in this box*

```
f_box = Text(value="exp(-x**2)", description=r'$f(x)=$')
n_slider = widgets.IntSlider(min=2,max=20, step=2, value=4, description=r'$n_{start}$')
method_type = Dropdown(options=['left', 'right', 'midpoint', 'trapezoid', 'simpson'],
                        value='left', description='Method:')
interact(AT.plot3Areas,f=f_box,a="-1.0",b="1.0",n=n_slider,method=method_type);

interactive(children=(Text(value='exp(-x**2)', description='$f(x)=$'), Text(value='-1.0', desc
```

Some quick explanations of how the notebook works:

The meaning of the variables are as follows: - "a": the left endpoint of the integration interval - "b": the right endpoint of the integration interval - "n" the number of sub-intervals to use for approximating the integral - "f(x)": the function to integrate.

How to define the function: - $x**p$ means x^p . For example, $x**0.5$ means \sqrt{x} - for a constant function $f(x) = c$ use $c * (x * 0)$ (every function definition needs an x in it!) - for e^p use $\exp(p)$ (so for e use $\exp(1)$) - for π use π - type trig functions the way you would on your calculator($\sin(x)$, $\cos(x)$, etc)

Some functions you might want to try: - $4-x**2$ - $\cos(x)$ - $\sin(x**2)$ - $\exp(x**2)$ - $\exp(-x**2/2)/\sqrt{2*\pi}$ (this is the famous "normal curve") - $\exp(x)/x$ (make sure that 0 is not part of your interval!)

$$\int_a^b f(x) dx \approx \sum_{i=1}^n w_i f(x_i)$$

The trapezoidal rule of numerical integration simply approximates the area by the sum of several equally spaced trapezoids under the curve between the limits of a and b.

The height of a trapezoid is found from the integrand, $y_j = y(x_j)$, evaluated at equally spaced points, x_j and x_{j+1} . Thus, a typical contribution is $Area = h \frac{y_j + y_{j+1}}{2}$, where $h = x_{j+1} - x_j$ is the spacing. Thus, for n points (and $n-1$ spaces), the well-known approximation is

$$I \approx h \left(\frac{1}{2} y_1 + y_2 + y_3 + \dots + y_{n-1} + \frac{1}{2} y_n \right)$$

where $w_j = h$, except $w_1 = w_n = \frac{h}{2}$.

In [4]: *## Do not change the text in this box*

```
# -3*x**2+2*x+6
f_box = Text(value="exp(-x**2)", description=r'$f(x)=$')
n_slider = widgets.IntSlider(min=1,max=20, step=2, value=2, description=r'$n_{start}$')
method_type = Dropdown(options=['trapezoid'],
                        value='trapezoid', description='Method:')
interact(AT.plot3Areas,f=f_box,a="-1.0",b="1.0",n=n_slider,method=method_type);

interactive(children=(Text(value='exp(-x**2)', description='$f(x)=$'), Text(value='-1.0', desc
```

2 Gauss integration

The famous mathematician Gauss posed this question:

- What is the minimum number of points, n , required to exactly integrate a polynomial?
- What are the corresponding function values and weights?

3 Example:

Let $[a, b] = [-1, 1]$ and $n = 2$

We want to choose x_1, x_2, w_1 and w_2 so that

$$\int_{-1}^1 f(x) dx \approx w_1 f(x_1) + w_2 f(x_2)$$

The approximation should be exact for any polynomial of degree 3 or less.

Let $f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3$

then

$$\int_{-1}^1 f(x) dx = a_0 \int_{-1}^1 1 dx + a_1 \int_{-1}^1 x dx + a_2 \int_{-1}^1 x^2 dx + a_3 \int_{-1}^1 x^3 dx$$

Each of the definite integrals on the right-hand side has an integrand of degree 3 or less. Gaussian Quadrature should be exact for each of these.

Applying Gaussian Quadrature to each remaining integral yields:

$$\int_{-1}^1 1 dx = 2 = w_1 + w_2$$

$$\int_{-1}^1 x dx = 0 = w_1 x_1 + w_2 x_2$$

$$\int_{-1}^1 x^2 dx = \frac{2}{3} = w_1 x_1^2 + w_2 x_2^2$$

$$\int_{-1}^1 x^3 dx = 0 = w_1 x_1^3 + w_2 x_2^3$$

$$w_1 + w_2 = 2$$

$$w_1 x_1 + w_2 x_2 = 0 \text{ we obtain } w_1 = \frac{2x_2}{x_2 - x_1} \text{ and } w_2 = \frac{2x_1}{x_1 - x_2}$$

Using these values for 3rd equation $x_1 x_2 = -\frac{1}{3}$ and 4th equation $x_1 = -x_2$.

Solving the nonlinear system gives us

$$w_1 = 1, w_2 = 1, x_1 = -\frac{\sqrt{3}}{3}, x_2 = \frac{\sqrt{3}}{3}$$

Therefore,

$$\int_{-1}^1 f(x) dx \approx f\left(\frac{-\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right)$$

Above Gauss quadrature integral has degree of precision 3. Trapezoidal rule has degree of precision 1.

Number of points, n	Weights, w_i	Points, x_i
1	2	0
2	1	$\pm \frac{1}{\sqrt{3}}$
3	0	$\pm \frac{8}{9}\sqrt{3}$
	$\pm \sqrt{\frac{3}{5}}$	$\pm \frac{5}{9}$
4	$\frac{18+\sqrt{30}}{36}$	$\pm \sqrt{\frac{3}{7} - \frac{2}{7}\sqrt{\frac{6}{5}}}$
	$\frac{18-\sqrt{30}}{36}$	$\pm \sqrt{\frac{3}{7} + \frac{2}{7}\sqrt{\frac{6}{5}}}$

4 Example

$$\int_{-1}^1 (x^2 + \cos x) dx = \left. \frac{x^3}{3} \right|_{-1}^1 + \sin x \Big|_{-1}^1 = \frac{2}{3} + 2\sin(1)$$

Trapezoidal rule :

$$\int_{-1}^1 (x^2 + \cos x) dx \approx \frac{2}{2} (f(-1) + f(1))$$

$$\text{Gauss quadrature: } \int_{-1}^1 (x^2 + \cos x) dx \approx f\left(\frac{-\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right)$$

```
In [22]: from scipy.integrate import quad
         from numpy import sqrt, sin, cos, pi
```

```
def f(x):
    return x**2 + cos(x) #integrant
```

```
In [23]: I_actual = 2/3 + 2*sin(1) #computed integral
```

```
print("Analytical computation of integral      is %1.10f" % I_actual)
```

```
#Trapezoidal rule for integral computation
```

```
h=2
```

```
I_trapezoidal = h*(f(-1)+f(1))/2
```

```
error_trapezoidal = abs(I_actual - I_trapezoidal)
```

```
print("Trapezoidal rule computation of integral is %1.10f and error is %1.10f" % (I_t
```

```
Analytical computation of integral      is 2.3496086363
```

```
Trapezoidal rule computation of integral is 3.0806046117 and error is 0.7309959755
```

```
In [24]: #Gauss quadrature rule for integral computation
```

```
I_Gauss_Quadrature = f(-sqrt(3)/3)+f(sqrt(3)/3)
```

```
error_gauss = abs(I_actual - I_Gauss_Quadrature)
```

```
print("Analytical computation of integral      is %1.10f" % I_actual)
```

```
print("Gauss quadrature computation of integral is %1.10f and error is %1.10f" % (I_G
```