

NICE PROOFS

SCALLOP YE

Proposition 1. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function with period T , and let $x_0 \in \mathbb{R}$. If $f(x_0) \neq 0$ and $\int_0^T f(x)dx = 0$, then f has at least two zeros in the interval $I = (x_0, x_0 + T)$.*

Proof. The function f is periodic with period T , so

$$\begin{aligned} f(x_0 + T) &= f(x_0) \neq 0, \\ \int_{x_0}^{x_0+T} f(x)dx &= \int_0^T f(x)dx = 0. \end{aligned}$$

Without loss of generality we may assume that $f(x_0 + T) = f(x_0) > 0$. By the mean value theorem for definite integrals, there exists a $c \in I$ such that

$$f(c) = \frac{1}{T} \int_{x_0}^{x_0+T} f(x)dx = 0.$$

Hence f has at least one zero in I . Suppose for sake of contradiction that f has only one zero in I . Again, suppose that there exists a $b \in I$ such that $f(b) < 0$. By the intermediate value theorem f has an extra zero in (x_0, b) or $(b, x_0 + T)$, a contradiction. Thus we have

$$f(x) \geq 0 \quad \forall x \in I.$$

Since f is continuous, there exists an $a \in I$ such that

$$f(x) \geq \frac{f(x_0)}{2} \quad \forall x \in (x_0, a).$$

By the properties of the Riemann integral

$$\begin{aligned} \int_{x_0}^a f(x)dx &\geq \int_{x_0}^a \frac{f(x_0)}{2} dx = \frac{a - x_0}{2} f(x_0) > 0, \\ \int_a^{x_0+T} f(x)dx &\geq 0, \\ \int_{x_0}^{x_0+T} f(x)dx &= \int_{x_0}^a f(x)dx + \int_a^{x_0+T} f(x)dx > 0. \end{aligned}$$

This contradicts the fact that $\int_{x_0}^{x_0+T} f(x)dx = 0$. Thus f has at least two zeros in I , as desired. \square