

Search for New Physics in All-hadronic Events with AlphaT in 8 TeV data at CERN

Yossof Eshaq

*Submitted in Partial Fulfillment of the
Requirements for the Degree Doctor of Philosophy*

Supervised by Professor Aran Garcia-Bellido

Department of Physics

Astronomy

Arts, Sciences and Engineering

University of Rochester

October 2, 2014

Abstract

An inclusive search for supersymmetric processes that produce final states with jets and missing transverse energy is performed in pp collisions at a centre-of-mass energy of $\sqrt{s} = 8$ TeV. The data sample corresponds to an integrated luminosity of 18.5 fb^{-1} collected by the CMS experiment at the LHC. In this search, a dimensionless kinematic variable, α_T , is used to discriminate between events with genuine and misreconstructed missing transverse energy. The search is based on an examination of the number of reconstructed jets per event, the scalar sum of transverse energies of these jets, and the number of these jets identified as originating from bottom quarks. The results are interpreted with various simplified models, with a special emphasis on models with a compressed mass spectrum.

0.1 Theoretical motivation

SM,Higgs,SUSY

Particle physics concerns itself with the study of particles and fields. Our current knowledge of their characteristics and interactions are formalized in the quantum field theory called the Standard Model. It through three symmetries: The color charge symmetry of Quantum Chromo Dynamics (QCD) represented in $SU(3)$, the flavor symmetry of Quantum Flavor Dynamics (QFD) represented in $SU(2)$ and the electric charge symmetry of Quantum Electro Dynamics represented in $U(1)$. Together, $SU(3) \times SU(2) \times U(1)$ represent the field theory.

0.2 LHC and CMS

LHC, CMS

0.3 Definition of $\alpha_{\mathbf{T}}$

0.4 Data sets and Monte Carlo samples

0.4.1 Data sets

The data analysed consist of the full run of 2012.

The following datasets are used to populate the hadronic signal and control samples. They correspond to the full data run of 2012 and an integrated luminosity of $19.45 \pm 0.8 \text{ fb}^{-1}$. The official JSON from the 22nd Jan 2013 is used to filter only certified luminosity sections with the run range 190456–208686.

Table 1: Datasets.

| Dataset | Luminosity (fb^{-1}) |
|----------|---------------------------------|
| Hadronic | 19.45 |
| Muon | 19.72 |
| Photon | 19.63 |

0.4.2 MC samples for signal and SM backgrounds

The SM background Monte Carlo samples for physics at 8 TeV are taken from the Summer12 simulation production run with CMSSW_5_3_X with the PU_S10 scenario. The effective luminosity of each MC sample is normalised to the integrated luminosity of the corresponding dataset, as listed in Table 2. The signal MADGRAPH Monte Carlo samples, listed in Table ??, are taken from a FastSim simulation production based on CMSSW_5_2_X. All MC samples are reweighted on an event-by-event basis such that the distribution of pile-up (PU) interactions matches that observed in data. This is done using the recommended recipe and the PU JSON of 13th December 2012.

Table 2: MC samples for Standard Model processes.

| Sample | HT (GeV) | Cross section (pb) | Corrected Cross section (pb) |
|--|----------------|--------------------|------------------------------|
| $W \rightarrow l\nu$ | Inclusive | 37509.0 | 34133.2 |
| $W \rightarrow l\nu$ | 150 - 200 | 253.8 | 234.53 |
| $W \rightarrow l\nu$ | 200 - 250 | 116.5 | 103.94 |
| $W \rightarrow l\nu$ | 250 - 300 | 57.6 | 51.34 |
| $W \rightarrow l\nu$ | 300 - 400 | 48.4 | 42.41 |
| $W \rightarrow l\nu$ | 400 - ∞ | 30.8 | 26.36 |
| $Z \rightarrow \nu\bar{\nu}$ | 50 - 100 | 452.8 | 405.21 |
| $Z \rightarrow \nu\bar{\nu}$ | 100 - 200 | 190.4 | 173.76 |
| $Z \rightarrow \nu\bar{\nu}$ | 200 - 400 | 45.1 | 42.41 |
| $Z \rightarrow \nu\bar{\nu}$ | 400 - ∞ | 6.26 | 5.81 |
| $t\bar{t}$ | Inclusive | 234.0 | 271.44 |
| $Z/\gamma^* \rightarrow l^+l^- (m_{ll} > 50)$ | Inclusive | 3503.7 | 3258.45 |
| $Z/\gamma^* \rightarrow l^+l^- (10 < m_{ll} < 50)$ | Inclusive | 13124.1 | 12205.4 |
| $Z/\gamma^* \rightarrow l^+l^-$ | 200 - 400 | 24.3 | 22.24 |
| $Z/\gamma^* \rightarrow l^+l^-$ | 400 - ∞ | 3.36 | 3.11 |
| $\gamma + \text{jets}$ | 200 - 400 | 1140.8 | 1060.9 |
| $\gamma + \text{jets}$ | 400 - ∞ | 124.7 | 115.97 |
| WW | Inclusive | 57.1 | 57.1 |
| WZ | Inclusive | 12.6 | 12.6 |
| ZZ | Inclusive | 8.26 | 8.26 |
| t (t-channel) | Inclusive | 56.4 | 56.4 |
| \bar{t} (t-channel) | Inclusive | 30.7 | 30.7 |
| t (s-channel) | Inclusive | 3.79 | 3.79 |
| \bar{t} (s-channel) | Inclusive | 1.76 | 1.76 |
| t (tW-channel) | Inclusive | 11.1 | 11.1 |
| \bar{t} (tW-channel) | Inclusive | 11.1 | 11.1 |

0.4.3 Corrections to cross sections for SM samples

A simulated event is weighted by the total number of events in the MC sample, the theoretical cross section, and total luminosity of the data being studied. The MC POG provides next-to-next-to leading order (NNLO) theoretical cross section for un-filtered (inclusive) SM samples [?]. In an attempt to provide higher statistics in tails of distribution analyses often cut on (ie H_T^{parton} , N^{parton} , \hat{p}_T), MC samples are provided binned in these variables. Only the leading-order (LO) cross sections are provided [?] but in general, the k -factors required to go from LO to NNLO cross sections are determined using corresponding inclusive samples and applied to each binned sample.

Studies conducted by other analyses [?] revealed that some LO cross sections calculated for MC samples binned according to H_T^{parton} are inaccurate to a level as large as 10%, leading to non-physical discontinuities in the H_T^{parton} distribution constructed from the binned samples of a


given process. Furthermore, due to an error in production, the $W \rightarrow l\nu$ H_T binned samples exhibit H_T -dependent biases. The following paragraphs describes a procedure to measure corrections to the discontinuities and biases of the $W \rightarrow l\nu$ H_T^{parton} binned samples as a function of H_T . Other analyses created similar procedures to correct the $Z \rightarrow \mu\mu$ H_T^{parton} binned samples, this analysis uses those measured corrections as k-factors.

As a first step, we wish to reweigh the cross sections of the H_T^{parton} binned samples such that their H_T^{parton} distributions match that of the inclusive sample's. Due to the inclusive sample's limited statistics in the tails of H_T distribution, we instead use the N^{parton} binned's H_T^{parton} distribution which is verified to agree well with the inclusive sample figure 1 (a). Additionally, to smooth statistical fluctuations, both distributions are fitted using a double exponential of the form $\exp(a + b * x + c * x^{1.05})$ for $H_T > 500$ GeV figure 1 (b). The ratio of the distributions figure 1(c) is applied as H_T^{parton} dependent event weight.


In the high- H_T and high- E_T corner of kinematic phase space of this analysis (and other SUSY analyses) the overall normalization of MC samples do not agree well with data. Therefore a data sideband in H_T is used to determine sample-specific corrections that are appropriate for the H_T - E_T phase space covered by this analysis. This correction is determined for the $W \rightarrow l\nu$ and $t\bar{t}$ samples by imposing requirements on the number of muons, jets, and b-tagged jets, to obtain samples rich in W + jets, and $t\bar{t}$ events. A sideband in H_T is used to determine both the yields in data and MC expectations. The sideband is defined by the region $200 < H_T < 225$ GeV and uses the jet p_T thresholds (73, 73, 37 GeV) to maintain comparable jet multiplicities, kinematics, and background admixtures as observed for the higher H_T bins. Trigger efficiency and b-tag scale factor corrections are determined and applied to the MC samples. The purity of the samples are $> 80\%$ and any contamination is taken into account. The correction is determined by taking the ratio of the data yield over the MC expectation in the sideband. Table 3 summarises the selection and corrections for the different samples.

Table 3: Correctins determined from a data sideband for the W + jets and $t\bar{t}$ samples. “Corrected yield” reflects the observed data yield minus the contamination as given by MC.


| Process | Selection | Purity | Corrected yield | MC expectation | Correction Factor |
|------------|--|--------|-----------------|--------------------|-------------------|
| W + jets | $\mu + \text{jets}, 2 \leq n_{\text{jet}} \leq 3, n_b = 0$ | 0.90 | 15682 | 18013.1 ± 85.9 | 0.87 ± 0.01 |
| $t\bar{t}$ | $\mu + \text{jets}, n_{\text{jet}} \geq 2, n_b \geq 2$ | 0.83 | 752 | 736.7 ± 11.5 | 1.02 ± 0.05 |



(a) Parton H_T distribution for the $W \rightarrow l\nu$ inclusive sample (blue) and $W \rightarrow l\nu N^{\text{parton}}$ binned sample (orange.)



(b) Fitted Parton H_T distribution for the $W \rightarrow l\nu H_T^{\text{parton}}$ sample (purple) and $W \rightarrow l\nu N^{\text{parton}}$ binned sample (orange.)



(c) Event weight determined from ration of figure 1 (b)

Figure 1: Generator-level H_T^{parton} distributions and measured weights

0.5 Hadronic Event Selection

As discussed in [REF], all-hadronic SUSY signatures consist of events with no isolated and detectable objects apart from energetic jets. To search for an excess in such events, it is convenient to use a jet-based variable, H_T , to quantify the energy in an event. H_T is defined as the scalar sum of the transverse energy of the jets in the event. The challenges due to large backgrounds in a search for an excess in all-hadronic events becomes evident when comparing observed data events overlaid with simulated events from SM processes as a function of H_T Figure [REF].

The dominant background are azimuthally-balanced multi-jet events, stemming from QCD processes. This search reduces this background to negligible levels in the selected region by employing the α_T variable further discussed in section [REF]. Furthermore, by construction, α_T also reduces backgrounds from severely mis-measured jets.

In absence of the multi-jet background, the remaining significant backgrounds are ...

0.5.1 Data

The data analyzed was recorded by CMS in 2012 between April 5th and Dec. 5th and totals $19.47 \pm 2.6\%$. The collected data is certified on a run-by-run basis, where initial automatic certification requires the LHC beams to be declared stable and all CMS subdetectors ON. Further monitoring of the data was done realtime by experts of each subdetector through analysis of histograms updated and filled each lumi section. Final certification was done offline, and lumi sections passing all criteria were listed in a Golden JSON file to be used by all analyses.

0.5.2 Event quality

Each event is subjected to a series of commonly used filters in CMS to ensure good quality data. Minimal requirements are that at least one primary vertex is identified and 25% of the reconstructed tracks to be of good quality. Additionally, various filters prescribed by the MET group[REF] are

applied. Events containing muons with inconsistent energy are flagged by the muon pog [REF] and filtered out of the analysis.

0.5.3 Triggers

Hadronic signal region and control samples

Only events passing one or more HLT triggers based on online quantities are recorded to be analyzed. For any analysis, in general, it is not expected that all recorded events reconstructed offline, pass the online trigger as detector conditions, energy corrections, and object-based quantities differ offline. In this analysis, cross-triggers at the HLT based on quantities H_T and α_T (labelled as HTxxx_AlphaT0pyy) are used with various thresholds to record candidate events for the hadronic (signal) region. In order to keep the cross-trigger's computational time low, the online quantities are constructed using simple calorimeter based jets (calo jets), and the use of particle-flow jets in this analysis is expected to introduce inefficiencies.

Each H_T bin is seeded by a single trigger chosen based on the efficiency of the trigger in that H_T bin. The α_T thresholds of the HTxxx_AlphaT0pyy triggers were tuned according to the threshold on the H_T leg in order to fully suppress QCD multijet events and simultaneously satisfying other criteria, such as maintaining acceptable trigger rates.

The HTxxx_AlphaT0pyy trigger efficiencies are measured with a reference (i.e., unbiased) event sample recorded by an unprescaled, loosely-isolated, eta-restricted single muon trigger, HLT_IsoMu24_eta2p1, within the SingleMu dataset. A sample of events containing at least one isolated muon with $p_T > 25 \text{ GeV}$ and $|\eta| < 2.1$ is used (similar to the $\mu + \text{jets}$ control sample defined in Section ??). A cut of $\Delta R > 0.5$ is placed between all muons and jets in each event, and only jets are considered in the calculation of H_T , H_T , and α_T , i.e. the muon is ignored.







(a) Differential, $375 < H_T < 475$ GeV(b) Cumulative, $375 < H_T < 475$ GeV(c) Differential, $475 < H_T < 525$ GeV(d) Cumulative, $475 < H_T < 525$ GeV(e) Differential, $525 < H_T < 675$ GeV(f) Cumulative, $525 < H_T < 675$ GeV

Figure 2: (Left) Differential and (Right) cumulative efficiency turn-on curves for the H_T - α_T cross triggers (as summarised in Table ??) that record events for the three lowest H_T bins for events satisfying $2 \leq n_{\text{jet}} \leq 3$.

Muon control samples

0.5.4 event selection







(a) Differential, $375 < H_T < 475$ GeV(b) Cumulative, $375 < H_T < 475$ GeV(c) Differential, $475 < H_T < 525$ GeV(d) Cumulative, $475 < H_T < 525$ GeV(e) Differential, $525 < H_T < 675$ GeV(f) Cumulative, $525 < H_T < 675$ GeV

Figure 3: (Left) Differential and (Right) cumulative efficiency turn-on curves for the H_T - α_T cross triggers (as summarised in Table ??) that record events for the three lowest H_T bins for events satisfying $n_{\text{jet}} \geq 4$.

0.6 Closure tests and systematic uncertainties on transfer factors

Limitations in simulating detector effects and event kinematics requires us to apply appropriate systematics uncertainties on the simulation-based translation factors. The following section describes how we obtain these systematic uncertainty through the method of closure tests.

0.6.1 Closure tests

At its core, the method compares an observed yield (N_{obs}) and a predicted yield (N_{pred}) in a subsample of a control region. The predicted yield is constructed by translating from a statistically independent data sample to the data sample of interest by the use of the proper translation factor. For example, for a given H_T bin, a prediction for the $n_{\text{jet}} \geq 4$, $n_b = 1$, $\mu + \text{jets}$ sample can be made by translating from the $2 \leq n_{\text{jet}} \leq 3$, $n_b = 1$, $\mu + \text{jets}$ in data via the translation factor:

$$\frac{N_{\text{MC}}^{\mu+\text{jets}}(H_T, n_{\text{jet}} \geq 4, n_b = 1)}{N_{\text{MC}}^{\mu+\text{jets}}(H_T, 2 \leq n_{\text{jet}} \leq 3, n_b = 1)} \quad (1)$$

The agreement between N_{obs} and N_{pred} is expressed as $(N_{\text{obs}} - N_{\text{pred}})/N_{\text{pred}}$. Assuming only statistical uncertainties on N_{obs} and N_{pred} , deviation of the ratio from zero defines our level of closure. A closure test set is defined as ratios for each H_T bin. Looking at the ratio as a function of H_T allows the measurement of statistical significant biases from zero and/or any dependence on H_T . If statistically significant biases are observed, further studies are required to understand and correct for these biases.

Eight sets of closure tests probe key ingredients of the simulation modelling of the SM backgrounds with genuine E_T as a function of H_T , as shown in Fig. 4. This is done for the two jet multiplicity bins separately: (a) $2 \leq n_{\text{jet}} \leq 3$ and (b) $n_{\text{jet}} \geq 4$.

Under the assumption of closure for the full ensemble of tests, systematic uncertainties on the transfer factors are derived for each n_{jet} category and H_T regions. The treatment for estimating the

systematic uncertainties on the transfer factors is described in Section 0.6.2.

As described in section ?? The α_T requirement is not imposed in the $\mu + \text{jets}$ control sample. Therefore it is important to verify the approach of using $\mu + \text{jets}$ samples without an α_T requirement to make background predictions in the signal region. The first set of closure tests (denoted by circles) attempts to do this by probing the modelling of the α_T distribution in genuine E_T events as a function of H_T . The tests compares data yields in the $\mu + \text{jets}$ sample with an α_T requirement against predictions determined in a $\mu + \text{jets}$ sample with the α_T requirement inverted.

The next three sets (triangles, crosses, squares) probe the sensitivity of the transfer factors to the relative admixture of events from the $W + \text{jets}$ and $t\bar{t}$ processes. These tests are conservative, since by construction, the admixture changes little when translating from the $\mu + \text{jets}$ control region to the signal region, whereas the closure tests use sub-samples with different b-tag requirements and therefore have very different admixtures of $W + \text{jets}$ and $t\bar{t}$ events. In the $2 \leq n_{\text{jet}} \leq 3$ bin, the test is sub-divided into separate jet categories. These tests also probe the modelling of the reconstruction of b-quark jets, although this also addressed more fully by dedicated studies that determine systematic uncertainties via the method described in Sec. ??.

The remaining tests probe the simulation modelling of the jet multiplicity in the $\mu + \text{jets}$ and $\gamma + \text{jets}$ samples, which is checked due to the exclusive binning in jet multiplicity. As in the case of the $W + \text{jets} / t\bar{t}$ admixture, this set of tests is a very conservative check, as predictions are always made from the same jet multiplicity bin, whereas the closure tests translate between the two bins.

Tables 4 and 5, which summarize the results obtained from fits of zeroeth order polynomials (i.e. a constant) to the sets of closure tests performed in the $2 \leq n_{\text{jet}} \leq 3$ and $n_{\text{jet}} \geq 4$ bins. Table 6 lists the fits result common to both jet multiplicities. The best fit value and its uncertainty is listed for each set of closure tests, along with the χ^2 , the number of degrees of freedom, and the p-value of the fit. The best fit value for the constant parameter is indicative of the level of closure, as averaged across the full H_T range considered in the analysis, and the p-value is indicative of

whether there is any significant dependence on H_T .

The closure tests demonstrate, within the statistical precision of each test, that there are no significant biases or dependencies on H_T inherent in the transfer factors obtained from simulation.

One set of tests does indicate a poor goodness of fit (indicated by a low p -value), which is the $n_b = 0 \rightarrow n_b = 1$ test in the $\mu + \text{jets}$ sample for the $n_{\text{jet}} \geq 4$ category, which has been identified as a upward (downward) fluctuation of event counts in the H_T bin 475–575 GeV (575–675 GeV) when $n_b = 1$. Combining these two bins yields an acceptable fit result, as indicated in Table 5, which points to a simple fluctuation rather than any systematic bias.

In addition to the fits described above, linear fits are also performed. The best fit values for the slope terms and the p -values obtained from each fit are summarised in Tables 4, 5, and 6. Typically, the best fit values are of the order 10^{-4} , which corresponds to a percent-level change per 100 GeV.

Table 4: A summary of the results obtained from fits of zeroeth order polynomials (i.e. a constant) to four sets of closure tests performed in the $2 \leq n_{\text{jet}} \leq 3$ bin.

| Closure test | Symbol | Constant fit | | | |
|---|----------|-------------------|----------|--------|------------|
| | | Best fit value | χ^2 | d.o.f. | p -value |
| $\alpha_T < 0.55 \rightarrow \alpha_T > 0.55$ ($\mu + \text{jets}$) | Circle | 0.007 ± 0.02 | 3.91 | 7 | 0.79 |
| 1 b-tags \rightarrow 2 b-tags ($\mu + \text{jets}$, nJet=3) | Triangle | -0.008 ± 0.04 | 3.20 | 7 | 0.87 |
| 0 b-tags \rightarrow 1 b-tags ($\mu + \text{jets}$, nJet=2) | Cross | 0.111 ± 0.03 | 5.87 | 7 | 0.55 |
| 0 b-tags \rightarrow 1 b-tags ($\mu + \text{jets}$, nJet=3) | Square | 0.040 ± 0.02 | 1.12 | 7 | 0.99 |

Table 5: A summary of the results obtained from fits of zeroeth order polynomials (i.e. a constant) to three sets of closure tests performed in the $n_{\text{jet}} \geq 4$ bin. [†]Further explanation of this fit can be found in the text.

| Closure test | Symbol | Constant fit | | | |
|---|----------|------------------|----------|--------|------------|
| | | Best fit value | χ^2 | d.o.f. | p -value |
| $\alpha_T < 0.55 \rightarrow \alpha_T > 0.55$ ($\mu + \text{jets}$) | Circle | 0.011 ± 0.04 | 5.81 | 7 | 0.56 |
| 1 b-tags \rightarrow 2 b-tags ($\mu + \text{jets}$) | Triangle | 0.045 ± 0.03 | 9.36 | 7 | 0.23 |
| 0 b-tags \rightarrow 1 b-tags ($\mu + \text{jets}$) | Square | 0.007 ± 0.03 | 25.30 | 7 | 0.00 |
| 0 b-tags \rightarrow 1 b-tags ($\mu + \text{jets}$) [†] | Square | 0.009 ± 0.03 | 10.12 | 6 | 0.12 |

Table 6: A summary of the results obtained from fits of zeroeth order polynomials (i.e. a constant) to four sets of closure tests ($2 \leq n_{\text{jet}} \leq 3 \rightarrow n_{\text{jet}} \geq 4$) that probe the accuracy of the MC modelling of the n_{jet} distribution observed in data, using the three data control samples.

| Closure test | Symbol | Constant fit | | | |
|---|------------------|-------------------|----------|--------|------------|
| | | Best fit value | χ^2 | d.o.f. | p -value |
| $2 \leq n_{\text{jet}} \leq 3 \rightarrow n_{\text{jet}} \geq 4$ (μ + jets, 1 b-tags) | Times | -0.053 ± 0.03 | 8.02 | 7 | 0.33 |
| $2 \leq n_{\text{jet}} \leq 3 \rightarrow n_{\text{jet}} \geq 4$ (μ + jets, 1 b-tags) | Invert. Triangle | 0.018 ± 0.04 | 6.23 | 7 | 0.51 |
| $2 \leq n_{\text{jet}} \leq 3 \rightarrow n_{\text{jet}} \geq 4$ (μ + jets, 0 b-tags) | Star | 0.034 ± 0.02 | 9.24 | 7 | 0.24 |
| $2 \leq n_{\text{jet}} \leq 3 \rightarrow n_{\text{jet}} \geq 4$ (γ + jets, 0 b-tags) | Diamond | 0.100 ± 0.04 | 12.20 | 7 | 0.09 |

0.6.2 Systematic uncertainties from closure tests

Once it is established that no significantly large bias or trend is observed for any set of closure tests, then systematic uncertainties are determined.

Systematics are determined for each H_T bin, as indicated in Table 7. For each H_T region, the systematic uncertainty is estimated by taking the quadrature sum of the weighted mean and sample variance for the closure tests within the given H_T region. This procedure yields the values quoted in Table 7.

The effect of uncertainties related to the modelling of b-quark jets in simulation on the transfer factors is found to be negligible, at the percent level as discussed in Section ??, in comparison to the aforementioned n_{jet} - and H_T -dependent systematic uncertainties.

Table 7: A summary of the magnitude of the systematic uncertainties (%) assigned to the transfer factors, according to n_{jet} and H_T region.

| n_{jet} | H_T region (GeV) | | | | | | | |
|------------------|--------------------|---------|---------|---------|---------|---------|-----------|--------|
| | 375–475 | 475–525 | 525–675 | 675–775 | 775–875 | 875–975 | 1075–1075 | > 1175 |
| 2–3 | 6 | 6 | 7 | 12 | 12 | 17 | 17 | 17 |
| ≥ 4 | 6 | 6 | 8 | 14 | 14 | 14 | 14 | 21 |

Figure 4 shows the sets of closure tests overlaid on top of grey bands that represent the H_T -dependent systematic uncertainties in Table 7. These systematic uncertainties are assumed to fully

uncorrelated between the different b jet multiplicity categories and also the seven H_T regions, which is a conservative approach given that one can expect some correlation between adjacent H_T bins (due to comparable kinematics). This approach of decorrelating the H_T regions should be contrasted against the fits shown in Figure ?? that do assume a correlated behaviour in H_T .



(a) $2 \leq n_{\text{jet}} \leq 3$ (b) $n_{\text{jet}} \geq 4$

Figure 4: Sets of closure tests (open symbols) overlaid on top of the systematic uncertainty used for each of the H_T region (shaded bands) and for the two different jet multiplicity bins: (a) $2 \leq n_{\text{jet}} \leq 3$ and (b) $n_{\text{jet}} \geq 4$.

0.7 Results

The likelihood described in ref:likelihood is used to relate yields, uncertainties. It is constructed using ROOTFIT [?] and maximized using MINUIT [?].

0.7.1 Standard Model


To test compatibility with a Standard Model only hypothesis, the signal term is removed from the likelihood model. The parameter values maximizing the likelihood function are listed in Tables 10–15 found in Appendix .1. The resulting SM yields along with the observed data yields are summarized in Tables .2. The uncertainty on the yields are obtained by constructing a probability density function (p.d.f) from the maximized likelihood, then generating an ensemble of pseudo-experiments from this p.d.f. and maximizing the same likelihood form for each pseudo-experiment, resulting in an ensemble of yields. The 68% quantile of each ensemble defines the quoted uncertainty on the corresponding yield.

Figures 6–11 show the H_T -binned observed data yields (black filled circles) and the SM expectations and uncertainties (dark blue solid line with light blue bands) as determined by the fit for the hadronic signal region and the $\mu + \text{jets}$ or both ($\mu + \text{jets}, \gamma + \text{jets}$) control samples, depending on the event category. The uncertainties in the SM expectations obtained from the ensemble of pseudo-experiments reflect the statistical uncertainties in the considered data samples and the systematic uncertainties in the transfer factors as discussed in section 0.6. Figures 6–11 are summarized in tabular format in Tables 16–21 in appendix .2 along with observed data yields and the fit result for all event categories and both signal region and control sample bins.


For each n_b, n_{jet} category, the goodness-of-fit of the SM-only hypothesis is determined by considering simultaneously all H_T bins entering the likelihood. The goodness-of-fit described in [?] is obtained by comparing the nominal maximized likelihood value $L_{\text{max}}^{\text{data}}$ to the corresponding ensemble of values, L_{max} . The quantile which $L_{\text{max}}^{\text{data}}$ falls in the distributions is interpreted as a

p-value. A p-value derived from a chi-square is also plotted for comparison.

The p-values obtained, shown in Figure 5 (Left), are found to be uniformly distributed in the range 0.0–1.0, with the lowest p-value determined to be 0.17.



(a) p-values per event category.



(b) Pull versus signal region bin.

Figure 5: Pulls and p-values. See text for details

Table 8: Summary of hadronic yields from fit.

| | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075–∞ |
|--------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|---------------------|---------------------|
| 0b le3j SM | 2637^{+50}_{-48} | 759^{+24}_{-23} | 252^{+14}_{-13} | $76.5^{+6.6}_{-4.7}$ | $33.7^{+3.5}_{-3.7}$ | $11.8^{+2.1}_{-2.3}$ | $6.3^{+1.4}_{-1.2}$ | $3.2^{+0.9}_{-0.9}$ |
| 0b le3j Data | 2627 | 762 | 253 | 77 | 32 | 9 | 9 | 4 |
| 1b le3j SM | 413^{+15}_{-14} | 111^{+6}_{-4} | $35.8^{+3.4}_{-2.7}$ | $10.1^{+1.6}_{-1.3}$ | $3.7^{+0.8}_{-0.8}$ | $1.6^{+0.6}_{-0.7}$ | $0.5^{+0.3}_{-0.4}$ | $0.1^{+0.1}_{-0.0}$ |
| 1b le3j Data | 440 | 116 | 37 | 15 | 3 | 2 | 1 | 0 |
| 2b le3j SM | $63.0^{+3.9}_{-4.1}$ | $18.0^{+1.3}_{-1.4}$ | $4.2^{+0.6}_{-0.5}$ | $1.1^{+0.2}_{-0.2}$ | $0.2^{+0.1}_{-0.1}$ | $0.0^{+0.0}_{-0.0}$ | $0.0^{+0.0}_{-0.0}$ | $0.0^{+0.0}_{-0.0}$ |
| 2b le3j Data | 80 | 19 | 8 | 3 | 0 | 0 | 0 | 0 |
| 0b ge4j SM | 460^{+16}_{-15} | 298^{+12}_{-11} | 146^{+7}_{-7} | $66.0^{+4.4}_{-5.5}$ | $27.1^{+3.2}_{-3.5}$ | $14.0^{+2.0}_{-1.9}$ | $6.5^{+1.4}_{-1.5}$ | $3.2^{+1.0}_{-1.0}$ |
| 0b ge4j Data | 470 | 302 | 158 | 66 | 28 | 15 | 6 | 2 |
| 1b ge4j SM | 192^{+9}_{-8} | 122^{+6}_{-6} | $44.8^{+3.2}_{-3.7}$ | $17.1^{+2.4}_{-2.0}$ | $6.8^{+1.2}_{-1.3}$ | $5.4^{+1.5}_{-1.5}$ | $2.4^{+1.0}_{-0.8}$ | $1.2^{+0.8}_{-0.8}$ |
| 1b ge4j Data | 196 | 132 | 44 | 14 | 8 | 6 | 2 | 0 |
| 2b ge4j SM | $74.2^{+4.2}_{-4.4}$ | $47.0^{+3.2}_{-3.3}$ | $20.1^{+2.0}_{-1.8}$ | $7.7^{+1.3}_{-1.1}$ | $1.9^{+0.3}_{-0.3}$ | $0.9^{+0.2}_{-0.2}$ | $0.4^{+0.1}_{-0.1}$ | $0.4^{+0.2}_{-0.1}$ |
| 2b ge4j Data | 72 | 52 | 31 | 12 | 1 | 2 | 0 | 1 |




Figure 6: Comparison of the H_T -binned observed data yields and SM expectations when requiring $2 \leq n_{\text{jet}} \leq 3$ and $n_b = 0$ for the (a-b) hadronic, (c) μ + jets, (d) $\mu\mu$ + jets and (e) γ + jets samples, as determined by a simultaneous fit to all data samples under the SM-only hypothesis. The observed event yields in data (black dots) and the expectations and their uncertainties (dark blue solid line with light blue bands), as determined by the simultaneous fit, are shown. For illustrative purposes only, the signal expectations (pink dashed line) for the model T2cc with $m_{\tilde{q}} = 250$ GeV and $m_{\text{LSP}} = 240$ GeV are stacked on top of the SM expectations.




Figure 7: Comparison of the H_T -binned observed data yields and SM expectations when requiring $2 \leq n_{jet} \leq 3$ and $n_b = 1$ for the (a-b) hadronic, (c) μ + jets, (d) $\mu\mu$ + jets and (e) γ + jets samples, as determined by a simultaneous fit to all data samples under the SM-only hypothesis. The observed event yields in data (black dots) and the expectations and their uncertainties (dark blue solid line with light blue bands), as determined by the simultaneous fit, are shown. For illustrative purposes only, the signal expectations (pink dashed line) for the model T2cc with $m_{\tilde{q}} = 250$ GeV and $m_{LSP} = 170$ GeV are stacked on top of the SM expectations.




Figure 8: Comparison of the H_T -binned observed data yields and SM expectations when requiring $2 \leq n_{\text{jet}} \leq 3$ and $n_b = 2$ for the (a-b) hadronic and $\mu + \text{jets}$ samples, as determined by a simultaneous fit to both the hadronic and $\mu + \text{jets}$ data samples under the SM-only hypothesis. The observed event yields in data (black dots) and the expectations and their uncertainties (dark blue solid line with light blue bands), as determined by the simultaneous fit, are shown.




Figure 9: Comparison of the H_T -binned observed data yields and SM expectations when requiring $n_{\text{jet}} \geq 4$ and $n_b = 0$ for the (a-b) hadronic, (c) μ + jets, (d) $\mu\mu$ + jets and (e) γ + jets samples, as determined by a simultaneous fit to all data samples under the SM-only hypothesis. The observed event yields in data (black dots) and the expectations and their uncertainties (dark blue solid line with light blue bands), as determined by the simultaneous fit, are shown. For illustrative purposes only, the signal expectations (pink dashed line) for the model T2cc with $m_{\tilde{q}} = 250$ GeV and $m_{\text{LSP}} = 170$ GeV are stacked on top of the SM expectations.




Figure 10: Comparison of the H_T -binned observed data yields and SM expectations when requiring $n_{\text{jet}} \geq 4$ and $n_b = 1$ for the (a-b) hadronic, (c) μ + jets, (d) $\mu\mu$ + jets and (e) γ + jets samples, as determined by a simultaneous fit to all data samples under the SM-only hypothesis. The observed event yields in data (black dots) and the expectations and their uncertainties (dark blue solid line with light blue bands), as determined by the simultaneous fit, are shown. For illustrative purposes only, the signal expectations (pink dashed line) for the model T2cc with $m_{\tilde{q}} = 250$ GeV and $m_{\text{LSP}} = 170$ GeV are stacked on top of the SM expectations.




Figure 11: Comparison of the H_T -binned observed data yields and SM expectations when requiring $n_{\text{jet}} \geq 4$ and $n_b = 2$ for the (a-b) hadronic and μ + jets samples, as determined by a simultaneous fit to both the hadronic and μ + jets data samples under the SM-only hypothesis. The observed event yields in data (black dots) and the expectations and their uncertainties (dark blue solid line with light blue bands), as determined by the simultaneous fit, are shown.

0.8 Signal models and efficiencies

fill

0.8.1 Introduction

As mentioned in section ??, SUSY is a theory of many particles and parameters. Simplified models (SMS) have been derived [?] which reduce both particles and parameters at the cost of losing generality of the full SUSY model. Yet, SMS models are beneficial experimentally, as they provide clear signatures to search for. Two SMS models have been studied 1. T2cc: pair produced stop sparticles each decaying into a charm quark and a neutralino. 2. T2tt: pair produced stop sparticles each decaying into a top quark and a neutralino.

Event samples for the simplified models are generated by the CMS SUSY MC group [REF] at leading order with MADGRAPH [?]. Inclusive, process-dependent, next-to-leading order calculations with next-to-leading logarithmic corrections [?] (NLO+NLL) of SUSY production cross sections are obtained with the program PROSPINO [?] and CTEQ6L [?] parton distribution functions. The FastSim framework is used, and the simulated signal events include multiple interactions per LHC bunch crossing (pileup) with the distribution of reconstructed vertices that match the one observed in data. The MC samples are listed in Section 0.4.

The SMS samples are produced in bins of stop mass (m_{Stop}) and neutralino mass (m_{LSP}). The analysis efficiency is studied in the usual n_{jet}, n_b binning but due to computational limitations not all categories are used to set a limit on a given model. The choice of categories for a model is made by computing the expected upper limit on the signal cross-section for each category separately using the much quicker asymptotic method [REF]. The categories are ranked by their expected upper limit. Depending on the number of mass bins and number of events per bin, two or more categories with the highest rank are chosen.

The simplified models, along with the event categories considered for each, is summarised in

Table 9.

Table 9: A summary of the simplified models considered for interpretation. The event categories considered for each model are listed.

| Model | Production/decay mode | $(n_{\text{jet}}, n_{\text{b}})$ event categories considered |
|-------|---|--|
| T2cc | $pp \rightarrow t\bar{t}^* \rightarrow c\tilde{\chi}^0 \bar{c}\tilde{\chi}^0$ | $(2-3,0), (\geq 4,0), (\geq 4,1)$ |
| T2tt | $pp \rightarrow t\bar{t}^* \rightarrow t\tilde{\chi}^0 \bar{t}\tilde{\chi}^0$ | $(\leq 3,1), (\leq 4,2)$ |

0.8.2 Efficiency times acceptance for T2cc

Figure 12 (Appendix .3.1) shows the expected signal efficiency times acceptance for the signal region and μ + jets control sample (i.e., signal contamination) for the model T2cc in the four most sensitive $(n_{\text{jet}}, n_{\text{b}})$ event categories: $(2-3,0)$, $(2-3,1)$, $(\geq 4,0)$, and $(\geq 4,1)$. The choice of which categories to use is made by inspecting Figure ?? (Appendix .3.1), which shows the expected significance per signal region bin following the injection of signal at the theoretical QCD production cross section for the T2cc model mass points $m_{\tilde{t}} = 250$ GeV and $m_{\text{LSP}} = 170$ GeV (Top) and $m_{\text{LSP}} = 240$ GeV (Bottom).

The efficiencies are typically at the percent level or less due to reliance on hard- p_{T} jets from initial state radiation for acceptance in the presence of a compressed mass spectrum and soft decay products. The largest efficiencies for the smallest mass splittings, $\Delta M = \sim 10$ GeV, are obtained with the $(2-3,0)$ category, while for larger mass splittings the $(2-3,1)$, $(\geq 4,0)$, and $(\geq 4,1)$ categories contribute due to the reduced backgrounds in these categories. The signal efficiency in the μ + jets control sample is negligible with respect to the signal region. By extension, the relative contamination for the $\mu\mu$ + jets sample is also considered to be negligible. Regardless, any potential contamination is accounted for in the likelihood model.

.1 Maximum likelihood parameter values

Table 10: SM-only maximum-likelihood parameter values (0b 1e3j).

| name | value | error |
|---------------------|----------|---------|
| EWK^0 | 2.64e+03 | 4.7e+01 |
| EWK^1 | 7.59e+02 | 2.2e+01 |
| EWK^2 | 2.52e+02 | 1.1e+01 |
| EWK^3 | 7.64e+01 | 6.0e+00 |
| EWK^4 | 3.37e+01 | 3.5e+00 |
| EWK^5 | 1.18e+01 | 2.0e+00 |
| EWK^6 | 6.32e+00 | 1.4e+00 |
| EWK^7 | 3.15e+00 | 8.9e-01 |
| f_{Zinv}^0 | 0.64 | 0.02 |
| f_{Zinv}^1 | 0.67 | 0.02 |
| f_{Zinv}^2 | 0.70 | 0.02 |
| f_{Zinv}^3 | 0.70 | 0.04 |
| f_{Zinv}^4 | 0.69 | 0.04 |
| f_{Zinv}^5 | 0.76 | 0.05 |
| f_{Zinv}^6 | 0.76 | 0.06 |
| f_{Zinv}^7 | 0.82 | 0.06 |
| $\rho_{\mu W}^0$ | 1.01 | 0.05 |
| $\rho_{\mu W}^1$ | 1.00 | 0.06 |
| $\rho_{\mu W}^2$ | 1.00 | 0.07 |
| $\rho_{\mu W}^3$ | 1.00 | 0.11 |
| $\rho_{\mu W}^4$ | 1.01 | 0.11 |
| $\rho_{\mu W}^5$ | 1.02 | 0.16 |
| $\rho_{\mu W}^6$ | 0.98 | 0.15 |
| $\rho_{\mu W}^7$ | 1.00 | 0.16 |
| $\rho_{\gamma Z}^0$ | 1.02 | 0.04 |
| $\rho_{\gamma Z}^1$ | 0.99 | 0.04 |
| $\rho_{\gamma Z}^2$ | 1.00 | 0.06 |
| $\rho_{\gamma Z}^3$ | 0.99 | 0.10 |
| $\rho_{\gamma Z}^4$ | 1.02 | 0.11 |
| $\rho_{\gamma Z}^5$ | 1.05 | 0.16 |
| $\rho_{\gamma Z}^6$ | 0.95 | 0.14 |
| $\rho_{\gamma Z}^7$ | 0.98 | 0.15 |

Table 11: SM-only maximum-likelihood parameter values (1b le3j).

| name | value | error |
|---------------------|----------|---------|
| EWK^0 | 4.13e+02 | 1.4e+01 |
| EWK^1 | 1.11e+02 | 5.9e+00 |
| EWK^2 | 3.58e+01 | 3.0e+00 |
| EWK^3 | 1.01e+01 | 1.6e+00 |
| EWK^4 | 3.66e+00 | 8.2e-01 |
| EWK^5 | 1.63e+00 | 6.0e-01 |
| EWK^6 | 4.87e-01 | 3.1e-01 |
| EWK^7 | 1.21e-01 | 4.7e-02 |
| f_{Zinv}^0 | 0.47 | 0.02 |
| f_{Zinv}^1 | 0.55 | 0.03 |
| f_{Zinv}^2 | 0.57 | 0.04 |
| f_{Zinv}^3 | 0.64 | 0.07 |
| f_{Zinv}^4 | 0.55 | 0.11 |
| f_{Zinv}^5 | 0.81 | 0.08 |
| f_{Zinv}^6 | 0.82 | 0.12 |
| f_{Zinv}^7 | 0.00 | 0.83 |
| $\rho_{\mu W}^0$ | 0.95 | 0.05 |
| $\rho_{\mu W}^1$ | 0.99 | 0.06 |
| $\rho_{\mu W}^2$ | 1.00 | 0.07 |
| $\rho_{\mu W}^3$ | 0.98 | 0.11 |
| $\rho_{\mu W}^4$ | 1.00 | 0.11 |
| $\rho_{\mu W}^5$ | 1.00 | 0.16 |
| $\rho_{\mu W}^6$ | 1.00 | 0.16 |
| $\rho_{\mu W}^7$ | 1.00 | 0.16 |
| $\rho_{\gamma Z}^0$ | 0.96 | 0.05 |
| $\rho_{\gamma Z}^1$ | 0.99 | 0.06 |
| $\rho_{\gamma Z}^2$ | 1.00 | 0.07 |
| $\rho_{\gamma Z}^3$ | 0.96 | 0.11 |
| $\rho_{\gamma Z}^4$ | 1.00 | 0.11 |
| $\rho_{\gamma Z}^5$ | 0.99 | 0.15 |
| $\rho_{\gamma Z}^6$ | 0.99 | 0.15 |
| $\rho_{\gamma Z}^7$ | 1.00 | 0.16 |

Table 12: SM-only maximum-likelihood parameter values (2b 1e3j).

| name | value | error |
|------------------|----------|---------|
| EWK^0 | 6.30e+01 | 3.8e+00 |
| EWK^1 | 1.80e+01 | 1.5e+00 |
| EWK^2 | 4.25e+00 | 5.4e-01 |
| EWK^3 | 1.11e+00 | 2.7e-01 |
| EWK^4 | 2.10e-01 | 5.9e-02 |
| EWK^5 | 3.83e-02 | 1.8e-02 |
| EWK^6 | 2.33e-02 | 1.7e-02 |
| EWK^7 | 1.30e-03 | 1.3e-03 |
| $\rho_{\mu W}^0$ | 0.94 | 0.05 |
| $\rho_{\mu W}^1$ | 1.00 | 0.06 |
| $\rho_{\mu W}^2$ | 0.98 | 0.07 |
| $\rho_{\mu W}^3$ | 0.98 | 0.11 |
| $\rho_{\mu W}^4$ | 1.00 | 0.11 |
| $\rho_{\mu W}^5$ | 1.00 | 0.16 |
| $\rho_{\mu W}^6$ | 1.00 | 0.16 |
| $\rho_{\mu W}^7$ | 1.00 | 0.16 |

Table 13: SM-only maximum-likelihood parameter values (0b ge4j).

| name | value | error |
|---------------------|----------|---------|
| EWK^0 | 4.60e+02 | 1.6e+01 |
| EWK^1 | 2.98e+02 | 1.2e+01 |
| EWK^2 | 1.46e+02 | 8.2e+00 |
| EWK^3 | 6.59e+01 | 5.7e+00 |
| EWK^4 | 2.71e+01 | 3.3e+00 |
| EWK^5 | 1.39e+01 | 2.2e+00 |
| EWK^6 | 6.47e+00 | 1.5e+00 |
| EWK^7 | 3.19e+00 | 9.5e-01 |
| f_{Zinv}^0 | 0.52 | 0.02 |
| f_{Zinv}^1 | 0.60 | 0.02 |
| f_{Zinv}^2 | 0.61 | 0.03 |
| f_{Zinv}^3 | 0.63 | 0.05 |
| f_{Zinv}^4 | 0.71 | 0.05 |
| f_{Zinv}^5 | 0.73 | 0.06 |
| f_{Zinv}^6 | 0.74 | 0.07 |
| f_{Zinv}^7 | 0.67 | 0.11 |
| $\rho_{\mu W}^0$ | 0.98 | 0.05 |
| $\rho_{\mu W}^1$ | 1.00 | 0.06 |
| $\rho_{\mu W}^2$ | 0.97 | 0.07 |
| $\rho_{\mu W}^3$ | 1.00 | 0.13 |
| $\rho_{\mu W}^4$ | 1.00 | 0.13 |
| $\rho_{\mu W}^5$ | 1.00 | 0.13 |
| $\rho_{\mu W}^6$ | 1.00 | 0.13 |
| $\rho_{\mu W}^7$ | 1.01 | 0.19 |
| $\rho_{\gamma Z}^0$ | 0.98 | 0.05 |
| $\rho_{\gamma Z}^1$ | 0.99 | 0.05 |
| $\rho_{\gamma Z}^2$ | 0.96 | 0.07 |
| $\rho_{\gamma Z}^3$ | 1.00 | 0.11 |
| $\rho_{\gamma Z}^4$ | 0.99 | 0.12 |
| $\rho_{\gamma Z}^5$ | 0.99 | 0.12 |
| $\rho_{\gamma Z}^6$ | 1.01 | 0.13 |
| $\rho_{\gamma Z}^7$ | 1.03 | 0.19 |

Table 14: SM-only maximum-likelihood parameter values (1b ge4j).

| name | value | error |
|---------------------|----------|---------|
| EWK^0 | 1.92e+02 | 8.5e+00 |
| EWK^1 | 1.22e+02 | 6.1e+00 |
| EWK^2 | 4.48e+01 | 3.5e+00 |
| EWK^3 | 1.71e+01 | 2.1e+00 |
| EWK^4 | 6.82e+00 | 1.3e+00 |
| EWK^5 | 5.43e+00 | 1.5e+00 |
| EWK^6 | 2.43e+00 | 7.9e-01 |
| EWK^7 | 1.21e+00 | 6.3e-01 |
| f_{Zinv}^0 | 0.23 | 0.02 |
| f_{Zinv}^1 | 0.29 | 0.03 |
| f_{Zinv}^2 | 0.38 | 0.05 |
| f_{Zinv}^3 | 0.36 | 0.08 |
| f_{Zinv}^4 | 0.49 | 0.10 |
| f_{Zinv}^5 | 0.58 | 0.12 |
| f_{Zinv}^6 | 0.70 | 0.11 |
| f_{Zinv}^7 | 0.72 | 0.16 |
| $\rho_{\mu W}^0$ | 0.99 | 0.05 |
| $\rho_{\mu W}^1$ | 0.98 | 0.05 |
| $\rho_{\mu W}^2$ | 1.00 | 0.07 |
| $\rho_{\mu W}^3$ | 1.03 | 0.13 |
| $\rho_{\mu W}^4$ | 0.99 | 0.13 |
| $\rho_{\mu W}^5$ | 1.00 | 0.13 |
| $\rho_{\mu W}^6$ | 1.00 | 0.13 |
| $\rho_{\mu W}^7$ | 1.01 | 0.19 |
| $\rho_{\gamma Z}^0$ | 1.00 | 0.06 |
| $\rho_{\gamma Z}^1$ | 0.99 | 0.06 |
| $\rho_{\gamma Z}^2$ | 1.00 | 0.08 |
| $\rho_{\gamma Z}^3$ | 1.02 | 0.13 |
| $\rho_{\gamma Z}^4$ | 0.99 | 0.13 |
| $\rho_{\gamma Z}^5$ | 0.99 | 0.13 |
| $\rho_{\gamma Z}^6$ | 1.01 | 0.13 |
| $\rho_{\gamma Z}^7$ | 1.03 | 0.19 |

Table 15: SM-only maximum-likelihood parameter values (2b ge4j).

| name | value | error |
|------------------|----------|---------|
| EWK ⁰ | 7.42e+01 | 4.4e+00 |
| EWK ¹ | 4.70e+01 | 3.1e+00 |
| EWK ² | 2.01e+01 | 1.8e+00 |
| EWK ³ | 7.71e+00 | 1.1e+00 |
| EWK ⁴ | 1.85e+00 | 3.4e-01 |
| EWK ⁵ | 9.29e-01 | 2.3e-01 |
| EWK ⁶ | 4.32e-01 | 1.4e-01 |
| EWK ⁷ | 4.09e-01 | 1.5e-01 |
| $\rho_{\mu W}^0$ | 1.01 | 0.05 |
| $\rho_{\mu W}^1$ | 0.98 | 0.05 |
| $\rho_{\mu W}^2$ | 0.94 | 0.07 |
| $\rho_{\mu W}^3$ | 0.93 | 0.11 |
| $\rho_{\mu W}^4$ | 1.01 | 0.13 |
| $\rho_{\mu W}^5$ | 0.98 | 0.13 |
| $\rho_{\mu W}^6$ | 1.01 | 0.13 |
| $\rho_{\mu W}^7$ | 0.98 | 0.18 |

.2 SM-only yield tables

The following tables compare the observations in the hadronic and control samples with the maximum-likelihood expectations obtained by the SM-only fit.

Table 16: 0b 1e3j

| H_T Bin (GeV) | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075– ∞ |
|---------------------|---------------------|--------------------|--------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| SM hadronic | 2744^{+48}_{-43} | 771^{+21}_{-23} | 254^{+13}_{-13} | $76.5^{+6.1}_{-4.8}$ | $33.7^{+3.7}_{-3.8}$ | $11.8^{+1.9}_{-2.1}$ | $6.3^{+1.4}_{-1.3}$ | $3.2^{+1.0}_{-0.9}$ |
| Data hadronic | 2728 | 766 | 257 | 77 | 32 | 9 | 9 | 4 |
| SM μ +jets | 9072^{+97}_{-113} | 3543^{+56}_{-62} | 1539^{+39}_{-41} | 686^{+25}_{-26} | 325^{+17}_{-17} | 158^{+13}_{-12} | $78.6^{+7.8}_{-8.3}$ | $54.1^{+7.0}_{-6.8}$ |
| Data μ +jets | 9078 | 3545 | 1538 | 686 | 326 | 159 | 78 | 54 |
| SM γ +jets | 3990^{+54}_{-61} | 1203^{+34}_{-37} | 410^{+17}_{-19} | 127^{+10}_{-10} | $48.8^{+6.0}_{-6.9}$ | $19.9^{+3.3}_{-4.2}$ | $12.1^{+3.0}_{-2.9}$ | $7.7^{+2.9}_{-2.7}$ |
| Data γ +jets | 4000 | 1206 | 408 | 127 | 50 | 22 | 10 | 7 |

Table 17: 0b ge4j

| H_T Bin (GeV) | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075– ∞ |
|---------------------|--------------------|-------------------|-------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| SM hadronic | 456^{+15}_{-14} | 291^{+12}_{-12} | 148^{+8}_{-7} | $66.0^{+5.6}_{-5.2}$ | $27.1^{+2.9}_{-3.4}$ | $14.0^{+1.9}_{-2.1}$ | $6.5^{+1.5}_{-1.2}$ | $3.2^{+1.0}_{-0.9}$ |
| Data hadronic | 480 | 299 | 158 | 66 | 28 | 15 | 6 | 2 |
| SM μ +jets | 1260^{+36}_{-36} | 891^{+29}_{-30} | 566^{+23}_{-23} | 308^{+20}_{-14} | 162^{+11}_{-12} | $81.3^{+8.6}_{-8.1}$ | $46.9^{+7.3}_{-6.1}$ | $28.6^{+6.4}_{-4.4}$ |
| Data μ +jets | 1249 | 888 | 562 | 308 | 162 | 81 | 47 | 29 |
| SM γ +jets | 439^{+20}_{-18} | 349^{+16}_{-18} | 161^{+10}_{-13} | $83.0^{+7.9}_{-8.5}$ | $32.6^{+5.7}_{-5.6}$ | $21.8^{+3.6}_{-4.2}$ | $7.7^{+2.4}_{-2.0}$ | $4.2^{+1.7}_{-1.7}$ |
| Data γ +jets | 427 | 344 | 155 | 83 | 32 | 21 | 8 | 5 |

Table 18: 1b le3j

| H_T Bin (GeV) | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075– ∞ |
|---------------------|--------------------|-------------------|----------------------|----------------------|----------------------|----------------------|----------------------|---------------------|
| SM hadronic | 426^{+15}_{-17} | 114^{+6}_{-6} | $35.5^{+3.3}_{-2.8}$ | $10.1^{+1.4}_{-1.5}$ | $3.7^{+0.9}_{-0.8}$ | $1.6^{+0.7}_{-0.6}$ | $0.5^{+0.3}_{-0.4}$ | $0.1^{+0.1}_{-0.0}$ |
| Data hadronic | 444 | 118 | 36 | 15 | 3 | 2 | 1 | 0 |
| SM μ +jets | 2282^{+50}_{-45} | 789^{+26}_{-27} | 325^{+20}_{-17} | 139^{+11}_{-14} | $62.7^{+9.2}_{-6.5}$ | $25.1^{+4.8}_{-5.1}$ | $16.1^{+3.9}_{-4.1}$ | $7.9^{+2.9}_{-2.9}$ |
| Data μ +jets | 2272 | 787 | 325 | 137 | 63 | 25 | 16 | 8 |
| SM γ +jets | 452^{+20}_{-22} | 146^{+9}_{-10} | $49.3^{+6.6}_{-5.9}$ | $18.1^{+3.8}_{-4.4}$ | $5.6^{+2.4}_{-2.4}$ | $4.3^{+1.9}_{-1.8}$ | $1.4^{+1.0}_{-1.4}$ | $0.0^{+0.0}_{-0.0}$ |
| Data γ +jets | 444 | 144 | 49 | 15 | 6 | 4 | 1 | 0 |

Table 19: 1b ge4j

| H_T Bin (GeV) | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075– ∞ |
|---------------------|--------------------|----------------------|----------------------|----------------------|-----------------------|----------------------|----------------------|----------------------|
| SM hadronic | 190^{+10}_{-8} | 120^{+6}_{-5} | $45.6^{+3.1}_{-3.8}$ | $17.1^{+2.6}_{-1.9}$ | $6.8^{+1.5}_{-1.3}$ | $5.4^{+1.3}_{-1.6}$ | $2.4^{+0.9}_{-0.9}$ | $1.2^{+0.7}_{-0.8}$ |
| Data hadronic | 206 | 135 | 45 | 14 | 8 | 6 | 2 | 0 |
| SM μ +jets | 1250^{+37}_{-28} | 891^{+27}_{-31} | 385^{+21}_{-20} | 200^{+12}_{-12} | $86.6^{+9.0}_{-10.1}$ | $55.3^{+7.7}_{-7.3}$ | $24.9^{+4.4}_{-5.3}$ | $10.7^{+3.1}_{-3.4}$ |
| Data μ +jets | 1238 | 881 | 385 | 202 | 86 | 55 | 25 | 11 |
| SM γ +jets | 102^{+10}_{-9} | $81.3^{+8.2}_{-7.9}$ | $32.7^{+5.5}_{-5.6}$ | $11.9^{+3.7}_{-3.0}$ | $7.6^{+2.6}_{-2.6}$ | $3.3^{+1.2}_{-1.6}$ | $3.7^{+1.9}_{-1.9}$ | $1.1^{+0.8}_{-1.1}$ |
| Data γ +jets | 98 | 77 | 33 | 13 | 7 | 3 | 4 | 2 |

Table 20: 2b le3j

| H_T Bin (GeV) | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075– ∞ |
|---------------------|----------------------|----------------------|----------------------|----------------------|----------------------|---------------------|---------------------|---------------------|
| SM hadronic | $65.0^{+4.3}_{-4.3}$ | $18.4^{+1.7}_{-1.6}$ | $4.2^{+0.6}_{-0.5}$ | $1.1^{+0.3}_{-0.2}$ | $0.2^{+0.1}_{-0.1}$ | $0.0^{+0.0}_{-0.0}$ | $0.0^{+0.0}_{-0.0}$ | $0.0^{+0.0}_{-0.0}$ |
| Data hadronic | 78 | 18 | 8 | 3 | 0 | 0 | 0 | 0 |
| SM μ +jets | 681^{+26}_{-31} | 217^{+14}_{-14} | $78.8^{+9.4}_{-9.3}$ | $19.9^{+4.1}_{-3.8}$ | $14.8^{+4.0}_{-3.9}$ | $5.0^{+2.0}_{-2.1}$ | $2.0^{+1.0}_{-1.0}$ | $1.0^{+1.0}_{-1.0}$ |
| Data μ +jets | 668 | 217 | 75 | 18 | 15 | 5 | 2 | 1 |
| Data γ +jets | 36 | 8 | 4 | 0 | 0 | 1 | 0 | 0 |

Table 21: 2b ge4j

| H_T Bin (GeV) | 375–475 | 475–575 | 575–675 | 675–775 | 775–875 | 875–975 | 975–1075 | 1075– ∞ |
|---------------------|----------------------|----------------------|----------------------|---------------------|----------------------|----------------------|----------------------|---------------------|
| SM hadronic | $73.6^{+4.2}_{-4.2}$ | $45.7^{+2.8}_{-2.9}$ | $20.4^{+1.8}_{-1.8}$ | $7.7^{+1.2}_{-1.0}$ | $1.9^{+0.3}_{-0.3}$ | $0.9^{+0.2}_{-0.2}$ | $0.4^{+0.1}_{-0.1}$ | $0.4^{+0.1}_{-0.2}$ |
| Data hadronic | 79 | 52 | 31 | 12 | 1 | 2 | 0 | 1 |
| SM μ +jets | 765^{+26}_{-27} | 521^{+23}_{-22} | 285^{+15}_{-17} | 128^{+12}_{-10} | $54.1^{+7.5}_{-8.0}$ | $20.1^{+4.6}_{-4.1}$ | $10.6^{+2.9}_{-2.9}$ | $9.6^{+2.9}_{-2.9}$ |
| Data μ +jets | 760 | 515 | 274 | 124 | 55 | 19 | 11 | 9 |
| Data γ +jets | 19 | 17 | 10 | 4 | 0 | 2 | 1 | 0 |

.3 Efficiencies and systematic uncertainties for simplified models

.3.1 T2cc




Figure 12: Hadronic selection efficiency times acceptance for T2cc for the relevant event categories defined by n_{jet} and n_b . Note the different z-axis scales.

.3.2 T2tt




Figure 13: The total systematic uncertainty in the signal efficiency times acceptance for all relevant event categories for the T2cc interpretation.




Figure 14: The total systematic uncertainty in the signal efficiency times acceptance for all relevant event categories for the T2tt interpretation.

3.3 T2tt




Figure 15: Hadronic selection efficiency times acceptance for the T2tt for the relevant event categories defined by n_{jet} and n_b . Note the different z-axis scales.

Bibliography

- [1] S. Chatrchyan et al. Missing transverse energy performance of the cms detector. *JINST*, 6:P09001, 2011.
- [2] Serguei Chatrchyan et al. Determination of Jet Energy Calibration and Transverse Momentum Resolution in CMS. *JINST*, 6:P11002, 2011.
- [3] V. Khachatryan et al. Search for Supersymmetry in pp Collisions at 7 TeV in Events with Jets and Missing Transverse Energy. *Phys. Lett. B*, 698:196, 2011.
- [4] L. Randall and D. Tucker-Smith. Dijet searches for supersymmetry at the large hadron collider. *Phys. Rev. Lett.*, 101:221803, 2008.