

Matrix Spaces; Rank 1; Small World Graphs

Vectors don't have to be columns of numbers. Whole matrices can be vectors too, and a covariance matrix lives in one such space. Then the simplest matrices of all, rank one, which is what a single-factor market looks like. And a first look at graphs.

PART 1 Index

What this lesson covers, in order. Click a topic to jump to it.

1. Conditions and important concepts

12 short definitions: matrix spaces, symmetric / upper / diagonal subspaces, rank one matrices, graphs.

2. Procedure

Find the dimension of a matrix space; test for rank one and split it into uv^T .

3. 2×2 examples

A two-asset covariance matrix; a one-factor return table.

4. 3×3 examples

A three-asset covariance matrix; one factor (rank 1) and two factors (rank 2).

5. Bigger example: the market model

Covariance = $\sigma^2 \beta \beta^T + D$. 1,275 numbers become 101. With a factor builder.

6. Small world graphs

A broker network: distances, the adjacency matrix and two-step links.

7. Final memory table

Everything on one screen.

METHODS IN THIS LESSON

[Matrix spaces](#)
[Symmetric matrices S](#)
[dim\(S + U\) formula](#)
[Rank one: \$uv^T\$](#)
[Single-factor model](#)
[Graphs and distance](#)
[Adjacency matrix](#)

PART 2 Prerequisites

Click any link to review it, then come back.

- Basis and dimension (Lesson 9) and the four subspaces (Lesson 10).
- Symmetric matrices and covariance (Lesson 5).

- Columns times rows (Lesson 3): a column times a row is a whole matrix.

PART 3 Explanation

THE NUMBERS USED IN THE EXAMPLES

Betas (how much a fund moves when the market moves 1%): Nifty 1, midcap 1.2, balanced 0.5, bonds 0. Market returns in three scenarios (bad, normal, good year): -10% , 5% , 20% . Market variance $\sigma^2 = 400$ (a 20% volatility, since $20^2 = 400$).

1. Conditions and important concepts

1. Vectors don't have to be columns

A vector space only needs two moves: **add** two of its things and **scale** one by a number, without leaving the space (Lesson 5). Matrices can do that. So can functions. So:

Anything you can add and scale can be a "vector".

2. The space M of all 3×3 matrices

DEFINITION · MATRIX SPACE M

M = all 3×3 matrices. Adding two gives a 3×3 matrix; scaling gives one; the zero matrix is in it. It has **dimension 9**: you choose 9 numbers.

A basis: the 9 matrices with a single 1 and zeros elsewhere, $E_{11}, E_{12}, \dots, E_{33}$ (E_{ij} has its 1 in row i , column j). Every matrix is $\sum a_{ij} E_{ij}$, and no E_{ij} is a mix of the others. M behaves just like $\mathbb{R}^9$.

3. Symmetric matrices S : where covariance lives

DEFINITION · THE SUBSPACE S OF SYMMETRIC MATRICES

S = all 3×3 matrices with $A^T = A$. Sums and multiples of symmetric matrices are symmetric, so S is a subspace. **Dimension 6**: choose the 3 diagonal entries and the 3 entries above the diagonal; the 3 below are copies.

A covariance matrix is symmetric (Lesson 5), so it lives in S . For n assets you need

$$\boxed{\dim S = \frac{n(n+1)}{2}} \quad n = 3 : 6 \quad n = 50 : \frac{50 \times 51}{2} = 1275$$

4. Upper triangular matrices U

All 3×3 matrices with zeros below the diagonal. A subspace of dimension 6 (3 diagonal + 3 above). Basis: $E_{11}, E_{12}, E_{13}, E_{22}, E_{23}, E_{33}$.

5. Diagonal matrices $D = S \cap U$

Matrices that are both symmetric and upper triangular must have zeros off the diagonal. Dimension 3. A diagonal covariance matrix means the assets are **uncorrelated**: only their own variances, no covariances.

6. Union vs sum

$S \cup U$ ("symmetric or upper triangular") is **not** a subspace: adding a symmetric matrix to an upper triangular one usually gives neither. The **sum** $S + U$ (all $s + u$) is a subspace, and here it is all of M .

RULE · DIMENSIONS OF A SUM AND AN INTERSECTION

$$\dim S + \dim U = \dim(S + U) + \dim(S \cap U) \quad 6 + 6 = 9 + 3 \checkmark$$

7. Rank one matrices

RULE · EVERY RANK ONE MATRIX IS UV^T

A matrix has rank 1 exactly when it is a column times a row: $A = uv^T$. Every column is a multiple of u ; every row is a multiple of v^T .

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 8 & 10 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 4 & 5 \end{bmatrix} \quad (\text{MIT's example})$$

8. Spotting rank one

Eliminate: if only **one** non-zero row survives, the rank is 1. Equivalently, every row is a multiple of the first non-zero row. For a 2×2 : $ad - bc = 0$ (and not all zero).

9. Rank one in markets: one factor

MODEL · SINGLE-FACTOR RETURNS

If every fund's return is its beta times the market's return, then the table of returns (funds $\times$ scenarios) is βr_m^T : a column of betas times a row of market returns. **One factor drives everything $\Rightarrow$ the table has rank 1.**

10. Rank one matrices are building blocks

A matrix of rank r can be written as a sum of r rank-one matrices. In markets: an r -factor model. But the sum of two rank-one matrices is usually rank 2, so the set of rank-one matrices is **not** a subspace (and neither is "all rank 4 matrices", MIT's example).

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{rank 1} + \text{rank 1} = \text{rank 2}$$

11. A subspace from a rule: zero-cost trades with four funds

All (v_1, v_2, v_3, v_4) with $v_1 + v_2 + v_3 + v_4 = 0$ is the null space of $A = \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}$. Rank 1, so its dimension is $4 - 1 = 3$, with basis $(-1, 1, 0, 0)$, $(-1, 0, 1, 0)$, $(-1, 0, 0, 1)$: switch Nifty into each other fund. The other three subspaces: $C(A) = \mathbb{R}^1$, row space of dimension 1, left null space $\{0\}$.

12. Graphs

DEFINITION · GRAPH AND DISTANCE

A **graph** is a set of **nodes** joined by **edges**. The **distance** between two nodes is the smallest number of edges on a path between them. A "small world" graph has short distances even with many nodes ("six degrees of separation").

2. Procedure

A. Dimension of a matrix space

Step	What to do
1	Check it's a subspace: the zero matrix is in it, and sums and multiples stay in it.
2	Count how many entries you can choose freely; the rest are forced.
3	Basis: one matrix per free choice (a 1 in that spot, plus whatever copies it forces).
4	Dimension = number of basis matrices.

B. Is it rank one? Split it into uv^T

Step	What to do
1	Take the first non-zero row: that's v^T .
2	For each row, find the number that turns v^T into it: those numbers form u .
3	If some row isn't a multiple of v^T , the rank is more than 1.
4	Check $uv^T = A$ entry by entry.

3. 2×2 examples

Example A: a two-asset covariance matrix (dimension 3)

Nifty variance 400, bonds variance 25, covariance 30:

$$\text{Cov} = \begin{bmatrix} 400 & 30 \\ 30 & 25 \end{bmatrix} = 400 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 25 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + 30 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Check entry by entry: top-left $400 + 0 + 0 = 400 \checkmark$, bottom-right $0 + 25 + 0 = 25 \checkmark$, both off-diagonal $0 + 0 + 30 = 30 \checkmark$. The three matrices on the right are a basis of the 2×2 symmetric matrices, so that space has **dimension 3** = $\frac{2 \times 3}{2}$: two variances and one covariance.

Example B: a one-factor return table (rank 1)

Nifty (beta 1) and midcap (beta 1.2), market return -10% in a bad year and 20% in a good year:

$$\begin{bmatrix} 1 \\ 1.2 \end{bmatrix} \begin{bmatrix} -10 & 20 \end{bmatrix} = \begin{bmatrix} 1(-10) & 1(20) \\ 1.2(-10) & 1.2(20) \end{bmatrix} = \begin{bmatrix} -10 & 20 \\ -12 & 24 \end{bmatrix}$$

- Elimination: row 2 $- 1.2 \times$ row 1 $= (-12 + 12, 24 - 24) = (0, 0)$. One non-zero row: **rank 1**.
- $ad - bc = (-10)(24) - (20)(-12) = -240 + 240 = 0 \checkmark$.
- Meaning: midcap's row is just $1.2 \times$ Nifty's row. One factor (the market) explains the whole table.

4. 3×3 examples

Example A: a three-asset covariance matrix (dimension 6)

Nifty, midcap, bonds:

$$\text{Cov} = \begin{bmatrix} 400 & 480 & 0 \\ 480 & 900 & 0 \\ 0 & 0 & 25 \end{bmatrix}$$

Six numbers decide it: the variances 400, 900, 25 and the covariances 480 (Nifty-midcap), 0 (Nifty-bonds), 0 (midcap-bonds). A basis of S : $E_{11}, E_{22}, E_{33}, E_{12} + E_{21}, E_{13} + E_{31}, E_{23} + E_{32}$. So

$$\text{Cov} = 400E_{11} + 900E_{22} + 25E_{33} + 480(E_{12} + E_{21}) + 0(E_{13} + E_{31}) + 0(E_{23} + E_{32}).$$

Example B: one factor, three funds (rank 1)

$$R = \beta r_m^T = \begin{bmatrix} 1 \\ 1.2 \\ 0.5 \end{bmatrix} \begin{bmatrix} -10 & 5 & 20 \end{bmatrix} = \begin{bmatrix} -10 & 5 & 20 \\ -12 & 6 & 24 \\ -5 & 2.5 & 10 \end{bmatrix} \begin{array}{l} \leftarrow \text{Nifty} \\ \leftarrow \text{midcap} \\ \leftarrow \text{balanced} \end{array}$$

- Row 2 $- 1.2 \times$ row 1: $(-12 + 12, 6 - 6, 24 - 24) = (0, 0, 0)$.
- Row 3 $- 0.5 \times$ row 1: $(-5 + 5, 2.5 - 2.5, 10 - 10) = (0, 0, 0)$.
- One non-zero row left: **rank 1**. Every column is a multiple of $\beta = (1, 1.2, 0.5)$; every row is a multiple of $r_m^T = (-10, 5, 20)$.

Example C: add a second factor (rank 2)

Suppose midcap also has a small-cap factor that adds $(4, -2, 6)$ to its returns. That's a second rank-one piece:

$$R_2 = \begin{bmatrix} 1 \\ 1.2 \\ 0.5 \end{bmatrix} \begin{bmatrix} -10 & 5 & 20 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \begin{bmatrix} 4 & -2 & 6 \end{bmatrix} = \begin{bmatrix} -10 & 5 & 20 \\ -8 & 4 & 30 \\ -5 & 2.5 & 10 \end{bmatrix}$$

Midcap's row: $(-12 + 4, 6 - 2, 24 + 6) = (-8, 4, 30)$. Is it a multiple of Nifty's row? $-8 / -10 = 0.8$, $4 / 5 = 0.8$, but $30 / 20 = 1.5$. No. Balanced is still $0.5 \times$ Nifty. So **rank 2**: two factors, two rank-one pieces.

5. Bigger example: the market model

Where did Example A's covariance matrix come from? From a **single-factor model**: each fund's risk is (its beta $\times$ the market's risk) plus its own private risk.

$$\text{Cov} = \sigma^2 \beta \beta^T + D \quad \beta = \begin{bmatrix} 1 \\ 1.2 \\ 0 \end{bmatrix} \text{ (Nifty, midcap, bonds)}, \quad \sigma^2 = 400$$

Step 1, the rank-one market part:

$$400 \beta \beta^T = 400 \begin{bmatrix} 1 \cdot 1 & 1 \cdot 1.2 & 1 \cdot 0 \\ 1.2 \cdot 1 & 1.2 \cdot 1.2 & 1.2 \cdot 0 \\ 0 & 0 & 0 \end{bmatrix} = 400 \begin{bmatrix} 1 & 1.2 & 0 \\ 1.2 & 1.44 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 400 & 480 & 0 \\ 480 & 576 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 2, the private (diagonal) part: Nifty 0 (it is the market), midcap 324, bonds 25:

$$D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 324 & 0 \\ 0 & 0 & 25 \end{bmatrix} \quad \text{Cov} = \begin{bmatrix} 400 & 480 & 0 \\ 480 & 576 + 324 & 0 \\ 0 & 0 & 25 \end{bmatrix} = \begin{bmatrix} 400 & 480 & 0 \\ 480 & 900 & 0 \\ 0 & 0 & 25 \end{bmatrix}$$

Why it matters. A full covariance matrix for 50 stocks lives in S , with $\dim = 1275$ numbers to estimate. The market model needs only 50 betas + 1 market variance + 50 private variances = 101. A rank-one piece plus a diagonal is a huge simplification, and it's how risk models start.

1275 numbers (full covariance) $\rightarrow$ 101 numbers (one factor)

Try it: build a factor table

Betas: Nifty Midcap Balanced

Market return (%): bad normal good

☐ Add a small-cap factor to midcap: $(4, -2, 6)$

Return table (rows: Nifty, midcap, balanced; columns: bad, normal, good):

$$\beta r_m^T = \begin{bmatrix} -10 & 5 & 20 \\ -12 & 6 & 24 \\ -5 & 2.5 & 10 \end{bmatrix}$$

Rank 1: one factor explains the whole table (every row is a multiple of $(-10, 5, 20)$).

6. Small world graphs

Six brokers A-F. An edge means two brokers trade directly with each other:

Edges	A-B, A-C, B-C, B-D, C-D, D-E, E-F
--------------	-----------------------------------

METHOD · ADJACENCY MATRIX

Put a 1 in row i , column j when nodes i and j are joined, 0 otherwise. It is symmetric. Its square counts **two-step** paths: entry (i, j) of A^2 = number of ways to go from i to j in exactly two steps.

$$A = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix} \quad A^2 = \begin{bmatrix} 2 & 1 & 1 & 2 & 0 & 0 \\ 1 & 3 & 2 & 1 & 1 & 0 \\ 1 & 2 & 3 & 1 & 1 & 0 \\ 2 & 1 & 1 & 3 & 0 & 1 \\ 0 & 1 & 1 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \end{bmatrix}$$

- $(A^2)_{AD} = 2$: A reaches D in two steps two ways, A-B-D and A-C-D. Check: row A $(0, 1, 1, 0, 0, 0) \cdot$ column D $(0, 1, 1, 0, 1, 0) = 0 + 1 + 1 + 0 + 0 + 0 = 2 \checkmark$.
- Diagonal of A^2 = number of trading partners: B has 3 (A, C, D) $\checkmark$.
- Distances from A: B 1, C 1, D 2, E 3, F 4. The farthest pair (A and F) is 4 steps apart.

Why a trader cares. In a financial network, distance is how many hops a shock takes to travel.

"Small world" networks, where any two banks are only a few hops apart, spread trouble fast.

Lesson 12 turns graphs into matrices properly, with directions on the edges.

7. Final memory table

Concept	Key fact	Example
Matrix space M (3×3)	dimension 9; basis E_{ij}	behaves like $\mathbb{R}^9$
Symmetric S	dimension $n(n + 1)/2$	covariance: 6 numbers for 3 assets, 1275 for 50
Upper triangular U	dimension 6 (for 3×3)	U from elimination
Diagonal $D = S \cap U$	dimension 3	uncorrelated assets
Sum rule	$\dim S + \dim U = \dim(S + U) + \dim(S \cap U)$	$6 + 6 = 9 + 3$
Rank one	$A = uv^T$; every row a multiple of one row	one-factor returns βr_m^T
Rank r	a sum of r rank-one pieces	an r -factor model
Not subspaces	$S \cup U$; all rank-1 matrices	rank 1 + rank 1 = rank 2
Graph	nodes + edges; distance = fewest edges	broker network, A to F = 4
Adjacency A	$(A^2)_{ij}$ counts two-step paths	A to D: 2 ways

▼ Check yourself (MIT problem 11.2): with $A = \begin{bmatrix} 1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$ acting on 3×3 matrices X , which X give $AX = 0$, and what are the dimensions?

A has rank 2 and $N(A)$ is the multiples of $(1, 1, 1)$. $AX = 0$ means every column of X is in $N(A)$, so $X = \begin{bmatrix} a & b & c \\ a & b & c \\ a & b & c \end{bmatrix}$: dimension 3. The matrices AX are those whose columns each add to 0: dimension 6. And $3 + 6 = 9 = \dim M$.

▼ Check yourself (MIT recitation): the 2×3 matrices whose null space contains $(2, 1, 1)$. A subspace? Dimension?

Yes: if $A(2, 1, 1) = 0$ and $B(2, 1, 1) = 0$, then $(A + B)(2, 1, 1) = 0$ and $(cA)(2, 1, 1) = 0$. Each row (r_1, r_2, r_3) must satisfy $2r_1 + r_2 + r_3 = 0$: 2 free numbers per row, 2 rows, so **dimension 4**.