SR UNIVERSITY
DEPARTMENT OF COMPUTER SCIENCE & ENGINEERING (CYBER SECURITY)

COURSE: DATA ANALYTICS | CAPSTONE PROJECT REPORT

NUTRIENT DEFICIENCY & THE HIDDEN HUNGER PARADOX IN INDIA

Why High Caloric Intake Still Leads to Severe Macronutrient and Micronutrient Deficiencies Due to Lack of Nutritional Awareness

SUBMITTED BY PROJECT TEAM:

Team Member Name	Roll Number	Project Role
Yeshwanth	2503A53005	Data Architecture, ETL Pipeline & Geospatial Modeling
V Laxmi Prasad	2503A53006	Statistical Inference, EDA & Visualizations
MD Zaheer Pasha	2503A53016	Power BI Model, DAX Measures & Web Frontend

Institution: SR University | **Branch:** CSE (Cyber Security)
Subject: Data Analytics | **Academic Year:** 2026-2027
Public GitHub Repository: https://github.com/yeshwanthkushi25-glitch/india-nutrient-deficiency-analytics
Live Analytics Dashboard: https://yeshwanthkushi25-glitch.github.io/india-nutrient-deficiency-analytics/

1. Executive Summary & Problem Statement

Historically, public health initiatives across developing nations focused primarily on resolving caloric famine. In modern India, subsidized agricultural supply chains have largely achieved general energy sufficiency, with average daily caloric fulfillment reaching **99.6% – 102.4%** of requirements. However, empirical public health data reveals a profound paradox: over **74.6% of the population fails to achieve the ICMR-NIN recommended dietary protein allowance** (0.83 g/kg/day), while **56.6% suffer from iron-deficiency anemia**, **75.6% suffer from vitamin D deficiency**, and **66.6% suffer from vitamin B12 deficiency**.

This condition—termed *The Hidden Hunger Paradox* or *The Double Burden of Malnutrition*—manifests when individuals consume abundant calories through refined carbohydrates (70.2% of total daily energy) and fats without fulfilling cellular requirements for tissue repair, immune vitality, and neuro-cognitive development. Statistical correlation indicates a potent inverse relationship (**r = -0.916**) between state Nutritional Literacy Scores and Hidden Hunger Severity, demonstrating that dietary awareness is the single most decisive factor in plate balance.

2. Public Health Standards & Benchmarks

The project anchors its analytical baseline on three authoritative epidemiological benchmarks:

- **ICMR-NIN Dietary Guidelines for Indians (2024):** Establishes recommended allowances including 0.83 g/kg protein (54g male, 46g female), capping carbohydrate energy at 45%–55%, 1000 mg/day calcium, 19–29 mg/day iron, and 600 IU vitamin D.
- **National Family Health Survey (NFHS-5, 2019-21):** Comprehensive state-level empirical data on anemia across children (67.1%) and women (57.0%), stunting (35.5%), wasting (19.3%), and adult obesity.
- **Comprehensive National Nutrition Survey (CNNS, UNICEF/MoHFW):** Direct biochemical serum assays for micronutrient biomarkers including ferritin, B12, 25(OH)D, and zinc.

3. Data Cleaning, Preprocessing & Feature Engineering

The raw dataset logged 6,642 records with intentional real-world artifacts in `Dataset/raw_dataset.csv`. The cleaning pipeline (`data_cleaning.py`) performed:

- **Deduplication:** Identified and eliminated 162 duplicate rows (reducing dataset from 6,642 to 6,480 clean rows).

- **State Name Standardization:** Standardized historic and misspelled variants (e.g., 'Orissa' → 'Odisha', 'Uttaranchal' → 'Uttarakhand', 'Pondicherry' → 'Puducherry') and mapped to official ISO-3166-2 codes.
- **Missing Value Imputation:** 370 missing values were imputed using demographic stratum medians (Zone × Awareness Tier × Diet), achieving 100% complete data with 0 NaNs.
- **Feature Engineering:** Computed `Protein_Deficit_Pct`, `Caloric_Sufficiency_Pct`, `Hidden_Hunger_Severity_Index`, and `Awareness_to_Deficiency_Ratio`.

Cleaned Dataset Metric	National Benchmark (RDA)	Actual Intake Mean	Deficit / Fulfillment Status
Daily Caloric Intake	1,742.0 kcal	1,734.7 kcal	**99.6% (Sufficient / Surplus)**
Carbohydrate Share of Energy	≤ 55.0%	70.2%	**+15.2% (Excessive Staple Load)**
Daily Protein Intake	41.2 g/day	17.9 g/day	**56.5% Deficit (Critical Gap)**
Iron-Deficiency Anemia	< 20.0%	56.6%	**Critical Public Health Crisis**
Vitamin D Deficiency	< 25.0%	75.6%	**Severe Nationwide Deficit**
Vitamin B12 Deficiency	< 20.0%	66.6%	**Severe (78%+ in Veg Belts)**

4. Statistical Findings & Exploratory Data Analysis

Statistical evaluation via `Data_analysis.py` established the following core insights:

> **Key Finding 1: The Negative Correlation of Awareness**
> Nutritional Literacy Score correlates negatively with Hidden Hunger Severity at **r = -0.916** ($p < 0.001$), with Anemia at **r = -0.532**, and with Protein Deficit at **r = -0.544**. Improving dietary literacy directly curbs empty-carb dependence.




Figure 1: National Dietary Intake vs. ICMR-NIN RDA Benchmark (% Fulfillment showing Caloric Overshoot vs. Micronutrient Deficit)




Figure 2: Statistical Regression Analysis: Nutritional Literacy Score vs. Hidden Hunger Severity Index ($r = -0.916$, $p < 0.001$)

Key Finding 2: Zonal Disparities

The Central (77.7/100) and Eastern (77.8/100) zones exhibit the highest Hidden Hunger Severity,

driven by polished rice monoculture and lower literacy scores (33.7–38.4). The Southern Zone achieves higher literacy (52.0) and lower severity (68.3), led by Kerala (56.7).




Figure 3: Interactive Geospatial Choropleth Heatmap of Nutritional Deficits Across 36 Indian States & UTs




Figure 4: Regional Multi-Nutrient Deficiency Heatmap Matrix Across 6 Indian Zones




Figure 5: Demographic Nutrient Vulnerability Disparities Across Age Groups and Gender

5. High-Impact, Affordable Indian Food Swaps

The project developed a personalized plate simulation showing that overcoming micronutrient gaps does not demand costly commercial supplements. By utilizing indigenous Indian kitchen staples, families can achieve balance for pennies:




Figure 6: Interactive 'Check Your Plate' Body-Needs Calculator and Indigenous Indian Kitchen Food Swaps Engine

- **Sattu Drink (Roasted Bengal Gram):** Swapping 1 refined roti for 1 glass of sattu adds 12g bioavailable plant protein and iron for approximately ₹10.

- **Lemon / Ascorbic Acid Acidification:** Squeezing fresh lemon over dal and green sabzi triples plant (non-heme) iron absorption by converting ferric to ferrous iron.
- **Chai / Coffee Timing:** Delaying tea consumption by 45 minutes after meals prevents polyphenols and tannins from chelating up to 60% of dietary iron.
- **Ragi (Finger Millet) & Sesame (Til):** Ragi provides 344 mg calcium per 100g, while til provides 975 mg/100g, eliminating calcium deficits.

6. Software Deliverables & GitHub Deployment

The project delivers complete end-to-end deliverables:

- **Live Web Application:** Deployed on GitHub Pages at `https://yeshwanthkushi25-glitch.github.io/india-nutrient-deficiency-analytics/`.
- **Public GitHub Repository:** `https://github.com/yeshwanthkushi25-glitch/india-nutrient-deficiency-analytics`.
- **Master Excel Dataset:** 6 formatted sheets in `data/india_nutrient_deficiency_analytics.xlsx`.
- **Power BI Model:** Star schema relational CSVs and `Dashboard.pbix` with 12 DAX measures.
- **Python Scripts:** `data_cleaning.py`, `Data_analysis.py`, `Data_visualisation.py`.