

11. Radium-226 ($^{226}_{88}\text{Ra}$) undergoes α -decay into radon (Rn).

- (a) Write a nuclear equation for the decay. (2 marks)

* (b) Given : mass of a radium nucleus = 226.0254 u
mass of a radon nucleus = 222.0176 u
mass of an α -particle = 4.0026 u

Calculate the energy released in the decay in MeV.


(2 marks)

- (c) 1 curie (Ci) is defined as the activity of 1 g of radium. The activity of a radium source used in laboratories is about 5 μCi . Estimate the number of radium atoms in this source and hence find its activity expressed in disintegrations per second. The half-life of radium-226 is 1600 years and take the mass of one mole of radium as 226 g. (1 $\mu\text{Ci} = 1 \times 10^{-6}$ Ci) (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

10. Scientists had been experimenting controlled fusion in a nuclear reactor in which deuterium (${}^2_1\text{H}$) and tritium (${}^3_1\text{H}$) undergo the following nuclear fusion:



Given: mass of a deuterium nucleus = 2.014102 u
 mass of a tritium nucleus = 3.016049 u
 mass of a helium nucleus = 4.002602 u
 mass of a neutron = 1.008665 u

* (a) Calculate the energy released, in MeV, in the above nuclear fusion. (2 marks)

(b) A deuterium nucleus and a tritium nucleus have to be within 10^{-15} m for nuclear fusion to occur and that a large amount of work done (about 0.4 MeV) is needed to bring two well separated nuclei to such a close distance.

(i) Explain why a large amount of work done is needed and state the kind of energy this work done has become. (2 marks)

In the nuclear reactor, deuterium and tritium exist as plasma, which is a mixture of ions at a very high temperature.

(ii) Explain why a very high temperature is needed for nuclear fusion to occur. (1 mark)

* (iii) Estimate the order of magnitude of the minimum temperature at which fusion of deuterium and tritium nuclei would be possible if the plasma can be regarded as an ideal gas. (Hint: For an ideal gas, the gas molecules each is assumed to have an average kinetic energy $E_K = \frac{3RT}{2N_A}$) (2 marks)

END OF PAPER

Sources of materials used in this paper will be acknowledged in the *Examination Report and Question Papers* published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

(b) As the first decay in the above chain from U to Th has a half-life much longer than those of subsequent decays, the decay from U-238 to Pb-206 can be simplified to a *single decay* with half-life 4.5×10^9 years:



Suppose that a uranium-bearing rock contains only U-238 and no Pb-206 at the time when it was formed long ago by solidification of molten material. In a particular sample of the rock, it is found that the ratio $\frac{\text{number of Pb - 206 atoms}}{\text{number of U - 238 atoms}}$ is $\frac{2}{3}$ at present.


(i) Estimate the age of the rock. Assume that all Pb-206 atoms come from the decay of U-238 originally present in the sample and ignore the small number of U-238 atoms which have decayed but have not yet become Pb-206. (2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

- (iii) The graph in Figure 9.1 shows how the number of U-238 atoms in the sample varies with time t subsequently while $t = 0$ denotes the *present time*. On Figure 9.1, sketch a graph to show the variation of the number of Pb-206 atoms in the sample with time. (2 marks)

Figure 9.1




END OF PAPER

Sources of materials used in this paper will be acknowledged in the *Examination Report and Question Papers* published by the Hong Kong Examinations and Assessment Authority at a later stage.

10. Dust may adhere to the surfaces of photos and films due to electrostatic attraction. To remove the dust effectively, a special brush with a thin slice of polonium-210 ($^{210}_{84}\text{Po}$) fixed near the brush hair as shown in Figure 10.1 may be used. Polonium-210 undergoes α decay and the daughter nucleus lead (Pb) is stable.

Figure 10.1



- (a) Write a nuclear equation for the decay of polonium-210. (2 marks)

- (b) Briefly explain how the α particles help clean the charged dust. (2 marks)

- (c) Briefly explain why the polonium-210 slice must be fixed near to the brush hair. (1 mark)


- *(d) The manufacturer recommends that the brush should be returned to the factory for replacement of the polonium-210 slice every year. Taking the activity of a newly replaced polonium-210 slice as 1 unit, find its activity after one year (365 days). Given: half-life of polonium-210 is 138 days. (2 marks)

END OF PAPER

Sources of materials used in this paper will be acknowledged in the booklet *HKDSE Question Papers* published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

10. (a) The equation below represents nuclear fission of uranium-235 (U-235).



- (i) What is the value of x ? (1 mark)

- (ii) State a necessary condition for chain reaction of fission to occur. (1 mark)

Scientists found evidence in Oklo, Africa that natural nuclear fission occurred two billion (2×10^9) years ago. The uranium mineral ore mined from Oklo **at present** is found to have 0.6% concentration by mass of U-235 (see the table below), which is much lower than usual.

- (b) The table gives the information of U-235 and U-238 in a sample of uranium mineral ore found in Oklo. Given: half-life of U-235 = 7.04×10^8 years

	2×10^9 years ago	at present
U-235	m_0 kg	0.060 kg (i.e. 0.6% concentration by mass)
U-238	13.556 kg	9.940 kg (i.e. 99.4% concentration by mass)

- *(i) Estimate the amount m_0 (in kg) of U-235 in the sample 2×10^9 years ago. (2 marks)

- (ii) Hence determine whether natural nuclear fission of U-235 was possible 2×10^9 years ago. For fission of U-235 to happen, its concentration by mass in the uranium mineral ore has to be at least 3%. (1 mark)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

There must be underground water in the vicinity of this uranium-rich mineral deposit for natural nuclear fission to be possible. Since water can slow down the fast neutrons from fission, these neutrons can easily be captured by U-235.

- (c) In fact the chain reaction stopped even before the concentration by mass of U-235 dropped to 3%. Explain why this occurred. (2 marks)

.....

.....

.....

.....

.....

END OF PAPER

Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.


Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

10. Given: mass of proton = 1.0073 u
 mass of α particle = 4.0015 u
 mass of $^{14}_7\text{N}$ nucleus = 13.9993 u
 mass of $^{17}_8\text{O}$ nucleus = 16.9947 u

When a stationary $^{14}_7\text{N}$ nucleus is bombarded by an α particle, the following nuclear reaction can be triggered with products $^{17}_8\text{O}$ and X fly off:



- (a) What is X ? (1 mark)

- *(b) Based on energy consideration, estimate the minimum kinetic energy, in MeV, of the α particle required for such a nuclear reaction to occur. (2 marks)

- (c) However, when conservation of momentum is also taken into account, the α particle must possess a kinetic energy greater than that found in (b) to bring about such a reaction. Explain. (2 marks)

END OF PAPER


Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

10. Deuterium (${}^2_1\text{H}$) and tritium (${}^3_1\text{H}$) are isotopes of hydrogen. The fusion of two deuterium nuclides can be represented by the following equation:



Given: mass of ${}^2_1\text{H}$ = 2.014102 u
 mass of ${}^3_2\text{He}$ = 3.016029 u
 mass of ${}^1_0\text{n}$ = 1.008665 u

- *(a) 1 in 6420 atoms of hydrogen in nature is a deuterium. Estimate the maximum energy, in MeV, that could be produced from this fusion reaction with 1 mole of hydrogen nuclides. (3 marks)

.....

.....

.....

.....

.....

.....

.....

.....

- (b) If conditions are altered, the fusion of two deuterium nuclides may **not** produce a helium (He) nucleus. Complete the equation below for such a possible fusion reaction. (1 mark)



- (c) Both fission and fusion produce energy. State **TWO** advantages of using fusion as an energy source over fission. (2 marks)

.....

.....

.....

.....

.....

END OF PAPER


Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

9. Part of the decay series of thorium-232 (Th-232) is shown below. Th-232 decays to radium-228 (Ra-228) with a half-life of 1.41×10^{10} years. The end product lead-208 (Pb-208) is stable.



- (a) (i) State the kind of radiation emitted by Th-232. (1 mark)

- (ii) How many α - and β - decay(s) has occurred before a Th-232 nucleus becomes a Pb-208 nucleus ? (2 marks)


- *(b) Estimate the proportion of Th-232 decaying in 10 years. Give your answer correct to 2 significant figures. (2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

- (c) By adding thorium oxide (a compound containing Th-232) to glass, thoriated glass having a higher refractive index is produced. Camera lenses manufactured before 1970 are often made from thoriated glass.



- (i) Although thoriated glass is radioactive, practically it is safe to use a camera with such lenses like the one shown in Figure 9.1. Why ? (1 mark)

- (ii) According to the decay series of thorium-232, thorium oxide would end up as lead oxide which is black in colour. Why is there no need to worry that a thoriated glass lens would turn black when used for several years ? (1 mark)

END OF PAPER

Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

(i) Write the nuclear equations corresponding to each of these two decay processes. (2 marks)

*(ii) The half-life of Bi-213 is 46 minutes. At time $t = 0$ s, its number of undecayed nuclei is 8.0×10^{11} . Find its activity (in Bq) at $t = 0$ s. (2 marks)

*(iii) Find the percentage decrease in the activity after 200 minutes. (2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Figure 12.2(a)

Figure 12.2(b)

- (i) Compared to placing it in a badge worn on the chest, what is the advantage of placing the chip in the ring worn on the hand ? (1 mark)

- (ii) The ring should be worn under the gloves. Explain briefly. (1 mark)

- (iii) Which type(s) of radiation (α , β or γ) can reach the chip of the ring ? (1 mark)

END OF PAPER

Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.


Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

14. In April 1986, a disastrous nuclear accident happened at the Chernobyl Nuclear Power Station. A large quantity of various radioactive substances was released and spread to neighbouring countries. The radiation levels recorded in these countries were much higher than the normal background count rate.

(a) State **ONE** source of background radiation. (1 mark)

- (b) One of the radioactive isotopes released in the accident was caesium-137 (Cs-137). The following equation shows how Cs-137 is produced :



Given : mass of one nuclide of ${}_{92}^{235}\text{U} = 235.0439 \text{ u}$

${}_{55}^{137}\text{Cs} = 136.9071 \text{ u}$

${}_{37}^{95}\text{Rb} = 94.9399 \text{ u}$

${}_0^1\text{n} = 1.0087 \text{ u}$

1 u is equivalent to 931 MeV

(i) What is the value of x ? (1 mark)

*(ii) Find the energy release in the fission of one U-235 nuclide in MeV. (2 marks)

*(iii) The half-life of Cs-137 is 30 years. A soil sample contaminated by Cs-137 has an activity of $1.2 \times 10^6 \text{ Bq}$ (disintegrations per second). A physicist comments that the contaminated sample will affect the environment for more than 350 years. Justify the physicist's claim with calculations. It is known that the activity of an uncontaminated soil sample is 200 Bq. (2 marks)

END OF PAPER

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.