

ch27 Nuclear Energy

Items:

Q11

Q10

Q9

Q10

Q10

Q10

Q10

Q9

Q12

11. Radium-226 ($^{226}_{88}\text{Ra}$) undergoes α -decay into radon (Rn).

(a) Write a nuclear equation for the decay.

(2 marks)

*(b) Given : mass of a radium nucleus = 226.0254 u

mass of a radon nucleus = 222.0176 u

mass of an α -particle = 4.0026 u

Calculate the energy released in the decay in MeV.

(2 marks)

(c) 1 curie (Ci) is defined as the activity of 1 g of radium. The activity of a radium source used in laboratories is about 5 μCi . Estimate the number of radium atoms in this source and hence find its activity expressed in disintegrations per second. The half-life of radium-226 is 1600 years and take the mass of one mole of radium as 226 g. ($1 \mu\text{Ci} = 1 \times 10^{-6} \text{ Ci}$)

(3 marks)


END OF PAPER

Sources of materials used in this paper will be acknowledged in the *Examination Report and Question Papers* published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Q10

10. Scientists had been experimenting controlled fusion in a nuclear reactor in which deuterium (${}^2_1\text{H}$) and tritium (${}^3_1\text{H}$) undergo the following nuclear fusion:



Given: mass of a deuterium nucleus = 2.014102 u
 mass of a helium nucleus = 4.002602 u
 mass of a neutron = 1.008665 u

- (a) Calculate the energy released, in MeV, in the above nuclear fusion.

(2 marks)

- (b) A deuterium nucleus and a tritium nucleus have to be within 10^{-15} m for nuclear fusion to occur and that a large amount of work done (about 0.4 MeV) is needed to bring two well separated nuclei to such a close distance.

- (i) Explain why a large amount of work done is needed and state the kind of energy this work done has become. (2 marks)

Answers written in the margins will not be marked.

In the nuclear reactor, deuterium and tritium exist as plasma, which is a mixture of ions at a very high

- (ii) Explain why a very high temperature is needed for nuclear fusion to occur. (1 mark)

- *(iii) Estimate the order of magnitude of the minimum temperature at which fusion of deuterium and tritium nuclei would be possible if the plasma can be regarded as an ideal gas. (Hint: For an ideal gas, the gas molecules each is assumed to have an average kinetic energy $E_K = \frac{3RT}{2N_A}$) (2 marks)

Sources of materials used in this paper will be acknowledged in the *Examination Report and Question Papers* published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.




Answers written in the margins will not be marked.

- Figure 9.1



Answers written in the margins will not be marked.

Figure 10.1



- (a) Write a nuclear equation for the decay of polonium-210. (2 marks)

- (b) Briefly explain how the α particles help clean the charged dust. (2 marks)

- (c) Briefly explain why the polonium-210 slice must be fixed near to the brush hair. (1 mark)

*(d) The manufacturer recommends that the brush should be returned to the factory for replacement of the polonium-210 slice every year. Taking the activity of a newly replaced polonium-210 slice as 1 unit, find its activity after one year (365 days). Given: half-life of polonium-210 is 138 days. (2 marks)

Answers written in the margins will not be marked

Sources of materials used in this paper will be acknowledged in the booklet *HKDSE Question Papers* published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked

List of data, formulae and relationships

Data

molar gas constant	$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
Avogadro constant	$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
acceleration due to gravity	$g = 9.81 \text{ m s}^{-2}$ (close to the Earth)
universal gravitational constant	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
speed of light in vacuum	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
charge of electron	$e = 1.60 \times 10^{19} \text{ C}$
electron rest mass	$m_e = 9.11 \times 10^{-31} \text{ kg}$
permeability of free space	$\mu_0 = 4\pi \times 10^{-7} \text{ C}^2 \text{ N}^{-1} \text{ A}^{-2}$
permeability of free space	$\mu_0 = 4\pi \times 10^{-7} \text{ N A}^{-2}$
atomic mass unit	$u = 1.66 \times 10^{-27} \text{ kg}$
atomic mass unit	$u = 1.50 \times 10^{-10} \text{ m}$
right angle	$\pi = 3.14$
parsec	$\text{pc} = 3.09 \times 10^{16} \text{ m} = 3.26 \text{ ly} = 206265 \text{ AU}$
Stefan constant	$\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
Planck constant	$h = 6.63 \times 10^{-34} \text{ J s}$

Rectilinear motion

Mathematics

For uniformly accelerated motion :

Equation of a straight line $y = mx + c$

$$y = x + \alpha$$

Q	R	S	T	U	V
1	2	3	4	5	6
7	8	9	10	11	12
13	14	15	16	17	18
19	20	21	22	23	24
25	26	27	28	29	30
31	32	33	34	35	36
37	38	39	40	41	42
43	44	45	46	47	48
49	50	51	52	53	54
55	56	57	58	59	60
61	62	63	64	65	66
67	68	69	70	71	72
73	74	75	76	77	78
79	80	81	82	83	84
85	86	87	88	89	90
91	92	93	94	95	96
97	98	99	100	101	102
103	104	105	106	107	108
109	110	111	112	113	114
115	116	117	118	119	120
121	122	123	124	125	126
127	128	129	130	131	132
133	134	135	136	137	138
139	140	141	142	143	144
145	146	147	148	149	150
151	152	153	154	155	156
157	158	159	160	161	162
163	164	165	166	167	168
169	170	171	172	173	174
175	176	177	178	179	180
181	182	183	184	185	186
187	188	189	190	191	192
193	194	195	196	197	198
199	200	201	202	203	204
205	206	207	208	209	210
211	212	213	214	215	216
217	218	219	220	221	222
223	224	225	226	227	228
229	230	231	232	233	234
235	236	237	238	239	240
241	242	243	244	245	246
247	248	249	250	251	252
253	254	255	256	257	258
259	260	261	262	263	264
265	266	267	268	269	270
271	272	273	274	275	276
277	278	279	280	281	282
283	284	285	286	287	288
289	290	291	292	293	294
295	296	297	298	299	300
301	302	303	304	305	306
307	308	309	310	311	312
313	314	315	316	317	318
319	320	321	322	323	324
325	326	327	328	329	330
331	332	333	334	335	336
337	338	339	340	341	342
343	344	345	346	347	348
349	350	351	352	353	354
355	356	357	358	359	360
361	362	363	364	365	366
367	368	369	370	371	372
373	374	375	376	377	378
379	380	381	382	383	384
385	386	387	388	389	390
391	392	393	394	395	396
397	398	399	400	401	402
403	404	405	406	407	408

$$E = M + \frac{1}{2}mv^2$$

$$\text{Surface area of cylinder} = 2\pi rh + 2\pi r^2$$

$$v^2 = v^2 + 2ax$$

Volume of cylinder = $\pi r^2 h$

Surface area of sphere = $4\pi r^2$

For small angles, $\sin \theta \approx \tan \theta \approx \theta$ (in radians)

Astronomy and Space Science

$U = -\frac{GMm}{r}$	gravitational potential energy
$P = \sigma AT^4$	Stefan's law
$\left \frac{\Delta f}{f_0} \right \approx \frac{v}{c} \left \frac{\Delta \lambda}{\lambda_0} \right $	Doppler effect

Energy and Use of Energy

$E = \frac{\Phi}{A}$	illuminance
$\frac{Q}{t} = -k \frac{A(T_H - T_C)}{d}$	rate of energy transfer by conduction
$U = \frac{K}{d}$	thermal transmittance U-value
$P = \frac{1}{2} \rho A v^3$	maximum power by wind turbine

Atomic World

$$\frac{1}{2} m_e v_{\text{max}}^2 = hf - \phi \quad \text{Einstein's photoelectric equation}$$

$$E_n = -\frac{1}{n^2} \left[\frac{m_e e^4}{8h^2 \epsilon_0^2} \right] = -\frac{13.6 \text{ eV}}{n^2} \quad \text{energy level equation for hydrogen}$$

$$\lambda = \frac{h}{p} = \frac{h}{mv} \quad \text{de Broglie's hypothesis}$$

$$\theta = \frac{1.22\lambda}{d} \quad \text{Rayleigh's criterion for resolving power}$$

6	Medical Physics
---	-----------------

$\theta \approx \frac{1.22\lambda}{d}$	Rayleigh criterion (resolving power)
$\text{power} = \frac{1}{f}$	power of a lens
$L = 10 \log \frac{I}{I_0}$	intensity level (dB)
$Z = \rho c$	acoustic impedance
$\alpha = \frac{I_r}{I_i} = \frac{(Z_2 - Z_1)^2}{(Z_2 + Z_1)^2}$	intensity reflection coefficient
$I = I_0 e^{-\mu x}$	transmitted intensity through a medium

A1. $E = mc \Delta T$	<i>heat energy during heating and cooling</i>	D1. $F = \frac{Gm_1m_2}{r^2}$	Coulomb's law
A2. $E_c = I_c \Delta t$	<i>energy transfer during charge</i>	D2. $\vec{F} = \frac{q\vec{E}}{4\pi\epsilon_0}$	<i>electric field strength due to a point charge</i>
A3. $pV = nRT$	<i>equation of state for an ideal gas</i>	D3. $E = \frac{1}{2} \epsilon_0 E^2$	<i>electric field between parallel plates</i>
A4. $pV = \frac{1}{2} N m \overline{v^2}$	<i>kinetic theory equation</i>	D4. $R = \frac{\rho \ell}{A}$	<i>resistance and resistivity</i>
A5. $E_k = \frac{1}{2} m v^2$	<i>molecular kinetic energy</i>	D5. $R = R_1 + R_2$	<i>resistors in series</i>
B1. $F = m \frac{\Delta v}{\Delta t} = \frac{\Delta p}{\Delta t}$	<i>force</i>	D6. $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$	<i>resistors in parallel</i>
B2. <i>moment</i> = $F \times d$	<i>moment of a force</i>	D7. $F = Bq v \sin \theta$	<i>force on a moving charge in a magnetic field</i>
B3. $E_p = mgh$	<i>gravitational potential energy</i>	D8. $F = B I \sin \theta$	<i>force on a current carrying wire in a magnetic field</i>
B4. $E_k = \frac{1}{2} m v^2$	<i>kinetic energy</i>	D9. $B = \frac{\mu_0 N I}{2r}$	<i>magnetic field strength due to a long straight wire</i>
B5. $E = pV$	<i>mechanical energy</i>	D10. $\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2}$	<i>capacitors in series</i>
B6. $a = \frac{v^2}{r} = \omega^2 r$	<i>centripetal acceleration</i>	D11. $\frac{V_1}{R_1} = \frac{V_2}{R_2}$	<i>ratio of terminal voltage to primary resistance in a transformer</i>
B7. $F = \frac{Gm_1m_2}{r^2}$	<i>Newton's law of gravitation</i>		
C1. $A_p = \frac{V_p}{\omega}$	<i>displacement amplitude</i>	E1. $N = N_0 e^{-\lambda t}$	<i>law of radioactive decay</i>
C2. $d \sin \theta = n \lambda$	<i>diffraction grating equation</i>	E2. $t_{1/2} = \frac{\ln 2}{\lambda}$	<i>half-life and decay constant</i>
C3. $\frac{1}{f} = \frac{1}{u} + \frac{1}{v}$	<i>equation for a single lens</i>	E3. $A = N \lambda$	<i>de Broglie wavelength of particles</i>

10. (a) The equation below represents nuclear fission of uranium-235 (U-235).

$${}^{235}_{92}\text{U} + {}^1_0\text{n} \rightarrow {}^{141}_{54}\text{Ba} + {}^{92}_{36}\text{Kr} + x {}^1_0\text{n} + 200\text{ MeV}$$

(i) What is the value of x ?

(1 mark)

(ii) State a necessary condition for chain reaction of fission to occur.

(1 mark)

Scientists found evidence in Oklo, Africa that natural nuclear fission occurred two billion (2×10^9) years ago. The uranium mineral ore mined from Oklo at present is found to have 0.6% concentration by mass of U-235 (see the table below), which is much lower than usual.

(b) The table gives the information of U-235 and U-238 in a sample of uranium mineral ore found in Oklo. Given: half-life of U-235 = 7.04×10^8 years.

	2×10^9 years ago	at present
U-235	m_0 kg	0.060 kg (i.e. 0.6% concentration by mass)
U-238	13.526 kg	9.940 kg (i.e. 99.4% concentration by mass)

* (i) Estimate the amount m_0 (in kg) of U-235 in the sample 2×10^9 years ago. (2 marks)

(ii) Hence determine whether natural nuclear fission of U-235 was possible 2×10^9 years ago. For fission of U-235 to happen, its concentration by mass in the uranium mineral ore has to be at least 3%. (1 mark)

Answers written in the margins will not be marked.

There must be underground water in the vicinity of this uranium-rich mineral deposit for natural nuclear fission to be possible. Since water can slow down the fast neutrons from fission, these neutrons can easily be captured by U-235.

(c) In fact the chain reaction stopped even before the concentration by mass of U-235 dropped to 3%. Explain why this occurred.

(2 marks)

END OF PAPER

Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Do not write on this page.

Answers written on this page will not be marked.

Q10

10. Given: mass of proton = 1.0073 u
 mass of α particle = 4.0015 u
 mass of $^{14}_7\text{N}$ nucleus = 13.9993 u
 mass of $^{17}_8\text{O}$ nucleus = 16.9947 u

When a stationary $^{14}_7\text{N}$ nucleus is bombarded by an α particle, the following nuclear reaction can be triggered with products $^{17}_8\text{O}$ and X fly off:



(a) What is X ? (1 mark)

*(b) Based on energy consideration, estimate the minimum kinetic energy, in MeV, of the α particle required for such a nuclear reaction to occur. (2 marks)

(c) However, when conservation of momentum is also taken into account, the α particle must possess a kinetic energy greater than that found in (b) to bring about such a reaction. Explain. (2 marks)

END OF PAPER


Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Do not write on this page.

Answers written on this page will not be marked.

10. Deuterium (${}^2_1\text{H}$) and tritium (${}^3_1\text{H}$) are isotopes of hydrogen. The fusion of two deuterium nuclides can be represented by the following equation:



Given: mass of ${}^2_1\text{H} = 2.014102 \text{ u}$

mass of ${}^3_2\text{He} = 3.016029 \text{ u}$

mass of ${}^1_0\text{n} = 1.008665 \text{ u}$

- *(a) 1 in 6420 atoms of hydrogen in nature is a deuterium. Estimate the maximum energy, in MeV, that could be produced from this fusion reaction with 1 mole of hydrogen nuclides. (3 marks)

.....

.....

.....

.....

.....

.....

.....

- (b) If conditions are altered, the fusion of two deuterium nuclides may **not** produce a helium (${}^4_2\text{He}$) nucleus. Complete the equation below for such a possible fusion reaction. (1 mark)



- (c) Both fission and fusion produce energy. State **TWO** advantages of using fusion as an energy source over fission. (2 marks)

.....

.....

.....

.....

.....

END OF PAPER


Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Do not write on this page.

Answers written on this page will not be marked.

9. Part of the decay series of thorium-232 (Th-232) is shown below. Th-232 decays to radium-228 (Ra-228) with a half-life of 1.41×10^{10} years. The end product lead-208 (Pb-208) is stable.



- (a) (i) State the kind of radiation emitted by Th-232. (1 mark)

- (ii) How many α - and β - decay(s) has occurred before a Th-232 nucleus becomes a Pb-208 nucleus? (2 marks)

- *(b) Estimate the proportion of Th-232 decaying in 10 years. Give your answer correct to 2 significant figures. (2 marks)

Answers written in the margins will not be marked.

- (c) By adding thorium oxide (a compound containing Th-232) to glass, thoriated glass having a higher refractive index is produced. Camera lenses manufactured before 1970 are often made from thoriated glass.

Figure 9.1




- (i) Although thoriated glass is radioactive, practically it is safe to use a camera with such lenses like the one shown in Figure 9.1. Why? (1 mark)

- (ii) According to the decay series of thorium-232, thorium oxide would end up as lead oxide which is black in colour. Why is there no need to worry that a thoriated glass lens would turn black when used for several years? (1 mark)

Q12

12. (a) Bismuth-213 (Bi-213) is a radionuclide currently undergoing clinical trials for cancer treatment. It is prepared in liquid and injected into the human body, where it targets and destroys cancer cells. Figure 12.1 shows a decay series of Bi-213 .

Figure 12.1



- (i) Write the nuclear equations corresponding to each of these two decay processes. (2 marks)

.....

.....

.....

- *(ii) The half-life of Bi-213 is 46 minutes. At time $t = 0$ s, its number of undecayed nuclei is 8.0×10^{11} . Find the activity (in Bq) at $t = 0$ s. (2 marks)

.....

.....

.....

.....

.....

- *(iii) Find the percentage decrease in the activity after 200 minutes. (2 marks)

.....

.....

.....

.....

.....

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

- (b) In hospital, technicians wear protective gloves to handle liquid radioactive materials, such as those containing Bi-213 . To monitor radiation exposure, they wear a specialised ring on their finger, as shown in Figure 12.2(a). Each ring contains a small chip in a plastic enclosure (Figure 12.2(b)), which detects and records radiation doses. The rings are collected every three months to monitor the radiation exposure of the technicians.



Figure 12.2(a)




Figure 12.2(b)

- (i) Compared to placing it in a badge worn on the chest, what is the advantage of placing the chip in the ring worn on the hand? (1 mark)

.....

.....

.....

- (ii) The ring should be worn under the gloves. Explain briefly. (1 mark)

.....

.....

.....

- (iii) Which type(s) of radiation (α , β or γ) can reach the chip of the ring? (1 mark)

.....

.....

END OF PAPER

Sources of materials used in this paper will be acknowledged in the *HKDSE Question Papers* booklet published by the Hong Kong Examinations and Assessment Authority at a later stage.

Answers written in the margins will not be marked.

Do not write on this page.

Answers written on this page will not be marked.