

*2. A gas bubble rises from the bottom of a lake to the water surface. Its radius increases from 0.8 cm to 1.0 cm.

- (a) If the gas pressure in the bubble at the water surface is 1.01×10^5 Pa, find the gas pressure in the bubble when it is at the bottom of the lake. Assume that the temperature of the gas in the bubble remains constant. (2 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Use kinetic theory to explain the change in gas pressure in the bubble as its volume increases. (2 marks)

.....

.....

.....

.....

.....

.....

.....

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Book 1 section 4.2: DSE long questions

2012 Q2 (page 2)

Figure 1 shows the variation of the magnetic field B with time t for a coil of N turns and area A placed in a uniform magnetic field. The coil is connected to a circuit containing a resistor of resistance R and a voltmeter. The voltmeter reading is V at time t .

Please stick the barcode label here.

*2. The aqua-lung (a cylinder containing compressed air) for divers has a capacity of $1.0 \times 10^4 \text{ cm}^3$. When the aqua-lung is filled, the air inside has a pressure of 210 atm (atmospheric pressure) at 24°C . The air in the aqua-lung is allowed to expand through a pressure-reducing valve until its pressure equals that of the surrounding water before it is supplied to divers. Assume that the temperature of the air inside the aqua-lung is always equal to that of the surrounding water.

- (a) A diver stays in water of temperature 24°C and pressure 2.0 atm at a depth of 10 m. Find the total volume of air (in cm^3) available for the diver from the aqua-lung at this water pressure. (2 marks)

- (b) The supply of air in (a) is sufficient for the diver to remain at such a depth for 1 hour.


- (i) If the diver breathed in the same volume V_0 (in cm^3) of air per minute, find V_0 . (1 mark)

- (ii) If the diver dives deeper where the water is of temperature 20°C and pressure 4.5 atm, estimate how long (in minutes) the air in a fully-filled aqua-lung would last. Assume that the diver breathes in the **same volume of air** per minute as that found in (b)(i). (3 marks)

Answers written in the margins will not be marked.

*2.

Figure 2.1



Judy uses the set-up shown in Figure 2.1 to study the relationship between the pressure and volume of a fixed mass of gas at constant temperature. The volume V of the gas trapped is read directly from the syringe and the corresponding pressure p is measured by a data-logger via a pressure sensor.

- (a) The initial volume and pressure of the gas are $6.0 \times 10^{-5} \text{ m}^3$ and $1.0 \times 10^5 \text{ Pa}$ respectively at a room temperature of 25°C . Estimate the number of gas molecules trapped in the syringe. (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.


Answers written in the margins will not be marked.

Please stick the barcode label here.

(b) The piston is then pushed in or pulled out to vary V and p such that several pairs of readings are recorded.

Figure 2.2 shows the graph of V against $\frac{1}{p}$ plotted.

Figure 2.2



(i) State ONE experimental precaution for keeping the gas temperature constant.

(1 mark)

Answers written in the margins will not be marked.

(ii) The straight line graph does not pass through the origin but cuts the vertical axis at $-V_0$ instead. Suggest what V_0 stands for.

(1 mark)

Answers written in the margins will not be marked.

(iii) If the experiment is repeated at a higher room temperature using this set-up with the same mass of the same gas, sketch the expected graph in Figure 2.2.

(2 marks)

Answers written in the margins will not be marked


Book 1 section 4.2: DSE long questions

2016 Q2 (page 3)

ANSWERS WRITTEN IN THE MARGINS WILL NOT BE MARKED.

2016-DSE-PHY 1B-5


5



Please stick the barcode label here.

- *2. Two vessels, A and B , of volumes $3V$ and $2V$ respectively are connected by a narrow tube with a tap as shown in Figure 2.1. Initially the tap is closed and both vessels are at the same temperature. Vessel A contains helium gas at a pressure p while vessel B contains 0.8 mol helium gas at a pressure $2p$. Assume that helium gas can be taken as an ideal gas.

Figure 2.1



- (a) Deduce the amount of helium gas (in mol) in vessel A . (2 marks)

.....

.....

.....

.....

.....

- (b) Now the tap is open and a steady state is reached. Assume that the temperature remains unchanged.
- (i) Find the gas pressure inside the vessels in terms of p . (2 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Account for the pressure change of the gas in vessel A using kinetic theory. (2 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....


Answers written in the margins will not be marked.

Book 1 section 4.2: DSE long questions

2018 Q2 (page 2)



- *2. A weather balloon of volume 0.52 m^3 is filled with helium gas of temperature 15°C and pressure 100 kPa at ground level.



- (a) Find the amount of helium gas (in mol) in the balloon.



(2 marks)

.....

.....

.....

- (b) The following graphs show the variation of air temperature T and atmospheric pressure p with height x above ground level.



The weather balloon is released and rises to the upper atmosphere. Assume that the temperature and pressure of the helium gas in the balloon are the same as those of the air outside at any height x .

- (i) A student believes that as the air temperature decreases in the first 10 km, the volume of the balloon decreases. Referring to the graphs above, explain qualitatively why this belief is not correct. (2 marks)

.....

.....

.....

.....

.....

.....

Answers written in the margins will not be marked.

Please stick the barcode label here.

- (ii) In fact the weather balloon keeps on expanding when it rises. The air temperature becomes steady at 216 K from a height of 12 km onwards. When the balloon rises further beyond 12 km and its volume reaches 8 m^3 ,

- (1) estimate the gas pressure in the balloon; (2 marks)


Answers written in the margins will not be marked.

- (2) hence find the corresponding height reached by the balloon. The variation of atmospheric pressure p with height x (in km) is given by


$$p = p_0 e^{-kx},$$

where p_0 is the atmospheric pressure at ground level and $k = 0.138 \text{ km}^{-1}$. (2 marks)

Answers written in the margins will not be marked.



2. Figure 2.1 shows a large gas tank connected with a cylindrical pipe open to the atmosphere. The pipe is fitted with a smooth piston AB . This well-insulated gas tank is filled with high-pressure steam at a temperature of $237\text{ }^{\circ}\text{C}$ under a pressure of $3.10 \times 10^6\text{ Pa}$ while the movable piston is held stationary by a force F_p .
Given: atmospheric pressure = $1.0 \times 10^5\text{ Pa}$



- (a) (i) On Figure 2.1 indicate the force F_p . (1 mark)

- *(ii) By considering the force acting on the piston due to the difference in pressure, find the value of F_p .
The piston has a cross-sectional area of 0.67 m^2 . (2 marks)

Answers written in the margins will not be marked.


- *(iii) Estimate the volume of the gas tank which contains 570 kg steam. You may treat the steam as an ideal gas. Given: mass of one mole of steam = 0.018 kg . (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Please stick the barcode label here.

- (b) This set-up can be used as a 'steam catapult' to launch jet fighters from an aircraft carrier. A jet fighter in position to be launched is connected to the piston via an inextensible cable as shown in Figure 2.2. When the holdback behind the jet fighter is released, the high-pressure steam in the gas tank expands and pushes the piston which in turn helps to accelerate the jet fighter.



In a trial run of the catapult, a jet fighter (with its engine shut down) acquires a final speed of 54 m s^{-1} in 1.5 s after running a distance horizontally on the deck. The mass of the jet fighter is $2.6 \times 10^4 \text{ kg}$.

- (i) Find the work done by the net force on the jet fighter during launch. (2 marks)


Answers written in the margins will not be marked.

- (ii) Calculate the average acceleration of the jet fighter during launch. (2 marks)


Answers written in the margins will not be marked.

- *(iii) State whether the acceleration of the jet fighter is increasing, decreasing or uniform during launch. Explain your answer. (3 marks)

Answers written in the margins will not be marked.



- *2. Figure 2.1 shows a 'water rocket' built from a 2-litre plastic bottle with fins attached. The bottle is filled with water such that 1.5 litres of air is trapped inside.



Initially the trapped air of temperature 27°C is at atmospheric pressure.
 Given: atmospheric pressure = $1.0 \times 10^5 \text{ Pa}$
 density of air = 1.20 kg m^{-3}

- (a) Find the mass of the trapped air m_0 . (2 marks)

Answers written in the margins will not be marked.

Air is then pumped slowly into the rocket until the total mass of trapped air inside the bottle becomes $4m_0$. Assume that the trapped air is kept at a temperature of 27°C (i.e. 300 K) and its volume remains constant.

- (b) (i) Find the pressure of the trapped air. (2 marks)

Answers written in the margins will not be marked.

- (ii) Use kinetic theory to explain why the pressure of the trapped air increases. (2 marks)

Answers written in the margins will not be marked.

Please stick the barcode label here.

- (iii) Given that the area of the water surface is 0.014 m^2 . Estimate the **increase** in the force acting on the water due to the trapped air. (2 marks)


- (c) Given that the mass of the plastic bottle is negligible.

- (i) After the rocket-launching control is pulled and a jet of water ejects from the nozzle, explain why the rocket can go upwards using Newton's laws of motion. (2 marks)

- (ii) **Just** after pulling the control, what is **most** of the energy stored in the pressurized air converted to? (1 mark)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.



Answers written in the margins will not be marked.

- *(b) At a certain location above the Earth in space where the pressure is $2.75 \times 10^{-10} \text{ N m}^{-2}$, 1 m^3 of the gas contains about 1.00×10^{11} molecules. Assuming the gas is ideal, estimate the temperature at the location (in K). (2 marks)

Answers written in the margins will not be marked.

