

Home

Dashboard

Instruction to get API KEY

Overview

Data Report

NTViz's functions

NTViz Tasks

Sections

Provider Instruction

Gemini API key:

.....

👁

Successfully connected to Gemini!

Tasks:

Functions:

Summarize & Goal

NTViz Tasks

 Filter Instruction  Requirements


Instruction: ▾



Select Model:


gemini-1.5-flash ▾

Upload a data file in .csv format:

 Drag and drop file here

Limit 200MB per file • CSV

Browse files

 weather.csv 118.6KB

✕

Successfully uploaded a CSV file with 2922 rows of data.

	location	date	precipitation	temp_max	temp_min	wind	weather
0	Seattle	2012-01-01	0	12.8	5	4.7	drizzle
1	Seattle	2012-01-02	10.9	10.6	2.8	4.5	rain
2	Seattle	2012-01-03	0.8	11.7	7.2	2.3	rain
3	Seattle	2012-01-04	20.3	12.2	5.6	4.7	rain
4	Seattle	2012-01-05	1.3	8.9	2.8	6.1	rain


No missing or duplicate values found in the data.

Generate Charts


★ Insight 1:

<pre>main() Goal(Goal(question='How does average daily temperature (`temp_max`, `temp_min`) vary across locations (`location`) throughout the year (`date`)?', visualization='Line chart showing average daily temperature (calculated as (temp_max + temp_min)/2) over time for each location.', rationale='1. **Data Types:**...'))</pre>	
A visualization goal	
index <code>int</code>	0
question <code>str</code>	'How does average daily temperature (`temp_max`, `temp_min`) vary across locations (`location`) throughout the year (`date`)?'
rationale <code>str</code>	'1. **Data Types:** `temp_max` and `temp_min` are numerical, `date` is datetime, and `location` is categorical.\n2. **Visualization Justification:** A line chart effectively visualizes trends over time for multiple categories. Using the average of `temp_max` and `temp_min` provides a representative...
visualization <code>str</code>	'Line chart showing average daily temperature (calculated as (temp_max + temp_min)/2) over time for each location.'

How does average daily temperature (`temp_max`, `temp_min`) vary across locations (`location`) throughout the year (`date`)?



°*↶•••?↷ Download Chart °*

 VizOps ▾

 Analyze Chart 1


✱ Insight 2:

```
main() Goal Goal(question='What is the relationship between precipitation
(`precipitation`) and maximum temperature (`temp_max`)?', visualization='Scatter plot of
`precipitation` vs. `temp_max` with a trend line.', rationale='1. **Data Types:**
`precipitation` and `temp_max` are both numerical.\n2. **Visualizat...
```

A visualization goal

index	int	1
question	str	'What is the relationship between precipitation (`precipitation`) and maximum temperature (`temp_max`)?'
rationale	str	'1. **Data Types:** `precipitation` and `temp_max` are both numerical.\n2. **Visualization Justification:** A scatter plot is ideal for exploring the relationship between two continuous variables. The trend line helps visualize any correlation.\n3. **Justification:** This visualization reveals if hi...
visualization	str	'Scatter plot of `precipitation` vs. `temp_max` with a trend line.'

What is the relationship between precipitation (`precipitation`) and maximum temperature (`temp_max`)?



°*↶•••?↷ Download Chart °*

⚙ VizOps ▾

🔍 Analyze Chart 2


★ Insight 3:

```
main() Goal Goal(question='How does the distribution of wind speed (`wind`) differ across weather conditions (`weather`)?', visualization='Box plot of `wind` for each
```

category in `weather`.', rationale='1. **Data Types:** `wind` is numerical, and `weather` is categorical.\n2. **Visualization Justification:** A ...

A visualization goal

index	int	2
question	str	'How does the distribution of wind speed (`wind`) differ across weather conditions (`weather`)?'
rationale	str	'1. **Data Types:** `wind` is numerical, and `weather` is categorical.\n2. **Visualization Justification:** A box plot effectively displays the distribution (median, quartiles, outliers) of a numerical variable for different categories. This allows for easy comparison of central tendencies and vari...
visualization	str	'Box plot of `wind` for each category in `weather`.'



[Download Chart](#)

VizOps

✳ Insight 4:

<pre>main() Goal Goal(question='What is the temporal trend of precipitation (`precipitation`) over the entire period (`date`), and can we predict future precipitation levels?', visualization='Line chart of `precipitation` over time (`date`) with a simple linear regression trend line and prediction interval.', ration...</pre>	
A visualization goal	
index <code>int</code>	3
question <code>str</code>	'What is the temporal trend of precipitation (`precipitation`) over the entire period (`date`), and can we predict future precipitation levels?'
rationale <code>str</code>	'1. Data Types: `precipitation` is numerical, and `date` is datetime.\n2. Visualization Justification: A line chart effectively displays trends over time. Adding a linear regression trend line provides a simple predictive element, showing the overall trend and potential future values. A pr...
visualization <code>str</code>	'Line chart of `precipitation` over time (`date`) with a simple linear regression trend line and prediction interval.'



 Analyze Chart 4

```
main() Goal Goal(question='Is there a correlation between minimum temperature
(`temp_min`) and wind speed (`wind`)? How does this vary by location (`location`)?',
visualization='Two separate scatter plots of `temp_min` vs. `wind`, one for each
location (`location`), with trend lines.', rationale="1. **Data Typ...
```

```
index int
```


4

question str

```
'Is there a correlation between minimum temperature (`temp_min`) and  
wind speed (`wind`)? How does this vary by location (`location`)?'

```

<div>rationalestr</div>	<div>"1. **Data Types:** `temp_min` and `wind` are numerical, and `location` is categorical.\n2. **Visualization Justification:** Scatter plots are best for visualizing the relationship between two numerical variables. Creating separate plots for each location allows for a comparison of the relationship..."</div>
<div>visualizationstr</div>	<div>'Two separate scatter plots of `temp_min` vs. `wind`, one for each location (`location`), with trend lines.'</div>



[Download Chart](#)

VizOps

Analyze Chart 5

Chart Analysis

Analysis Report	^
Analysis:	
1. Chart Description:	

- The chart displays the relationship between minimum temperature (`temp_min`) and wind speed (`wind`) for two locations: New York and Seattle.
- It uses two separate subplots, one for each location, allowing for a side-by-side comparison.
- Each subplot contains a scatter plot showing individual data points, with a black trend line overlaid.
- The x-axis represents minimum temperature, and the y-axis represents wind speed. The titles clearly indicate the location of each subplot.

2. Noticeable Patterns:

- **Trend Lines:** The trend lines (linear regressions) show the overall direction of the relationship between temperature and wind speed for each location. The R^2 value (coefficient of determination) displayed on each trend line indicates the strength of the linear relationship. A higher R^2 value (closer to 1) suggests a stronger linear relationship. Without the actual chart image, the precise R^2 values and slopes cannot be stated.
- **Scatter Plot Density:** The density of the scatter points in each subplot gives a visual indication of the data distribution. A tightly clustered scatter plot suggests a strong relationship (though not necessarily linear), while a widely dispersed plot suggests a weaker relationship. Again, specific observations require the chart image.
- **Potential Correlations:** The direction of the trend line indicates the type of correlation. A positive slope suggests a positive correlation (higher temperatures associated with higher wind speeds), while a negative slope indicates a negative correlation (higher temperatures associated with lower wind speeds). The strength of the correlation is indicated by the R^2 value.
- **Outliers:** The chart might reveal outliers—data points significantly deviating from the overall trend. These outliers warrant further investigation to understand their causes.

3. Contextual Insights:

- To provide specific contextual insights, we need the actual R^2 values from the generated chart. However, we can make some general statements based on the provided dataset summary:
- **Data Range:** The `temp_min` ranges from -7.7 to 37.8 degrees, and `wind` ranges from 0.4 to 16.2. This wide range suggests potential variability in the relationship between temperature and wind across different seasons and weather conditions.
- **Location Differences:** The dataset indicates that there are 1461 data points for each location (New York and Seattle). The chart allows us to visually compare how the relationship between temperature and wind

differs between these two geographically distinct locations. We expect to see different patterns due to varying weather systems and geographical features.

- **Crucial Information:** The crucial information lies in the comparison of the R^2 values between New York and Seattle. A significant difference in R^2 would indicate that the relationship between minimum temperature and wind speed is location-dependent. For example, if Seattle shows a stronger negative correlation than New York, it might suggest a stronger influence of temperature on wind patterns in Seattle's climate.
- **Meaningful Insights:** Combining the chart with the knowledge that the dataset includes weather information ("weather" column) allows for a deeper analysis. We could investigate if the relationship between temperature and wind varies significantly across different weather types (rain, sun, fog).

4. External Influences:

- **Seasonality:** Seasonal changes significantly impact both temperature and wind speed. The data likely shows different patterns across seasons (e.g., stronger winds in winter months with lower temperatures).
- **Geographical Factors:** The geographical location of each city (coastal vs. inland, proximity to mountains, etc.) influences local wind patterns and temperature variations. Seattle's coastal location, for instance, might lead to different wind patterns compared to New York's inland location.
- **Large-Scale Weather Systems:** Major weather events (storms, pressure systems) can cause temporary spikes in wind speed regardless of temperature. These events could create outliers in the data.

5. Recommendations:

- **Investigate Location Differences:** Focus on understanding why the relationship between temperature and wind differs (or doesn't) between New York and Seattle. This could involve analyzing geographical factors and typical weather patterns for each location.
- **Seasonal Analysis:** Analyze the data seasonally to understand how the relationship changes over the year. This will provide a more nuanced understanding of the correlation.
- **Weather Type Analysis:** Examine the relationship between temperature and wind for each weather type (rain, sun, fog) separately. This will help determine if specific weather conditions modify the correlation.
- **Outlier Investigation:** Identify and investigate any outliers. Understanding the causes of these outliers can improve the accuracy of future predictions.

6. Conclusion:

The chart provides a visual comparison of the relationship between minimum temperature and wind speed in New York and Seattle. By analyzing the trend lines (specifically their slopes and R^2 values) and the scatter plot density, we can determine the strength and direction of the correlation for each location. Further analysis, incorporating seasonal and weather type data, will provide a more comprehensive understanding of the factors influencing this relationship and lead to more accurate predictions. The key takeaway will be whether the correlation is consistent across locations or significantly influenced by geographical and seasonal factors.
