An online adaptive sampling algorithm for nonsmooth stochastic optimization with difference-of-convex objectives *

Yuhan Ye†

November 27, 2024

Abstract

We propose an adaptive online algorithm for stochastic nonsmooth nonconvex optimization problems with difference-of-convex (DC) objectives that achieves state-of-art sample complexity. The sample size and step size are determined by the current estimates of stochastic quantities. The Sample Average Approximation (SAA) error for the linearized concave parts is closely related to the convergence of the SAA for subdifferential mappings. We derive a novel $O(\sqrt{p/n})$ point-wise rate (modulo logarithmic factors) where p is the dimension and n is the sample size. Our result complements the state-of-art $O(\sqrt[4]{p/n})$ result, matching the same SAA convergence rate in the smooth case. Due to the different convergence rates and DCA's special structure, we should use different sampling sizes to achieve the optimal error-sample size trade-off. Numerical experiments show that our adaptive strategy improves the algorithm's performance and reduces its worst-case sampling size.

Keywords: stochastic, nonsmooth, sampling, adaptive, difference-of-convex, sample average approximation, subdifferential mapping

1 Introduction

Consider the class of nonsmooth nonconvex optimization problems in the form of

$$\underset{x\in C}{\text{minimize}} \quad f(x),$$

where $C \subset \mathbb{R}^p$ is a convex compact set. The nonsmooth stochastic objective function $f : \mathbb{R}^p \to \mathbb{R}$ is given as $f(x) = \mathbb{E}_{\xi,\zeta}[F(x,\xi,\zeta)]$, where the expectation $\mathbb{E}_{\xi,\zeta}$ is the expected value of the function $F(x,\xi,\zeta)$ with respect to ξ and ζ. The random vector ξ and ζ is defined over probability spaces $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ and $(\Omega_2, \mathcal{F}_2, \mathbb{P}_2)$, respectively. In this work, we focus on $F : (\mathbb{R}^p, \Omega_1, \Omega_2) \to \mathbb{R}$, which are difference-of-convex functions with the decomposition

$$F(x,\xi,\zeta) = G(x,\xi) - H(x,\zeta),$$

where $G(x,\xi)$ and $H(x,\zeta)$ are Carathéodorian, i.e., it is continuous in $x \in C$ for all $\xi \in \Omega_1$, $\zeta \in \Omega_2$ and Borel measurable in ξ and ζ for all $x \in C$.

Let $g = \mathbb{E}_\xi[G(x,\xi)]$ and $h = \mathbb{E}_\zeta[H(x,\zeta)]$. The objective $f(x) = g(x) - h(x)$ itself is a difference-of-convex function.

We consider the above stochastic DC program in the online fashion: at each iteration, new data is used to construct a stochastic approximation of G and one of a subgradient of H, and previous samples are no longer available. Since the solution of the proposed algorithms is updated via DCA with proximal terms and the update steps require new fresh samples from the distribution of ξ and ζ, we refer to our algorithm as osproxDCA. The proposed osproxDCA enjoys the double benefit of an online algorithm: it is suitable to perform streaming data from an unknown distribution. Therefore, even if the distribution of ξ and ζ changes at a certain time (suppose that, due to some real-world events, ξ becomes ξ' or ζ becomes ζ' at the k-th iteration), the algorithm will automatically solve the problem with the updated ξ' and ζ'.

We summarize the contribution of the paper as follows:

*Submitted to the editors.

†School of Mathematics, Peking University, Beijing, China (2100010664@stu.pku.edu.cn).

- We derive a novel $O(\sqrt{p/n})$ convergence rate (modulo logarithmic factors) in expectation of the pointwise SAA error for subdifferential mappings, matching the same rate of differential mappings in the smooth case, complementing the state-of-art $O(\sqrt[4]{p/n})$ result, where p is the dimension and n is the sample size. Compared to the previous work [54, 21] under uniform setting, we employed a new and simple proof technique by transforming the one-sided deviation of subdifferential set-valued functions into the SAA form of support functions..
- Based on this, we propose an adaptive online stochastic proxDCA framework complementing previous work [33, 35]. Under the same setting as [35], our work improves either the sampling scale or the assumptions required.
- From the perspective of practical numerical experiments, our adaptive strategy improves the algorithm's performance and reduces its worst-case sampling size.

1.1 Related Literature

Sample Average Approximation (SAA) and Convergence of Subdifferential Mappings This paper draws on the extensive body of work surrounding Sample Average Approximation (SAA) within the realms of stochastic programming and statistical learning. While the results in both fields share many similarities, they primarily differ in their focus on various convergence criteria. In stochastic programming, significant attention has been dedicated to the convergence of (approximate) minimizers and optimal values, with considerations of distributional and almost sure limits [32, 58, 25, 56, 51, 49, 57, 31, 53]. In contrast, the statistical learning community has typically focused more on deriving excess risk bounds that hold with high probability, often independent of the problem's dimensionality [55, 27, 61, 38, 39, 40, 10, 50, 68, 76, 62, 30].

Several studies have investigated (sub)gradient-based convergence, which is also the focus of our work. For instance, [72] demonstrates non-asymptotic, dimension-dependent high-probability bounds on the distance between the empirical and population subdifferentials under the Hausdorff metric. However, the main assumption in this paper essentially requires the population objective to be smooth. Alternatively, [73] directly smooths the empirical loss function $f(x, z)$ and shows that the gradient of this smoothed empirical risk converges to an element of the population subdifferential. However, no finite-sample bounds are provided in [73]. The paper [41] discusses uniform convergence of gradients for smooth objectives under the assumption that the gradient $\nabla f(x, z)$ is sub-Gaussian with respect to the population data. The bounds established in [41] are dimension-dependent and rely on covering arguments. More recently, [23] provides dimension-independent high-probability convergence rates of gradients for smooth Lipschitz generalized linear models, utilizing a "chain rule" for Rademacher complexity. The resulting rates are of the form $n^{-1/2}$ for smooth generalized linear models. However, these works do not directly examine the convergence behavior of subdifferential sets.

The most general asymptotic convergence result that we are aware of is given in [59]. This study proves that, with probability one, the limit of an enlarged empirical subdifferential converges to the enlarged subdifferential of the population risk under the Hausdorff metric. In the uniform setting, [54] achieves a tight $\sqrt{p/n}$ rate (modulo logarithmic factors) for the uniform convergence of subdifferential mappings, where p is the dimension and n is the sample size. This result is based on a weaker notion of subdifferential convergence and assumes stronger conditions [54, Assumption C]. This complements the $\sqrt[4]{p/n}$ rate in [21], which is based on the graphical distance between subdifferentials, derived using Attouch's epigraphical convergence theorem [2] and Moreau envelope smoothing [44]. However, the method used in [21] often leads to suboptimal sample complexity guarantees [21, Section 5].

Our work advances the literature on the convergence of subdifferential mappings in the context of Sample Average Approximation. We achieve the $O(\sqrt{p/n})$ rate, as found in [54], but without relying on the assumption [54, Assumption C]. Moreover, our result surpasses the $O(\sqrt[4]{p/n})$ rate in [21], offering a new perspective for establishing convergence of subdifferential mappings. Specifically, our approach applies to stochastic weakly-convex classes, which are central to many data-driven applications [20].

Stochastic and Online DC Optimization DC programming, first introduced by [66], has been the subject of extensive study over the years. A thorough review of DC programming's evolution can be found in [36]. While there has been considerable work on DC programming, its application in stochastic settings has only recently gained attention. The majority of existing research on stochastic DC optimization builds on the classical DC Algorithm (DCA) used in deterministic DC problems. The central idea behind DCA is to approximate the DC problem as a convex problem by linearizing the second component. This results in a

double-loop algorithm, where the update rule is given by $\min_x \{\phi(x) - \langle \nabla\phi(x_k), x\rangle\}$. The method was first extended to stochastic DC problems (SDCA) in [67] for large-sum problems involving non-convex smooth functions, which was later generalized to handle large-sum non-smooth problems in [34]. The first work that allowed both components in a DC problem to be non-smooth was presented in [33], where a SDCA scheme was proposed that stores all past information for constructing future subproblems. This approach achieves near-optimal sample complexity by adding just one sample per DCA subproblem. In [35], the efficiency of the SDCA scheme was improved by eliminating the need to store historical information, instead resampling multiple samples at each DCA subproblem and using Sample Average Approximation (SAA) to approximate the linearized DC function. The resulting algorithms exhibit adaptive capabilities, a significant advantage over those proposed in [33]. However, the approach in [35] relies on strong assumptions, as outlined in [35, Assumption 1]. Our work is conducted under similar conditions to [35], improving upon the sampling scale and the assumptions required.

Moreover, non-asymptotic convergence analysis for stochastic DC optimization has been explored in [46, 74]. These studies propose stochastic proximal DC algorithms (SPD), which modify SDCA by adding an extra quadratic term after linearizing the second component. The main limitation of these analyses is their reliance on the smoothness or Hölder continuity of the gradient, assumptions that are often too strong for many commonly used non-smooth functions. Recent work in non-smooth weakly-convex optimization [19, 63, 45, 75] introduced Moreau envelope smoothing approximations for both components, leading to non-asymptotic convergence analysis to nearly ϵ-critical points under deterministic settings. However, these works have yet to establish non-asymptotic convergence for non-smooth DC problems, as a gap remains between nearly ϵ-critical points and true critical points.

Adaptive Sampling in Stochastic Optimization Adaptive sampling is an effective approach in stochastic optimization, designed to manage the variance in approximations throughout the optimization process. The core concept is intuitive yet impactful: when the current iterate is far from the optimal solution, it is sufficient to use less precise and computationally inexpensive (gradient) information. However, as the algorithm approaches the solution, higher accuracy in (gradient) estimates becomes essential for both theoretical guarantees and practical performance. The effectiveness of such methods lies in the strategy used to determine the sample size or approximation accuracy.

Algorithms with predefined (often geometric) rules for increasing sample sizes have been explored in works such as [24, 11]. Among these, [11] demonstrates that such methods achieve the optimal worst-case first-order complexity in unconstrained optimization. Additionally, adaptive sampling methods offer advantages over fixed-sample approaches, such as leveraging parallelism and generating iterates with reduced variance due to progressively increasing sample sizes.

Other adaptive strategies rely on gradient approximation tests to regulate the accuracy of approximations. Examples include norm-based tests [12, 11], inner product tests [8, 9], and other techniques [13, 6, 29]. Moreover, adaptive sampling has been successfully applied to simulation-based optimization problems [28, 48]. More recently, methods for constrained stochastic optimization problems with convex feasible sets have been proposed in [4, 71]. For a comprehensive overview of adaptive sampling techniques, refer to [18].

2 Background and notations

We first summarize the notation used throughout the paper. We write $\mathbb{R}^p$ as the p-dimensional Euclidean space equipped with the inner product $\langle x, y\rangle = x^\top y$ and the induced norm $\|x\| \triangleq \sqrt{x^\top x}$. The symbol $\mathbb{B}(x, \delta)$ is used to denote the closed ball of radius $\delta > 0$ centered at a vector $x \in \mathbb{R}^p$. Let A and C be two nonempty subsets of $\mathbb{R}^n$. The distance from a vector $x \in \mathbb{R}^n$ to A is defined as $\operatorname{dist}(x, A) \triangleq \inf_{y\in A} \|y - x\|$. The one-sided deviation of A from C is defined as $\mathbb{D}(A, C) \triangleq \sup_{x\in A} \operatorname{dist}(x, C)$. The Hausdorff distance between A and C is defined as $\mathbb{H}(A, C) := \max\{\mathbb{D}(A, C), \mathbb{D}(C, A)\}$.

Next, we introduce the concepts of **generalized derivatives and subdifferentials for nonsmooth functions**. Interested readers are referred to the monographs [16, 52, 43] for thorough discussions on these subjects. Consider a function $r : \mathcal{O} \to \mathbb{R}$ defined on an open set $\mathcal{O} \subseteq \mathbb{R}^p$. The classical one-sided directional derivative of r at $\bar{x} \in \mathcal{O}$ along the direction $d \in \mathbb{R}^p$ is defined as $r'(\bar{x}; d) \triangleq \lim_{t\downarrow 0} \dfrac{r(\bar{x} + td) - r(\bar{x})}{t}$ if this limit exists. The function r is said to be directionally differentiable at $\bar{x} \in \mathcal{O}$ if it is directionally differentiable

along any direction $d \in \mathbb{R}^p$. In contrast, the Clarke directional derivative of r at $\bar{x} \in \mathcal{O}$ along the direction $d \in \mathbb{R}^p$ is defined as $r^\circ(\bar{x}; d) \triangleq \limsup_{x \to \bar{x},\, t \downarrow 0} \dfrac{r(x+td) - r(x)}{t}$, which is finite when r is Lipschitz continuous near $\bar{x}$. The Clarke subdifferential of r at $\bar{x}$ is the set $\partial_C r(\bar{x}) \triangleq \{v \in \mathbb{R}^n \mid r^\circ(\bar{x}; d) \geq v^\top d \text{ for all } d \in \mathbb{R}^p\}$, which coincides with the usual subdifferential in convex analysis for a convex function. If r is strictly differentiable at $\bar{x}$, then $\partial_C r(\bar{x}) = \{\nabla r(\bar{x})\}$. We say that r is Clarke regular at $\bar{x} \in \mathcal{O}$ if r is directionally differentiable at $\bar{x}$ and $r^\circ(\bar{x}; d) = r'(\bar{x}; d)$ for all $d \in \mathbb{R}^p$. This Clarke regularity at $\bar{x}$ is equivalent to have $r(x) \geq r(\bar{x}) + \bar{v}^\top(x - \bar{x}) + o(\|x - \bar{x}\|)$ for any $\bar{v} \in \partial_C r(\bar{x})$. Therefore, if a function fails to satisfy the Clarke regularity at $\bar{x}$, there does not exist an approximate linear lower bound of the original function based on the Clarke subdifferentials with a small o error locally.

Then, we introduce the concepts of **outer semicontinuity**. Let $\mathcal{A} : \mathbb{R}^p \rightrightarrows \mathbb{R}^m$ be a set-valued mapping. Its outer limit at $x \in \mathbb{R}^n$ is defined as $\limsup_{x \to \bar{x}} \mathcal{A}(x) := \bigcup_{x^\nu \to \bar{x}} \limsup_{\nu \to \infty} \mathcal{A}(x^\nu) = \big\{u \mid \exists\, x^\nu \to \bar{x}, \exists\, u^\nu \to u \text{ with } u^\nu \in \mathcal{A}(x^\nu)\big\}$. We say $\mathcal{A}$ is outer semicontinuous (osc) at $\bar{x} \in \mathbb{R}^n$ if $\limsup_{x \to \bar{x}} \mathcal{A}(x) \subseteq \mathcal{A}(\bar{x})$. Clarke subdifferential is outer semicontinuous, which is necessary for establishing subsequential convergence [52, Proposition 6.6]. In addition, for a L_r-Lipschitz r, $\partial r(\bar{x})$ is locally bounded by L_r [52, Theorem 9.13]. For any $\varepsilon > 0$, we denote the Goldstein subdifferential as $\partial^\varepsilon r(x) := \bigcup_{x' \in \mathbb{B}(x,\varepsilon)} \partial r(x')$., which is also outer semicontinuous [26],

We next review some basics about **random set-valued mappings and their expectation**. Let $(\Omega, \mathcal{F}, P)$ be a probability space. For fixed x, let $\mathcal{A}(x, \omega) : \Omega \to 2^{\mathbb{R}^n}$ be a set-valued mapping which takes values of closed subsets in $\mathbb{R}^n$. Let $\mathfrak{B}(\mathbb{R}^n)$ or simply $\mathfrak{B}$ denote the space of closed bounded subsets of $\mathbb{R}^n$ endowed with topology $\tau_{\mathbb{H}}$ generated by the Hausdorff distance $\mathbb{H}$. We consider the Borel σ-field $\mathcal{G}(\mathfrak{B}, \tau_{\mathbb{H}})$ generated by the $\tau_{\mathbb{H}}$-open subsets of $\mathfrak{B}$. A set-valued mapping $\mathcal{A}(x, \omega) : \Omega \to 2^{\mathbb{R}^n}$ is said to be $\mathcal{F}$-measurable if, for every member $\mathcal{W}$ of $\mathcal{G}(\mathfrak{B},\ \tau_{\mathbb{H}})$, one has $\mathcal{A}^{-1}(\mathcal{W}) \in \mathcal{F}$. By a measurable selection of $\mathcal{A}(x, \omega)$, we refer to a vector $A(x, \omega) \in \mathcal{A}(x, \omega)$, which is measurable. Note that such measurable selections exist if $\mathcal{A}(x, \omega)$ is measurable, see [1] and references therein. For a general set-valued mapping which is not necessarily measurable, the expectation of $\mathcal{A}(x, \omega)$, denoted by $\mathbb{E}[\mathcal{A}(x, \omega)]$, is defined as the collection of $\mathbb{E}[A(x, \omega)]$ where $A(x, \omega)$ is an integrable selection. The integrability is in the sense of Aumann [3]. $\mathbb{E}[\mathcal{A}(x, \omega)]$ is regarded as well defined if $\mathbb{E}[\mathcal{A}(x, \omega)] \in \mathfrak{B}$ is nonempty. A sufficient condition of the well-definedness of the expectation is that $\mathcal{A}(x, \omega)$ is measurable and $\mathbb{E}[\|\mathcal{A}(x, \omega)\|] := \mathbb{E}[\mathbb{H}(0, \mathcal{A}(x, \omega))] < \infty$, see [3, Theorem 2]. Let $r(x, \xi) : \mathbb{R}^n \times \Xi \to \mathbb{R}$ be a random lsc function, where $\xi : (\Omega, \mathcal{F}, P) \to \Xi$ is a random vector with support set $\Xi \subset \mathbb{R}^m$. Suppose: (a) r is Lipschitz w.r.t. x for every ξ with modulus $\kappa(\xi)$, where $\kappa : \Xi \to \mathbb{R}_+$ is a measurable function that satisfies $\mathbb{E}[\kappa(\xi(\omega))] < \infty$, (b) for any $x \in \mathbb{R}^n$, $r(x, \xi)$ is Clarke regular at x for a.e. $\xi \in \Xi$. Then at every $x \in \mathbb{R}^n$, $\mathbb{E}[r(x, \xi(\omega))]$ is also Clarke regular and $\partial_x \mathbb{E}[r(x, \xi(\omega))] = \mathbb{E}\left[\partial_x r(x, \xi(\omega))\right]$ [16, Theorem 2.7.2].

Finally, we introduce **DC criticality**. A point $x^* \in \mathbb{R}^p$ is called *a DC critical* point if $0 \in \partial g(x^*) - \partial h(x^*)$, or equivalently $\partial g(x^*) \cap \partial h(x^*) \neq \emptyset$, while it is called *a strongly DC critical point* of $(\mathcal{P})$ if $\emptyset \neq \partial h(x^*) \subset \partial g(x^*)$. In the framework of DC programming, the terminology "critical point" is referred to the notion of DC criticality. The notion of DC criticality is close to Clarke, Fréchet, limiting stationarity in the sense $\partial^F f(x^*) \subseteq \partial f(x^*) \subseteq \partial g(x^*) - \partial h(x^*)$ (whenever h is continuous at x^*) and $\partial^C f(x^*) \subseteq \partial g(x^*) - \partial h(x^*)$, where equalities hold under technical assumptions. Furthermore, the commonly used *directional stationarity* is equivalent to strong DC criticality, which is the strongest necessary condition for local DC optimality.

3 osproxDCA framework and SAA error analysis

We first make the following assumptions on $G(x, \xi)$ and $H(x, \zeta)$:

Assumption 1. *(Assumptions For Problem)*

1. *The feasible region C is convex and close, and $f(x) > f_m$ is lower bounded.*
2. *For almost all $\xi \in \Omega_1$, $G(x,\xi)$ is ρ_g-convex (possibly weakly convex if $\rho_g < 0$) over the feasible region C, i.e., for all $x, y \in C$ and $\alpha \in [0,1]$, we have:*

$$G(\alpha x + (1-\alpha)y, \xi) \leq \alpha G(x,\xi) + (1-\alpha)G(y,\xi) - \frac{\rho_g}{2}\alpha(1-\alpha)\|x-y\|^2.$$

   Similarly, $H(x,\zeta)$ is ρ_h-convex (possibly weakly convex ? if $\rho_h < 0$) over C for almost all $\zeta \in \Omega_2$.
3. *For almost all $\xi \in \Omega_1$, $G(x,\xi)$ is L_g-Lipschitz continuous over C, i.e., for all $x, y \in C$, we have:*

$$\|G(x,\xi) - G(y,\xi)\| \leq L_g\|x-y\|.$$

   Similarly, $H(x,\zeta)$ is L_h-Lipschitz continuous over C for almost all $\zeta \in \Omega_2$.

3.1 osproxDCA framework

The DC algorithm is an iterative method with a locally approximated convex objective. At iteration k and iterate x_k, the local approximation model to the objective with the exact g and a true subgradient $y_k \in \partial h(x_k)$ is

$$m_k(x) = g(x) - h(x_k) - y_k^T(x - x_k) + \frac{1}{2}\mu_k\|x - x_k\|^2, \tag{1}$$

where $\mu_k > 0$ is a scalar quadratic coefficient. In the stochastic setting, we use stochastic estimate $\bar{g}_k(\cdot)$ of $g(\cdot)$ and a stochastic subgradient estimate $\bar{y}_k$ of y_k in the local model $\bar{m}_k(\cdot)$:

$$\bar{m}_k(x) = \bar{g}_k(x) - \bar{h}_k(x_k) - \bar{y}_k^T(x - x_k) + \frac{1}{2}\mu_k\|x - x_k\|^2. \tag{2}$$

The stochastic estimate of function value $\bar{h}_k$ does not affect the optimization subproblem and its solution. Hence, we hereby use it interchangeably with $h(x_k)$. Denoting $d = x - x_k$, $\bar{m}_k(x)$ and $m_k(x)$ can be rewritten as $\bar{M}_k(d)$ and $M_k(d)$, respectively,

$$\begin{aligned} \bar{M}_k(d) &= \bar{g}_k(x_k + d) - \bar{h}_k(x_k) - \bar{y}_k^T d + \frac{1}{2}\mu_k\|d\|^2, \\ M_k(d) &= g(x_k + d) - h(x_k) - y_k^T d + \frac{1}{2}\mu_k\|d\|^2. \end{aligned} \tag{3}$$

The subproblem to be solved at iteration k is

$$\begin{aligned} \underset{d}{\text{minimize}} \quad & \bar{M}_k(d) \\ \text{subject to} \quad & x_k + d \in C. \end{aligned} \tag{4}$$

The first-order optimality conditions of subproblem (4) are

$$\begin{aligned} \bar{z}_{k+1} - \bar{y}_k + \mu_k\bar{d}_k + \bar{v}_k &= 0, \\ x_k, x_k + \bar{d}_k \in C, \bar{v}_k &\in \partial i_C(x_k + \bar{d}_k), \end{aligned} \tag{5}$$

where $\bar{z}_{k+1} \in \partial\bar{g}_k(x_k + \bar{d}_k)$.

The stochastic quantities are estimated at each iteration with independent and identically distributed (i.i.d.) general samples of ξ and ζ. We use $S_{g,k}$ to denote the set of samples of ξ at iteration k when obtaining estimation $g(x)$, and $S_{h,k}$ to denote the set of samples of ζ at iteration k when obtaining estimation $\bar{y}_k$. The cardinality or sample size of $S_{g,k}, S_{h,k}$ is defined as $N_{g,k} = |S_{g,k}|$ and $N_{h,k} = |S_{h,k}|$. Let $\mathcal{F}_k \triangleq \sigma\left(S_{g,1}, S_{h,1}, S_{g,2}, S_{h,2}, \ldots, S_{g,k-1}, S_{h,k-1}\right)$ be a filtration, i.e., an increasing sequence of σ-fields generated by samples used in the previous $k-1$ iterations.

Our proposed stochastic DC algorithm framework is presented in Algorithm 1, while the exact rule when updating $\mu_k, N_{g,k}, N_{h,k}$ is analyzed and presented in Section 4 and 5. Additionally, the quadratic term $\mu_k\|d\|^2$ can be replaced by $d^T B_k d$, where B_k is a positive definite matrix.

It is trivial to verify that $\bar{g}_k(x)$ is L_g-Lipschitz, $\bar{h}_k(x)$ is L_h-Lipschitz, and $\|\bar{y}_k\| \leq L_h$.

Algorithm 1 osproxDCA framework

1: Initialize $x_0, \mu_0, N_{g,0}, N_{h,0}$.
2: **for** $k = 0, 1, 2, \cdots$ **do**
3: Generate iid samples $S_{g,k} = \{\xi^{k,i}\}_{i=1}^{N_{g,k}}$ and $S_{h,k} = \{\zeta^{k,i}\}_{i=1}^{N_{h,k}}$ from the distribution of ξ and ζ that are independent of previous samples.
4: Construct the local approximation model $\bar{M}_k(d)$ by SAA:
Set $\bar{g}_k(x) = \frac{1}{N_{g,k}} \sum_{i=1}^{N_{g,k}} G\left(x, \xi^{k,i}\right)$, $\bar{h}_k(x) = \frac{1}{N_{h,k}} \sum_{i=1}^{N_{h,k}} H\left(x, \zeta^{k,i}\right)$, and select $\bar{y}_k \in \partial \bar{h}_k(x) = \frac{1}{N_{h,k}} \sum_{i=1}^{N_{h,k}} \partial_x H\left(x, \zeta^{k,i}\right)$.
5: Solve the $\rho_g + \mu_k$-convex optimization subproblem (4) to obtain $\bar{d}_k$.
6: Take the step $x_{k+1} = x_k + \bar{d}_k$.
7: Call the $\mu_k, N_{g,k}, N_{h,k}$ update rules to obtain $\mu_{k+1}, N_{g,k+1}, N_{h,k+1}$.
8: **end for**

Remark 1. *If there exists an isomorphic mapping ϕ from $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ to $(\Omega_2, \mathcal{F}_2, \mathbb{P}_2)$, Step 3 in Algorithm 1 could be simplified to:*

If $N_{g,k} \geq N_{h,k}$,

1. *generate iid samples $S_{g,k} = \{\xi^{k,i}\}_{i=1}^{N_{g,k}}$ from the distribution of ξ that are independent of previous samples;*
2. *for $i = 1, 2, \cdots, N_{h,k}$, let $\zeta^{k,i} = \phi(\xi^{k,i})$ and $S_{h,k} = \{\zeta^{k,i}\}_{i=1}^{N_{h,k}}$.*

The counterpart is similar.

3.2 Uniform convergence of the SAA for g in expectation

To continue, we first introduce a lemma regarding the convergence rate of the sample average approximation in expectation. This result is derived from the Rademacher average of the random function $\psi(x;\omega)$, as discussed in [22, Corollary 3.2] and further explored in [17, Theorem 10.1.5]. For a random vector function $\psi(\cdot,\omega) : \mathcal{X} \to \mathbb{R}$ and i.i.d. random variables $(\omega^1, \ldots, \omega^n) = \bar{\omega}^n$ drawn from the distribution of ω, we define the quantity

$$\delta_n\left(\psi, \bar{\omega}^n\right) := \sup_{x \in \mathcal{X}} \left\| \frac{1}{n} \sum_{i=1}^{n} \boldsymbol{\psi}\left(x, \omega^i\right) - \mathbb{E}\boldsymbol{\psi}(x,\omega) \right\|.$$

We have the following basic estimates [22, Theorem 3.1].

Lemma 1. *(Basic Estimates). If functions $\psi(\cdot,\omega)$ are bounded by constant M and Lipschitz continuous with constant L in the first variable $x \in \mathcal{X} \subseteq [0,r]^p \subseteq \mathbb{R}^d$ uniformly in ω, then for any $\alpha \in (0, 1/2)$ it holds that*

$$\begin{aligned}
&\mathbb{E}_{\bar{\omega}^n} \delta_n\left(\psi, \bar{\omega}^n\right) \leq 2\sqrt{p}(Lr + M/\sqrt{(1-2\alpha)e})/n^{\alpha}, \\
&\mathbb{P}\left\{\sqrt{n}\left|\delta_n\left(\psi, \bar{\omega}^n\right) - \mathbb{E}_{\bar{\omega}^n} \delta_n\left(\psi, \bar{\omega}^n\right)\right| \geq t\right\} < 2\exp\left\{-t^2/\left(2M^2\right)\right\} \quad \forall t \geq 0.
\end{aligned}$$

By Lemma 1, we could derive the SAA error in expectation for g.

Theorem 1. *Assume that $\bar{d}_k \in [-r_k, r_k]^d$, for any $\alpha \in (0, 1/2)$,*

$$\mathbb{E}_{S_{g,k}} \left|\bar{g}_k(x_{k+1}) - \bar{g}(x_k) - g(x_{k+1}) + g_k(x_k)\right| \leq \frac{C_g}{\mu_k N_{g,k}^{\alpha}},$$

where $C_g \geq 4\sqrt{p} L_g (L_g + L_h)\left(2 + \frac{L_g}{\sqrt{(1-2\alpha)e}}\right)$.

Proof. First we prove that $\|\bar{d}_k\| \leq \frac{2(L_g+L_h)}{\mu_k}$. Since $\bar{d}_k$ is the solution of subproblem (4), we have

$$\bar{g}_k(x_k + \bar{d}_k) - \bar{h}_k(x_k) - \bar{y}_k^T \bar{d}_k + \frac{1}{2}\mu_k \|\bar{d}_k\|^2 = \bar{M}_k(\bar{d}_k) \leq \bar{M}_k(0) = \bar{g}_k(x_k) - \bar{h}_k(x_k).$$

Hence we have $\frac{1}{2}\mu_k\|\bar{d}_k\|^2 \leq \bar{g}_k(x_k) - \bar{g}_k(x_k+\bar{d}_k) + \bar{y}_k^T\bar{d}_k \leq \|\bar{d}_k\|(L_g+L_h)$, which implies $\|\bar{d}_k\| \leq \frac{2(L_g+L_h)}{\mu_k}$. Let $r_k = \frac{2(L_g+L_h)}{\mu_k}$ and $\psi(x,\xi) = G(x,\xi) - G(x_k,\xi)$, which is L_g-Lipschitz. By Lemma 1,

$$\mathbb{E}_{Sg,k} \sup_{\Delta_x \in [-r_k, r_k]^d} \left| \frac{1}{n}\sum_{i=1}^{n} \psi\left(x_k + \Delta_x, \xi^{k,i}\right) - \mathbb{E}_\xi \psi(x_k+\Delta_x,\xi) \right| \leq \frac{\hat{C}_g}{N_{g,k}^{\alpha}},$$

where $\hat{C}_g = 2\sqrt{p}(2L_g r_k + L_g r_k/\sqrt{(1-2\alpha)e})$. Noticed that

$$|\bar{g}_k(x_{k+1}) - \bar{g}(x_k) - g(x_k) + g_k(x_k)| \leq \sup_{\Delta_x \in [-r_k, r_k]^d} \left| \frac{1}{n}\sum_{i=1}^{n} \psi\left(x_k + \Delta_x, \xi^{k,i}\right) - \mathbb{E}_\xi \psi(x_k+\Delta_x,\xi) \right|,$$

Hence, we finish the proof by substituting r_k for its definition. □

3.3 Convergence of the SAA for subdifferential mapping ∂h in expectation

In the following discussion, we investigate the convergence rate of the sample average subdifferential mapping $\partial\bar{h}_k(x) = \frac{1}{N_{k,h}}\sum_{k=1}^{N} \partial_x h\left(x, z^k\right)$ to $\partial h(x) = \mathbb{E}_\zeta\left[\partial_x h(x,\zeta)\right]$ in expectation. To achieve this, we analyze the quantity $\mathbb{D}\left(\frac{1}{N}\sum_{k=1}^{N} \partial_x H\left(x,\zeta^k\right), \mathbb{E}_\zeta\left[\partial_x H(x,\zeta)\right]\right)$ converging to 0 in expectation as N increases.

Define the support function of a set S as $\sigma(u,S) := \sup_{s\in S} u^T s$. It is well known that $\sigma(u,S) = \sigma(u, \operatorname{conv} S)$ and for nonempty sets S and S', the following holds:

$$\sigma\left(u, S+S'\right) = \sigma(u,S) + \sigma\left(u,S'\right), \tag{6}$$

as discussed in [15, p. 621]. Furthermore, the well-established Hömander's formula [14, Theorem II-18] states that for any two nonempty convex and compact subsets A and B of $\mathbb{R}^n$:

$$\mathbb{D}(A,B) = \max_{\|u\|\leqslant 1}\left(\sigma(u,A) - \sigma(u,B)\right). \tag{7}$$

Using the above formula, we have the following lemma that converts our targeted quantity into the form of an SAA error; see, e.g., [72, Theorem 4.3].

Lemma 2. *For any $x \in C$, we have*

$$\mathbb{D}\left(\partial\bar{h}_k(x), \partial h(x)\right) = \max_{\|u\|\leqslant 1}\left[\frac{1}{N_{k,h}}\sum_{i=1}^{N_{k,h}} \sigma\left(u, \partial_x H\left(x,\zeta^{k,i}\right)\right) - \mathbb{E}_\zeta\left[\sigma\left(u, \partial_x H(x,\zeta)\right)\right]\right].$$

Proof.

$$\begin{aligned}
&\mathbb{D}\left(\frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \partial_x H\left(x,\zeta^k\right), \operatorname{conv}\mathbb{E}_\zeta\left[\partial_x H(x,\zeta)\right]\right) \\
&= \mathbb{D}\left(\operatorname{conv}\left\{\frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \partial_x H\left(x,\zeta^{k,i}\right)\right\}, \operatorname{conv}\mathbb{E}_\zeta\left[\partial_x H(x,\zeta)\right]\right) \\
&= \sup_{\|u\|\leqslant 1} \sigma\left(u, \frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \partial_x H\left(x,\zeta^{k,i}\right)\right) - \sigma\left(u, \mathbb{E}_\zeta\left[\partial_x H(x,\zeta)\right]\right) \\
&= \sup_{\|u\|\leqslant 1}\left[\frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \sigma\left(u, \partial_x H\left(x,\zeta^{k,i}\right)\right) - \sigma\left(u, \mathbb{E}_\zeta\left[\partial_x H(x,\zeta)\right]\right)\right] \\
&= \sup_{\|u\|\leqslant 1}\left[\frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \sigma\left(u, \partial_x H\left(x,\zeta^{k,i}\right)\right) - \mathbb{E}_\zeta\left[\sigma\left(u, \partial_x H(x,\zeta)\right)\right]\right].
\end{aligned}$$

By the convexity of $H(x,\zeta)$, $\partial_x H\left(x,\zeta^k\right)$ is convex and compact, hence $\frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}}\partial_x H\left(x,\zeta^{k,i}\right)$ is convex and the first equality stands. The second equality is due to (7). The third equality is due to (6). The last equality is due to the interchangeability of $\mathbb{E}_\zeta$ and σ, see [47, Proposition 3.4] for details. □

We now derive the subdifferentials convergence of the SAA for h in expectation:

Lemma 3. *For any $\alpha \in (0,1/2)$, we have*

$$\sup_{x\in C}\mathbb{E}_{S_{h,k}}\left[\mathbb{D}\left(\partial\bar{h}_k(x),\partial h(x)\right)\right] \le \frac{\hat{C}_h}{N_{h,k}^{\alpha}},$$

where $\hat{C}_h = 2\sqrt{p}(2L_h + L_h/\sqrt{(1-2\alpha)e})$.

Moreover,

$$\mathbb{P}\left\{N_{h,k}^{\alpha_h}\mathbb{D}\left(\partial\bar{h}_k(x),\partial h(x)\right) \ge \hat{C}_h + t\right\} \le \exp\left\{-t^2/\left(2L_h^2\right)\right\}.$$

Proof. For any $x\in C$, by Lemma 2,

$$\mathbb{E}_{S_{h,k}}\left[\mathbb{D}\left(\partial\bar{h}_k(x),\partial h(x)\right)\right] \le \mathbb{E}_{S_{h,k}}\sup_{\|u\|\leqslant 1}\left\|\frac{1}{N_{k,h}}\sum_{i=1}^{N_{k,h}}\psi(u,\zeta^{k,i}) - \mathbb{E}_\zeta\left[\psi(u,\zeta)\right]\right\|,$$

where $\psi(u,\zeta) = \sigma\left(u,\partial_x h\left(x,\zeta\right)\right)$. To satisfy the condition in Lemma 1, we first verify that $\psi(\cdot,\zeta)$ are uniformly bounded by constant L_h and Lipschitz continuous with constant L_h in the first variable $u\in\mathbb{B}(0,1)\subseteq[-1,1]^d$ uniformly in ζ.

The first property is trivial since $\sup_{s\in\partial_x h(x,\zeta)}\|s\| \le L_h$. Now, we prove the second property. For any $u,v\in\mathbb{B}(0,1)$, suppose that $\sigma\left(u,\partial_x h\left(x,\zeta\right)\right) = u^T s$ where $s\in\partial_x h\left(x,\zeta\right)$. Then we have

$$\sigma\left(v,\partial_x h\left(x,\zeta\right)\right) \ge v^T s \ge u^T s - \|u-v\|\|s\| \ge \sigma\left(u,\partial_x h\left(x,\zeta\right)\right) - L_h\|u-v\|.$$

Similarly, $\sigma\left(u,\partial_x h\left(x,\zeta\right)\right) \ge \sigma\left(v,\partial_x h\left(x,\zeta\right)\right) - L_h\|u-v\|$, hence $\psi(u,\zeta)$ is L_h-Lipschitz continuous in $\mathbb{B}(0,1)$. We thus finish the proof by using Lemma 1. □

We then strengthen the result to bound the square error in expectation as follows. This is the main result of this section.

Theorem 2. *For any $\alpha\in(0,1/2)$, we have*

$$\sup_{x\in C}\mathbb{E}_{S_{h,k}}\left[\mathbb{D}^2\left(\partial\bar{h}_k(x),\partial_x h(x)\right)\right] \le \frac{C_h}{N_{h,k}^{2\alpha}},$$

where $C_h \ge \hat{C}_h\left(\hat{C}_h + \sqrt{2}L_h\frac{\sqrt{\alpha'}}{2\sqrt{(\alpha'-\alpha)e}}\right) + L_h^2$ with $\hat{C}_h = \sqrt{p}(2L_h + L_h/\sqrt{(1-2\alpha')e})$ for any $\alpha'\in(\alpha,1/2)$.

Proof. For any $x\in C$, let $\delta = \mathbb{D}\left(\partial\bar{h}_k(x),\partial_x h(x)\right)$, which is bounded by L_h. By Lemma 3, $\mathbb{E}_{S_{h,k}}[\delta] \le \frac{\hat{C}_h}{N_{h,k}^{\alpha}}$ and $\mathbb{P}\left\{\delta \ge \frac{\hat{C}_h+t}{N_{h,k}^{\alpha_h}}\right\} \le \exp\left\{-t^2/\left(2L_h^2\right)\right\}$ stands for any $t>0$, $\alpha'\in(\alpha,1/2)$.

Hence for any $t>0$, we have

$$\begin{aligned}
\mathbb{E}_{S_{h,k}}[\delta^2] &= \mathbb{E}_{S_{h,k}}\left[\delta^2\mathbf{1}(\delta < \frac{\hat{C}_h+t}{N_{h,k}^{\alpha_h}})\right] + \mathbb{E}_{S_{h,k}}\left[\delta^2\mathbf{1}(\delta \ge \frac{\hat{C}_h+t}{N_{h,k}^{\alpha'}})\right] \\
&\le \mathbb{E}_{S_{h,k}}[\delta]\frac{\hat{C}_h+t}{N_{h,k}^{\alpha'}} + L_h^2\mathbb{P}\left\{\delta \ge \frac{\hat{C}_h+t}{N_{h,k}^{\alpha'}}\right\} \\
&\le \frac{\hat{C}_h(\hat{C}_h+t)}{N_{h,k}^{2\alpha'}} + L_h^2\exp\left\{-t^2/\left(2L_h^2\right)\right\},
\end{aligned}$$

where the first inequality is due to $\delta \le L_h$, and the last inequality is due to Lemma 3.

Take $t = \sqrt{2}L_h\sqrt{\alpha' \ln N_{h,k}}$, we could obtain that

$$\mathbb{E}_{S_{h,k}}[\delta^2] \le \frac{\hat{C}_h\left(\hat{C}_h + \sqrt{2}L_h\sqrt{\alpha' \ln N_{h,k}}\right) + L_h^2}{N_{h,k}^{2\alpha'}}.$$

Since $\sqrt{\ln N_{h,k}} \le \frac{N^{2\alpha'-2\alpha}}{2\sqrt{(\alpha'-\alpha)e}}$, we have for any $\alpha' \in (\alpha, 1/2)$,

$$\mathbb{E}_{S_{h,k}}[\delta^2] \le \frac{\hat{C}_h\left(\hat{C}_h + \sqrt{2}L_h\frac{\sqrt{\alpha'}}{2\sqrt{(\alpha'-\alpha)e}}\right) + L_h^2}{N_{h,k}^{2\alpha}},$$

where $\hat{C}_h = \sqrt{p}(2L_h + L_h/\sqrt{(1-2\alpha')e})$. □

Remark 2. *Notice that the second concentration-type result of Lemma 1 and 3 are due to McDiarmid's bounded difference inequality, and they play an essential role in the proof of Theorem 2.*

Remark 3. *When H is smooth, Theorem 2 could be simplified to*

$$\sup_{x\in C} \mathbb{E}_{S_{h,k}}\left[\|\nabla \bar{h}_k(x) - \nabla h(x)\|^2\right] \le \frac{L_h^2}{N_{h,k}},$$

that is, $C_h = L_h^2$ and $\alpha = 1/2$.

4 Convergence analysis

4.1 Descent and error analysis per iteration

This section analyzes the descent and error at the k-th iteration. We could derive the following sufficiently descent property, similar to the original deterministic version, see, e.g., [64, Theorem 3] and [65, Theorem 3.7].

Lemma 4. *(Sufficiently Descent Property) For any $y_k \in \partial h(x_k)$, the step x_{k+1} from Algorithm 1 satisfies*

$$f(x_k) - f(x_{k+1}) \ge (y_k - \bar{y}_k)^T \bar{d}_k + \mu_{\rho,k}\left\|\bar{d}_k\right\|^2 + \Delta g_k(x_k) - \Delta g_k(x_{k+1}),$$

where $\mu_{\rho,k} = \mu_k + \frac{\rho_g+\rho_h}{2}$ and $\Delta g_k(x) = g(x) - \bar{g}_k(x)$.

Proof. From the convexity of $\bar{g}_k(\cdot)$ and $h(\cdot)$, we have

$$\begin{aligned} &\bar{g}_k(x_k) - \bar{g}_k(x_k + \bar{d}_k) + \bar{z}_{k+1}^T\bar{d}_k \ge \frac{1}{2}\rho_g\|\bar{d}_k\|^2, \\ &h(x_k + \bar{d}_k) - h(x_k) - y_k^T\bar{d}_k \ge \frac{1}{2}\rho_h\|\bar{d}_k\|^2, \end{aligned} \tag{8}$$

for $\bar{z}_{k+1} \in \bar{\partial}\bar{g}_k(x_k + \bar{d}_k)$, $x_k + \bar{d}_k \in C$, $d_k \in \mathbb{R}^n$. Therefore,

$$\begin{aligned} &f(x_k) - f(x_{k+1}) = g(x_k) - h(x_k) - g(x_{k+1}) + h(x_{k+1}) \\ =&\bar{g}_k(x_k) - \bar{g}_k(x_{k+1}) - h(x_k) + h(x_{k+1}) + [g(x_k) - \bar{g}_k(x_k)] - [g(x_{k+1}) - \bar{g}_k(x_{k+1})] \\ \ge& -\bar{z}_{k+1}^T\bar{d}_k + y_k^T\bar{d}_k + \frac{1}{2}(\rho_g + \rho_h)\|\bar{d}_k\|^2 + \Delta g_k(x_k) - \Delta g_k(x_{k+1}). \end{aligned} \tag{9}$$

Taking the dot product with $-\bar{d}_k$ of the first line of (5), we have

$$\begin{aligned} -\bar{z}_{k+1}^T\bar{d}_k + \bar{y}_k^T\bar{d}_k =& \mu_k\|\bar{d}_k\|^2 + \bar{v}_k\bar{d}_k \\ \ge& \mu_k\|\bar{d}_k\|^2 + [i_C(x_k) - i_C(x_k + \bar{d}_k) - \bar{v}_k^T(-\bar{d}_k)] \\ \ge& \mu_k\|\bar{d}_k\|^2, \end{aligned} \tag{10}$$

where the first inequality uses the second line of (5) and the second inequality comes from the convexity of C and $i_C(\cdot)$. Applying (10) to (9) leads to

$$f(x_k) - f(x_{k+1}) \geq (y_k - \bar{y}_k)^T \bar{d}_k + \mu_{\rho,k}\|\bar{d}_k\|^2 + \Delta g_k(x_k) - \Delta g_k(x_{k+1}). \tag{11}$$

□

Remark 4. *In practice, experience from the deterministic problems suggests that μ_k be reduced when possible to improve convergence behavior. For details, see section 3.1 of [70].*

To further the analysis, the estimation error bound in expectation from sampling, which is derived in Section 3, comes into play. In [37], with no specific form of $\bar{\phi}_k(\cdot)$, the variance of the stochastic objectives is assumed to be bounded. The variance bounds then come from sample size N_k. Similarly, in [7], an unbiased estimate with bounded variance is made on the gradient with SQP. In [60], the Monte Carlo estimate of the objective is assumed to be unbiased, and its variance is uniformly bounded.

We present the sufficient descent property in expectation as follows.

Lemma 5. *(Sufficiently Descent Property in Expectation) At the k-th iteration, the following stands for any $c > 0$:*

$$\mathbb{E}\left[f\left(x_k\right) - f\left(x_{k+1}\right)|\mathcal{F}_k\right] \geq (\mu_{\rho,k} - c)\mathbb{E}\left[\left\|\bar{d}_k\right\|^2|\mathcal{F}_k\right] - \frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} - \frac{C_h}{4cN_{h,k}^{\alpha_h}}, \tag{12}$$

where $\alpha_g \in (0, 1/2)$, $\alpha_h \in (0, 1]$ with corresponding C_g, C_h are defined in Theorem 1 and 2.

Proof. By Lemma 4, for any $y_k \in \partial h(x_k)$, $c > 0$,

$$\begin{aligned} f\left(x_k\right) - f\left(x_{k+1}\right) &\geq \left(y_k - \bar{y}_k\right)^T \bar{d}_k + \mu_{\rho,k}\left\|\bar{d}_k\right\|^2 + \Delta g_k\left(x_k\right) - \Delta g_k\left(x_{k+1}\right) \\ &\geq \left(\mu_{\rho,k} - c\right)\left\|\bar{d}_k\right\|^2 + \Delta g_k\left(x_k\right) - \Delta g_k\left(x_{k+1}\right) - \frac{1}{4c}\left\|y_k - \bar{y}_k\right\|^2. \end{aligned}$$

Take y_k such that $dist(\bar{y}_k, \partial h(x_k)) = \|y_k - \bar{y}_k\|$, we have

$$\begin{aligned} f\left(x_k\right) - f\left(x_{k+1}\right) &\geq \left(\mu_{\rho,k} - c\right)\left\|\bar{d}_k\right\|^2 + \Delta g_k\left(x_k\right) - \Delta g_k\left(x_{k+1}\right) - \frac{1}{4c} dist(\bar{y}_k, \partial h(x_k))^2 \\ &\geq \left(\mu_{\rho,k} - c\right)\left\|\bar{d}_k\right\|^2 + \Delta g_k\left(x_k\right) - \Delta g_k\left(x_{k+1}\right) - \frac{1}{4c}\mathbb{D}^2\left(\partial\bar{h}_k(x_k), \partial_x h(x_k)\right) \end{aligned}$$

Taking expectation for both sides under $\mathcal{F}_k$, we finish the proof after Theorem 1 and Theorem 2. □

4.2 Almost sure convergence to critical points

From Lemma 5, to obtain a convergent step $\bar{d}_k \to 0$, the sequences $N_{g,k}$ and $N_{h,k}$ need to be controlled so that $\sum_{k\geq 0}\left(\frac{1}{\mu_k N_{g,k}^{\alpha_g}} + \frac{1}{N_{h,k}^{\alpha_h}}\right) < \infty$. In order to obtain a finite $N_{g,k}$ at each iteration, we make the necessary assumption that μ_k is bounded below by a constant greater than 0.

Assumption 2. *μ_k is bounded below by $\check{\mu} > 0$.*

Under Assumption 2, there exist a number of combinations of conditions to ensure such is the case. For example, this can be achieved as standard practice by increasing the sample size $N_{g,k}$ and $N_{h,k}$ as the following condition:

Condition 1. ***(Summable Condition)*** *Given pre-determined sequences $\{\hat{N}_{g,k}\}$ and $\{\hat{N}_{h,k}\}$ satisfy $\sum_{k\geq 0}\left(\hat{N}_{h,k}^{-\alpha_h} + \hat{N}_{g,k}^{-\alpha_g}\right) < \infty$ and a pre-determined constant $c_l, c_\mu > 0$, we say that* ***Summable Condition*** *is satisfied at the k-th iteration if parameters $c_k, \mu_k, N_{g,k}, N_{h,k}$ are selected to satisfy*

$$N_{g,k} \geq \hat{N}_{g,k},\ N_{h,k} \geq \hat{N}_{h,k},\ \text{and}\ c_l \leq c_k \leq \mu_{\rho,k} - c_\mu. \tag{13}$$

We could ensure the convergence of osproxDCA by directly setting a sublinear growth for the sample size. However, this approach is not adaptive. We propose a practical **Stepsize Norm Condition** for adaptive sampling in the following part. Compared to the gradient accuracy condition and other variance-based tests in [**?**, **?**], our method bridges the gap between theory and practical implementation.

Condition 2. ***(Stepsize Norm Condition)*** *Given a pre-determined constant $c_\mu > 0$, we say that* ***Stepsize Norm Condition*** *is satisfied at the k-th iteration if parameters $c_k, \mu_k, N_{g,k}, N_{h,k}$ are selected to satisfy*

$$(\mu_{\rho,k-1} - c_\mu - c_{k-1}) \|\bar{d}_{k-1}\|^2 \geq \frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} + \frac{C_h}{4c_k N_{h,k}^{\alpha_h}}, \text{ and } c_k \leq \mu_{\rho,k} - c_\mu. \tag{14}$$

Remark 5. *Here, $\{c_k\}$ are intermediate variables for parameter updates. They provide the interrelationship between μ_k, $N_{g,k}$, and $N_{h,k}$ to ensure the convergence of osproxDCA in an expectation. Note that it is only used to determine μ_k, $N_{g,k}$, and $N_{h,k}$, and is irrelevant to the solution of the subproblems.*

We present the convergence result in the following theorems, which are significant for designing practical adaptive osproxDCA.

Theorem 3. *(Square Summable Property) Under Assumption 2, if either* ***Summable Condition*** (13) *or* ***Stepsize Norm Condition*** (14) *is satisfied for sufficiently large k, then we have*

$$\lim_{k\to\infty} \mathbb{E}\left[\sum_{k\geq 0} \left\|\bar{d}_k\right\|^2 |\mathcal{F}_0\right] < \infty \text{ , hence } \mathbb{E}\left[\left\|\bar{d}_k\right\| |\mathcal{F}_0\right] \to 0 \text{ as } k \to \infty.$$

Further, $\lim_{k\to\infty}\|\bar{d}_k\| = 0$ with probability 1.

Proof. Let $\mathcal{K}_1$ be the set of k that satisfies Summable Condition, $\mathcal{K}_2$ be the set of k that satisfies Stepsize Norm Condition. Supposed that when $k \geq K$, $k \in \mathcal{K}_1 \cup \mathcal{K}_2$. If $k \in \mathcal{K}_2$, we have

$$\mathbb{E}\left[f(x_k) - f(x_{k+1}) |\mathcal{F}_k\right] \geq (\mu_{\rho,k} - c_k)\mathbb{E}\left[\left\|\bar{d}_k\right\|^2 |\mathcal{F}_k\right] - (\mu_{\rho,k-1} - c_\mu - c_k) \|\bar{d}_{k-1}\|^2.$$

Taking expectation under $\mathcal{F}_0$ from both side, we have

$$\mathbb{E}\left[f(x_k) - f(x_{k+1}) |\mathcal{F}_0\right] \geq (\mu_{\rho,k} - c_k)\mathbb{E}\left[\left\|\bar{d}_k\right\|^2 |\mathcal{F}_0\right] - (\mu_{\rho,k-1} - c_\mu - c_k)\, \mathbb{E}\left[\|\bar{d}_{k-1}\|^2 |\mathcal{F}_0\right].$$

If $k \in \mathcal{K}_1 \cup \{0, 1, \cdots, K-1\}$, from Lemma 5 and take expectation under $\mathcal{F}_0$, we have

$$\mathbb{E}\left[f(x_k) - f(x_{k+1}) |\mathcal{F}_0\right] \geq (\mu_{\rho,k} - c_k)\mathbb{E}\left[\left\|\bar{d}_k\right\|^2 |\mathcal{F}_0\right] - \frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} - \frac{C_h}{4c_k N_{h,k}^{\alpha_h}}.$$

Taking sum for each k, we have

$$\begin{aligned}\mathbb{E}\left[\sum_{k=0}^{n-1} (\mu_{\rho,k} - c_k) \|\bar{d}_k\|^2 |\mathcal{F}_0\right] &\leq \mathbb{E}\left[f(x_0) - f(x_n) |\mathcal{F}_0\right] + \sum_{k=0}^{K-1}\left(\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} + \frac{C_h}{4c_k N_{h,k}^{\alpha_h}}\right) \\ &+ \sum_{k=K,k\in\mathcal{K}_1}^{n-1}\left(\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} + \frac{C_h}{4c_k N_{h,k}^{\alpha_h}}\right) + \mathbb{E}\left[\sum_{k=K,k\in\mathcal{K}_2}^{n-1} (\mu_{\rho,k-1} - c_{k-1} - c_\mu) \|\bar{d}_{k-1}\|^2 |\mathcal{F}_0\right].\end{aligned} \tag{15}$$

Hence, we could obtain that

$$\begin{aligned}\mathbb{E}\left[\sum_{k=K-1}^{n-1} c_\mu \|\bar{d}_k\|^2 |\mathcal{F}_0\right] &\leq \mathbb{E}\left[f(x_0) - f(x_n) |\mathcal{F}_0\right] \\ &+ \sum_{k=0}^{K-1}\left(\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} + \frac{C_h}{4c_k N_{h,k}^{\alpha_h}}\right) + \sum_{k=K,k\in\mathcal{K}_1}^{n-1}\left(\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}} + \frac{C_h}{4c_l N_{h,k}^{\alpha_h}}\right).\end{aligned} \tag{16}$$

By Assumption 1, $\mathbb{E}\left[f\left(x_0\right)-f\left(x_n\right)|\mathcal{F}_0\right]<f(x_0)-f_m$. By the definition of Summable Condition and Assumption 2, we have

$$\sum_{k=K,k\in\mathcal{K}_1}^{n-1}\left(\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}}+\frac{C_h}{4c_l N_{h,k}^{\alpha_h}}\right)\leq\sum_{k=K,k\in\mathcal{K}_1}^{n-1}\left(\frac{C_g}{\tilde{\mu}\hat{N}_{g,k}^{\alpha_g}}+\frac{C_h}{4c_l \hat{N}_{h,k}^{\alpha_h}}\right).$$

Take $n\to\infty$, the right-hand side of (16) is bounded. It follows that $\lim_{n\to\infty}\mathbb{E}[\|\bar{d}_n\|^2|\mathcal{F}_0]=0$. The first part of the theorem is proven.

We proceed by contradiction for the second part of the theorem. Suppose there exists $\epsilon>0$ and $a>0$ such that

$$\mathbb{P}(\limsup_{k\to\infty}\|\bar{d}_k\|\geq\epsilon)\geq a. \tag{17}$$

By Chebyshev's inequality, we have $\mathbb{P}(\|\bar{d}_k\|\geq\epsilon)\leq\frac{\mathbb{E}\left[\|\bar{d}_k\|^2|\mathcal{F}_0\right]}{\epsilon^2}$. Since $\mathbb{E}\left[\|\bar{d}_k\|^2|\mathcal{F}_0\right]$ is finitely summable, there exists $N>0$ such that $\sum_{k=N}^{\infty}\mathbb{P}(\|\bar{d}_k\|\geq\epsilon)\leq\sum_{k=N}^{\infty}\frac{\mathbb{E}\left[\|\bar{d}_k\|^2|\mathcal{F}_0\right]}{\epsilon^2}<a$. Therefore,

$$\mathbb{P}\left(\limsup_{k\to\infty}\|\bar{d}_k\|\geq\epsilon\right)=\mathbb{P}\left(\limsup_{k\to\infty:k\geq N}\|\bar{d}_k\|\geq\epsilon\right)\leq\sum_{k=N}^{\infty}\mathbb{P}(\|\bar{d}_k\|\geq\epsilon)<a. \tag{18}$$

This is a contradiction against (17). Hence, we could obtain that $\lim_{k\to\infty}\|\bar{d}_k\|=0$ with probability 1. □

In the next part, we prove the convergence to critical points. The following assumption is natural and necessary; without it, the algorithm may not converge to critical points, even in the deterministic case.

Assumption 3. *μ_k is bounded above by $\hat{\mu}>0$.*

To proceed, we first prove a technical lemma.

Lemma 6. *Under Assumption 3, if either* ***Summable Condition*** (13) *or* ***Stepsize Norm Condition*** (14) *is satisfied for sufficiently large k, then $N_{g,k},N_{h,k}\to\infty$ as $k\to\infty$ with probability 1.*

Proof. Supposed that when $k\geq K$, $k\in\mathcal{K}_1\cup\mathcal{K}_2$. If $k\in\mathcal{K}_1$, we have $N_{g,k}\geq\hat{N}_{g,k}$ and $N_{h,k}\geq\hat{N}_{h,k}$. Since $\sum_{k\geq0}\left(N_{h,k}^{-\alpha_h}+N_{g,k}^{-\alpha_g}\right)<\infty$, we have $N_{g,k},N_{h,k}\to\infty$ as $k\to\infty$.

If $k\in\mathcal{K}_2$, we have

$$\begin{aligned}\frac{C_g}{\hat{\mu}N_{g,k}^{\alpha_g}}+\frac{C_h}{4(\hat{\mu}+\frac{\rho_g+\rho_h}{2}-c_\mu)N_{h,k}^{\alpha_h}}&\leq\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}}+\frac{C_h}{4(\mu_k+\frac{\rho_g+\rho_h}{2}-c_\mu)N_{h,k}^{\alpha_h}}\\&\leq\frac{C_g}{\mu_k N_{g,k}^{\alpha_g}}+\frac{C_h}{4c_k N_{h,k}^{\alpha_h}}\\&\leq\left(\mu_{\rho,k-1}-c_\mu-c_{k-1}\right)\|\bar{d}_{k-1}\|^2\\&\leq\left(\hat{\mu}+\frac{\rho_g+\rho_h}{2}-c_\mu-c_{k-1}\right)\|\bar{d}_{k-1}\|^2,\end{aligned}$$

where the first and last inequality is due to Assumption 3. The second and third inequality is due to the definition of Stepsize Norm Condition (14).

By Theorem 3, $\lim_{k\to\infty}\|\bar{d}_k\|=0$ with probability 1, thus $N_{g,k},N_{h,k}\to\infty$ with probability 1. □

We then introduce a result concerning the law of large numbers (LLN) for subdifferentials, which follows from the LLN for random set-valued mappings as discussed in [1]. This result can be interpreted as a pointwise version of [59, Theorem 2].

Lemma 7. *For any $r > r' \geqslant 0$ the following limits hold:*

$$\sup_{x\in\mathcal{C}} \mathbb{D}\left(\partial^{r'}\bar{g}_k(x), \partial^r g(x)\right) \to 0 \ \ w.p.\ 1.$$

$$\sup_{x\in\mathcal{C}} \mathbb{D}\left(\partial^{r'}\bar{h}_k(x), \partial^r h(x)\right) \to 0 \ \ w.p.\ 1.$$

Now, we derive our main result as follows.

Theorem 4. *Under Assumption 2 and 3, if either* ***Summable Condition*** (13) *or* ***Stepsize Norm Condition*** (14) *is satisfied for sufficiently large k, every accumulation point of the sequence $\{x_k\}$ produced by Algorithm 1 is a DC critical point of f with probability 1.*

Proof. Let x be an accumulation point of $\{x_k\}$. Firstly we prove that for any given ε, $0 \in \partial^\varepsilon g(x) - \partial^\varepsilon h(x) + \partial i_C(x)$.

From the optimal condition (5), there exist $\bar{z}_{k+1} \in \partial\bar{g}_k(x_{k+1})$ and $\bar{v}_k \in \partial i_C(x_{k+1})$ for each k, such that

$$\bar{z}_{k+1} - \bar{y}_k + \mu_k\bar{d}_k + \bar{v}_k = 0. \tag{19}$$

By Lemma 7, there exists $y_k \in \partial^\varepsilon h(x_k)$, $z_k \in \partial^\varepsilon g(x_{k+1})$ such that $\lim_{k\to\infty} \bar{y}_k - y_k = 0$ and $\lim_{k\to\infty} \bar{z}_k - z_k = 0$ with probability 1. We can assume $\lim_{k\to\infty} x_{n_k} = x$ where $x \in C$. Further, y_k and z_k are bounded due to the Lipschitz continuity of $G(x,\xi)$ and $H(x,\zeta)$. Thus, $\bar{v}_k$ is also bounded, and there exist accumulation points for $\{y_k\}$ and $\{z_k\}$. Passing on to a subsequence, we assume $y_{n_{m_k}} \to \bar{y}$ and $z_{n_{m_k}} \to \bar{z}$. Therefore $\bar{y}_{n_{m_k}} \to \bar{y}$ and $\bar{z}_{n_{m_k}} \to \bar{z}$. By (19), we have

$$0 = \bar{z}_{n_{m_k}} - \bar{y}_{n_{m_k}} + \mu_{n_{m_k}}\bar{d}_{n_{m_k}} + \bar{v}_{n_{m_k}}. \tag{20}$$

By Theorem 3 and Assumption 3, $\lim_{k\to\infty} \mu_k\bar{d}_k = 0$ with probability 1. Thus, $\lim_{k\to\infty} \bar{v}_{n_{m_k}} = \bar{y} - \bar{z}$ with probability 1. Given that $\bar{v}_{n_{m_k}} \in \partial i_C(x_{n_{m_k}+1})$, the outer semicontinuity of $\partial i_C(\cdot)$ leads to $\lim_{k\to\infty} \bar{v}_{n_{m_k}} \in \partial i_C(x)$. By the outer semicontinuity of Goldstein subdifferential, $\bar{z} = \lim_{k\to\infty} z_{n_{m_k}} \in \partial^\varepsilon g(x)$, since $x_{n_{m_k}+1} \to x$ and $z_{n_{m_k}} \in \partial^\varepsilon g(x_{n_{m_k}+1})$. Similarly, $\bar{y} = \lim_{k\to\infty} y_{n_{m_k}} \in \partial h^\varepsilon(x)$.

Therefore, with probability 1, $0 \in \partial^\varepsilon g(x) - \partial^\varepsilon h(x) + \partial i_C(x)$. By the outer semicontinuity of Clarke subdifferential, we obtain that

$$\bigcap_{\epsilon>0} (\partial^\epsilon g(x) - \partial^\epsilon h(x) + \partial i_C(x)) = \partial g(x) - \partial h(x) + \partial i_C(x).$$

Thus we have $0 \in \partial g(x) - \partial h(x) + \partial i_C(x)$ with probability 1. □

Remark 6. *Indeed, the current solution can be considered as the initial point for restart. Hence, even if the distribution of ξ and ζ changes at a certain time, the algorithm continues operating based on new samples from the new distribution. Theorem 3 and 4 are still valid, and the subsequential convergence with probability one to DC critical points of the DC problem associated with the new distribution is guaranteed. This is indeed an advantage of the osproxDCA. In contrast, intuitively, stochastic algorithms using aggregated updates (still using old samples to compute the current solution) barely have this kind of adaptivity.*

5 Adaptive sampling osproxDCA

The previous theorems are enough to get a simple adaptive scheme for osproxDCA, following a similar strategy of [8, 11, 5, 69]. We present the adaptive sampling osproxDCA as Algorithm 2.

When $\rho_g + \rho_h > 0$, given any predetermined sequence $\{\mu_k\}$ that satisfy $0 < \check{\mu} \leq \mu_k \leq \hat{\mu}$, we present a simplified version for Algorithm 2 as follows by fixing $c_k = \frac{\rho_g+\rho_h}{2} = c_l$, $c_\mu = \frac{\check{\mu}}{2}$, see Algorithm 3.

There has now arisen a natural question:

What is the optimal strategy for updating the sample sizes $N_{g,k}$ and $N_{h,k}$?

The following lemma provides the optimal order between $N_{g,k}$ and $N_{h,k}$ to minimize the total number of samples at each iteration.

Algorithm 2 Adaptive osproxDCA

Require: Initial point x_0, initial parameter μ_0, c_0, $N_{g,0}$ and $N_{h,0}$, error estimation parameter $\alpha_g \in (0, 1/2)$, $\alpha_h \in (0, 1]$ with corresponding C_g, C_h defined in Theorem 1 and 2, step size norm condition parameter $c_\mu, c_l > 0$, sample size upper bound sequence $\{\hat{N}_{g,k}\}$ and $\{\hat{N}_{h,k}\}$ which satisfy $\sum_{k\geq 0}\left(\hat{N}_{h,k}^{-\alpha_h} + \hat{N}_{g,k}^{-\alpha_g}\right) < \infty$, step size upper bound $\hat{\mu}$ and lower bound $\check{\mu}$ that satisfy $\hat{\mu} > \check{\mu}$.

1: **for** $k = 0, 1, 2, \cdots$ **do**

2: Generate iid samples $\{\xi^{k,i}\}_{i=1}^{N_{g,k}}$ and $\{\zeta^{k,i}\}_{i=1}^{N_{h,k}}$ from the distribution of ξ and ζ that are independent of previous samples.

3: Construct the local approximation model $\bar{M}_k(d)$:
Set $\bar{g}_k(x) = \frac{1}{N_{g,k}}\sum_{i=1}^{N_{g,k}} G\left(x, \xi^{k,i}\right)$, $\bar{h}_k(x) = \frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} H\left(x, \zeta^{k,i}\right)$, and select $\bar{y}_k \in \partial\bar{h}_k(x) = \frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \partial_x H\left(x, \zeta^{k,i}\right)$.

4: Solve the $(\rho_g + \mu_k)$-convex optimization subproblem (4) to obtain $\bar{d}_k$:

$$\begin{aligned} \underset{d}{\text{minimize}} \quad & \bar{g}_k(x_k + d) - \bar{h}(x_k) - \bar{y}_k^T d + \frac{1}{2}\mu_k\|d\|^2, \\ \text{subject to} \quad & x_k + d \in C. \end{aligned}$$

5: Take the step $x_{k+1} = x_k + \bar{d}_k$.

6: Update $\check{\mu} \leq \mu_{k+1} \leq \hat{\mu}$, $c_{k+1}, N_{g,k+1}$ and $N_{h,k+1}$ such that one of the followings stands:
1. $\left(\mu_k + \frac{\rho_g+\rho_h}{2} - c_\mu - c_k\right)\|\bar{d}_k\|^2 \geq \frac{C_g}{\mu_{k+1}N_{g,k+1}^{\alpha_g}} + \frac{C_h}{4c_k N_{h,k+1}^{\alpha_h}}$, and $c_{k+1} \leq \mu_{k+1} + \frac{\rho_g+\rho_h}{2} - c_\mu$,
2. $N_{g,k+1} \geq \hat{N}_{g,k+1}$, $N_{h,k} \geq \hat{N}_{h,k+1}$, and $c_l \leq c_{k+1} \leq \mu_{k+1} + \frac{\rho_g+\rho_h}{2} - c_\mu$.

7: **end for**

Algorithm 3 Simplified Adaptive osproxDCA

Require: Initial point x_0, error estimation parameter $\alpha_g \in (0, 1/2)$, $\alpha_h \in (0, 1]$ with corresponding C_g, C_h defined in Theorem 1 and 2, sample size upper bound sequence $\{\hat{N}_{g,k}\}$ and $\{\hat{N}_{h,k}\}$ which satisfy $\sum_{k\geq 0}\left(\hat{N}_{h,k}^{-\alpha_h} + \hat{N}_{g,k}^{-\alpha_g}\right) < \infty$, predetermined quadratic term sequence $\{\mu_k\}$ with upper bound $\hat{\mu}$ and lower bound $\check{\mu}$.

1: **for** $k = 0, 1, 2, \cdots$ **do**

2: Generate iid samples $\{\xi^{k,i}\}_{i=1}^{N_{g,k}}$ and $\{\zeta^{k,i}\}_{i=1}^{N_{h,k}}$ from the distribution of ξ and ζ that are independent of previous samples.

3: Construct the local approximation model $\bar{M}_k(d)$:
Set $\bar{g}_k(x) = \frac{1}{N_{g,k}}\sum_{i=1}^{N_{g,k}} G\left(x, \xi^{k,i}\right)$, $\bar{h}_k(x) = \frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} H\left(x, \zeta^{k,i}\right)$, and select $\bar{y}_k \in \partial\bar{h}_k(x) = \frac{1}{N_{h,k}}\sum_{i=1}^{N_{h,k}} \partial_x H\left(x, \zeta^{k,i}\right)$.

4: Solve the $\rho_g + \mu_k$-convex optimization subproblem (4) to obtain $\bar{d}_k$:

$$\begin{aligned} \underset{d}{\text{minimize}} \quad & \bar{g}_k(x_k + d) - \bar{h}(x_k) - \bar{y}_k^T d + \frac{1}{2}\mu_k\|d\|^2, \\ \text{subject to} \quad & x_k + d \in C. \end{aligned}$$

5: Take the step $x_{k+1} = x_k + \bar{d}_k$.

6: Update $N_{g,k+1}$ and $N_{h,k+1}$ such that one of the followings stands:
1. $\left(\mu_k - \frac{\check{\mu}}{2}\right)\|\bar{d}_k\|^2 \geq \frac{C_g}{\mu_{k+1}N_{g,k+1}^{\alpha_g}} + \frac{C_h}{2(\rho_g+\rho_h)N_{h,k+1}^{\alpha_h}}$,
2. $N_{g,k+1} \geq \hat{N}_{g,k+1}$, and $N_{h,k} \geq \hat{N}_{h,k+1}$.

7: **end for**

Lemma 8. *(Optimal Order of N_g and N_h) Consider the following subproblem when updating the sample size:*

$$\begin{aligned} &\textit{minimize} \quad N_{g,k} + N_{h,k}, \\ &\textit{subject to} \quad \frac{C_1}{N_{g,k}^{1/2}} + \frac{C_2}{N_{h,k}} \leq C_0, \\ &N_{g,k} \geq 0, \quad N_{h,k} \geq 0. \end{aligned}$$

We could derive that $N_{h,k} = \sqrt{2}\left(\frac{C_2}{C_1}\right)^{1/2} N_{g,k}^{3/4}$. Hence the optimal order of $N_{h,k}$ is $O(s_{g,k}^{3/4})$.

Remark 7. *If the updating rule of $N_{g,k}$ and $N_{h,k}$ is based on Lemma 8, then sample size upper bound sequence $\hat{N}_{h,k}$ is no longer required.*

6 Numerical applications

6.1 Objective with smooth concave part

Consider the expected problem of principal component analysis (denoted by (E-PCA)) [42],

$$\min\left\{f(x) = -(1/2)\mathbb{E}_{s\sim\mathbb{P}}\left(\langle x, s\rangle^2\right) : x \in \mathbb{R}^p \|x\| \leq 1\right\},$$

where s is a normalized random vector, i.e., $\|s\| = 1$. We formulate the (E-PCA) problem as an SDC program with $G(x,s) = \frac{L}{2}\|x\|^2 - \frac{1}{2}\langle x, s\rangle^2, H(x,s) = \frac{L}{2}\|x\|^2 + \frac{1}{2}\langle x, s\rangle^2$, feasible region $C = \mathbb{B}(0,1)$, where $L \geq 1$. Then for all s, $G(\cdot, s)$ is Lipschitz with modulus L and convex with modulus $L-1$ on C. $H(\cdot, s)$ is Lipschitz with modulus $L+1$ and convex with modulus L on C. In Algorithm 3, we take constant quadratic term $\mu_k = \mu$, $\alpha_g = 0.45$ with corresponding $C_g = \sqrt{p}8L(L+1)(2L+1)$. Since $H(\cdot, s)$ is smooth, we take $\alpha_h = 1$ with corresponding $C_h = (L+1)^2$.

6.2 Objective with nonsmooth concave part

Acknowledgments

Yuhan Ye is partially supported by the elite undergraduate training program of School of Mathematical Sciences at Peking University.

References

[1] Z. Artstein and R. A. Vitale, *A strong law of large numbers for random compact sets*, The Annals of Probability, (1975), pp. 879–882.

[2] H. Attouch, *Convergence de fonctions convexes, des sous-différentiels et semi-groupes associés*, CR Acad. Sci. Paris, 284 (1977), p. 13.

[3] R. J. Aumann, *Integrals of set-valued functions*, Journal of Mathematical Analysis and Applications, 12 (1965), pp. 1–12.

[4] F. Beiser, B. Keith, S. Urbainczyk, and B. Wohlmuth, *Adaptive sampling strategies for risk-averse stochastic optimization with constraints*, arXiv preprint arXiv:2012.03844, (2020).

[5] A. S. Berahas, R. Bollapragada, and B. Zhou, *An adaptive sampling sequential quadratic programming method for equality constrained stochastic optimization*, 2023, `https://arxiv.org/abs/2206.00712`.

[6] A. S. Berahas, L. Cao, and K. Scheinberg, *Global convergence rate analysis of a generic line search algorithm with noise*, SIAM Journal on Optimization, 31 (2021), pp. 1489–1518.

[7] A. S. Berahas, F. E. Curtis, D. Robinson, and B. Zhou, *Sequential quadratic optimization for nonlinear equality constrained stochastic optimization*, SIAM J. Optim., 31 (2021), pp. 1352–79.

[8] R. Bollapragada, R. Byrd, and J. Nocedal, *Adaptive sampling strategies for stochastic optimization*, SIAM Journal on Optimization, 28 (2018), pp. 3312–3343.

[9] R. Bollapragada, J. Nocedal, D. Mudigere, H.-J. Shi, and P. T. P. Tang, *A progressive batching l-bfgs method for machine learning*, in International Conference on Machine Learning, PMLR, 2018, pp. 620–629.

[10] O. Bousquet and A. Elisseeff, *Stability and generalization*, J. Mach. Learn. Res., 2 (2002), pp. 499–526.

[11] R. H. Byrd, G. M. Chin, J. Nocedal, and Y. Wu, *Sample size selection in optimization methods for machine learning*, Mathematical programming, 134 (2012), pp. 127–155.

[12] R. G. Carter, *On the global convergence of trust region algorithms using inexact gradient information*, SIAM Journal on Numerical Analysis, 28 (1991), pp. 251–265.

[13] C. Cartis and K. Scheinberg, *Global convergence rate analysis of unconstrained optimization methods based on probabilistic models*, Mathematical Programming, 169 (2018), pp. 337–375.

[14] C. Castaing, M. Valadier, et al., *Convex analysis and measurable multifunctions [electronic resource]*.

[15] H. Christian, *Chapter 14 - set-valued integration and set-valued probability theory: An overview*, in Handbook of Measure Theory, E. PAP, ed., North-Holland, Amsterdam, 2002, pp. 617–673.

[16] F. H. Clarke, *Optimization and Nonsmooth Analysis*, SIAM, 1990.

[17] Y. Cui and J.-S. Pang, *Modern Nonconvex Nondifferentiable Optimization*, SIAM, 2021.

[18] F. E. Curtis and K. Scheinberg, *Adaptive stochastic optimization: A framework for analyzing stochastic optimization algorithms*, IEEE Signal Processing Magazine, 37 (2020), pp. 32–42.

[19] D. Davis and D. Drusvyatskiy, *Stochastic model-based minimization of weakly convex functions*, 2018, `https://arxiv.org/abs/1803.06523`.

[20] D. Davis and D. Drusvyatskiy, *Subgradient methods under weak convexity and tame geometry*, SIAG/OPT Views and News, 28 (2020), pp. 1–10.

[21] D. Davis and D. Drusvyatskiy, *Graphical convergence of subgradients in nonconvex optimization and learning*, Mathematics of Operations Research, 47 (2022), pp. 209–231.

[22] Y. M. Ermoliev and V. I. Norkin, *Sample average approximation method for compound stochastic optimization problems*, SIAM Journal on Optimization, 23 (2013), pp. 2231–2263.

[23] D. Foster, A. Sekhari, and K. Sridharan, *Uniform convergence of gradients for non-convex learning and optimization*, in Advances in Neural Information Processing Systems, 2018. Accepted, arXiv:1810.11059.

[24] M. P. Friedlander and M. Schmidt, *Hybrid deterministic-stochastic methods for data fitting*, SIAM Journal on Scientific Computing, 34 (2012), pp. A1380–A1405.

[25] C. J. Geyer, *On the asymptotics of constrained M-estimation*, Annals of Statistics, 22 (1994), pp. 1993–2010.

[26] A. A. Goldstein, *Optimization of lipschitz continuous functions*, Mathematical Programming, 13 (1977), pp. 14–22.

[27] P. D. Grünwald and N. A. Mehta, *Fast rates for general unbounded loss functions: From erm to generalized bayes*, arXiv preprint arXiv:1605.00252, (2016).

[28] F. S. Hashemi, S. Ghosh, and R. Pasupathy, *On adaptive sampling rules for stochastic recursions*, in Proceedings of the Winter Simulation Conference 2014, IEEE, 2014, pp. 3959–3970.

[29] B. Jin, K. Scheinberg, and M. Xie, *High probability complexity bounds for line search based on stochastic oracles*, arXiv preprint arXiv:2106.06454, (2021).

[30] S. M. Kakade, K. Sridharan, and A. Tewari, *On the complexity of linear prediction: Risk bounds, margin bounds, and regularization*, in Advances in Neural Information Processing Systems, 2009, pp. 793–800.

[31] Y. M. Kaniovski, A. J. King, and R. J.-B. Wets, *Probabilistic bounds (via large deviations) for the solutions of stochastic programming problems*, Annals of Operations Research, 56 (1995), pp. 189–208.

[32] A. J. King and R. T. Rockafellar, *Asymptotic theory for solutions in statistical estimation and stochastic programming*, Mathematics of Operations Research, 18 (1993), pp. 148–162.

[33] H. A. Le Thi, V. N. Huynh, T. P. Dinh, and H. P. Hau Luu, *Stochastic difference-of-convex-functions algorithms for nonconvex programming*, SIAM Journal on Optimization, 32 (2022), pp. 2263–2293, https://doi.org/10.1137/20M1385706, https://doi.org/10.1137/20M1385706, https://arxiv.org/abs/https://doi.org/10.1137/20M1385706.

[34] H. A. Le Thi, H. M. Le, D. N. Phan, and B. Tran, *Stochastic dca for minimizing a large sum of dc functions with application to multi-class logistic regression*, Neural Networks, 132 (2020), pp. 220–231, https://doi.org/https://doi.org/10.1016/j.neunet.2020.08.024, https://www.sciencedirect.com/science/article/pii/S0893608020303233.

[35] H. A. Le Thi, H. P. H. Luu, and T. P. Dinh, *Online stochastic dca with applications to principal component analysis*, IEEE Transactions on Neural Networks and Learning Systems, 35 (2024), pp. 7035–7047, https://doi.org/10.1109/TNNLS.2022.3213558.

[36] H. A. Le Thi and T. Pham Dinh, *Dc programming and dca: thirty years of developments*, Mathematical Programming, 169 (2018), https://doi.org/10.1007/s10107-018-1235-y.

[37] J. Liu, Y. Cui, and J.-S. Pang, *Solving nonsmooth and nonconvex compound stochastic programs with applications to risk measure minimization*, Mathematics of Operations Research, 47 (2022).

[38] M. Liu, X. Zhang, L. Zhang, R. Jin, and T. Yang, *Fast rates of erm and stochastic approximation: Adaptive to error bound conditions*, arXiv preprint arXiv:1805.04577, (2018).

[39] N. A. Mehta, *Fast rates with high probability in exp-concave statistical learning*, arXiv preprint arXiv:1605.01288, (2016).

[40] N. A. Mehta and R. C. Williamson, *From stochastic mixability to fast rates*, in Proceedings of the 27th International Conference on Neural Information Processing Systems - Volume 1, NIPS'14, Cambridge, MA, USA, 2014, MIT Press, pp. 1197–1205.

[41] S. Mei, Y. Bai, and A. Montanari, *The landscape of empirical risk for nonconvex losses*, Annals of Statistics, 46 (2018), pp. 2747–2774.

[42] A. Montanari and E. Richard, *Non-negative principal component analysis: Message passing algorithms and sharp asymptotics*, IEEE Transactions on Information Theory, 62 (2016), pp. 1458–1484.

[43] B. S. Mordukhovich, *Variational Analysis and Generalized Differentiation I: Basic Theory*, vol. 330, Springer Science & Business Media, 2006.

[44] J.-J. Moreau, *Proximité et dualité dans un espace hilbertien*, Bulletin de la Société mathématique de France, 93 (1965), pp. 273–299.

[45] A. Moudafi, *A Regularization of DC Optimization.*, Pure and Applied Functional Analysis, (2022), https://amu.hal.science/hal-03581239.

[46] A. Nitanda and T. Suzuki, *Stochastic Difference of Convex Algorithm and its Application to Training Deep Boltzmann Machines*, in Proceedings of the 20th International Conference on Artificial Intelligence and Statistics, A. Singh and J. Zhu, eds., vol. 54 of Proceedings of Machine Learning Research, PMLR, 20–22 Apr 2017, pp. 470–478, https://proceedings.mlr.press/v54/nitanda17a.html.

[47] N. S. Papageorgiou, *On the theory of banach space valued multifunctions. 1. integration and conditional expectation*, Journal of Multivariate Analysis, 17 (1985), pp. 185–206.

[48] R. Pasupathy, P. Glynn, S. Ghosh, and F. S. Hashemi, *On sampling rates in simulation-based recursions*, SIAM Journal on Optimization, 28 (2018), pp. 45–73.

[49] S. T. Rachev and W. Römisch, *Quantitative stability in stochastic programming: The method of probability metrics*, Mathematics of Operations Research, 27 (2002), pp. 792–818.

[50] A. Rakhlin, S. Mukherjee, and T. Poggio, *Stability results in learning theory*, Analysis and Applications, 03 (2005), pp. 397–417, https://doi.org/10.1142/S0219530505000650.

[51] S. M. Robinson, *Analysis of sample-path optimization*, Mathematics of Operations Research, 21 (1996), pp. 513–528.

[52] R. T. Rockafellar and R. J.-B. Wets, *Variational Analysis*, Springer-Verlag, Berlin Heidelberg, 1998.

[53] W. Römisch and R. Wets, *Stability of ε-approximate solutions to convex stochastic programs*, SIAM Journal on Optimization, 18 (2007), pp. 961–979.

[54] F. Ruan, *Subgradient convergence implies subdifferential convergence on weakly convex functions: With uniform rates guarantees*, 2024, https://arxiv.org/abs/2405.10289.

[55] S. Shalev-Shwartz, O. Shamir, N. Srebro, and K. Sridharan, *Stochastic convex optimization*, in Proceedings of the Conference on Learning Theory (COLT), 2009.

[56] A. Shapiro, *On the asymptotics of constrained local M-estimators*, Annals of Statistics, 28 (2000), pp. 948–960.

[57] A. Shapiro, *Stochastic Programming by Monte Carlo Simulation Methods*, Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät II, Institut für Mathematik, 2000.

[58] A. Shapiro and T. H. de Mello, *On the rate of convergence of optimal solutions of monte carlo approximations of stochastic programs*, SIAM Journal on Optimization, 11 (2000), pp. 70–86.

[59] A. Shapiro and H. Xu, *Uniform laws of large numbers for set-valued mappings and subdifferentials of random functions*, Journal of Mathematical Analysis and Applications, 325 (2007), pp. 1390–1399.

[60] S. Shashaani, F. S. Hashemi, and R. Pasupathy, *ASTRO-DF: A class of adaptive sampling trust-region algorithms for derivative-free stochastic optimization*, SIAM Journal on Optimization, 28 (2018), pp. 3145–3176.

[61] N. Srebro, K. Sridharan, and A. Tewari, *Smoothness, low noise and fast rates*, in Advances in Neural Information Processing Systems 23, J. D. Lafferty, C. K. I. Williams, J. Shawe-Taylor, R. S. Zemel, and A. Culotta, eds., Curran Associates, Inc., 2010, pp. 2199–2207.

[62] K. Sridharan, S. Shalev-shwartz, and N. Srebro, *Fast rates for regularized objectives*, in Advances in Neural Information Processing Systems 21, D. Koller, D. Schuurmans, Y. Bengio, and L. Bottou, eds., Curran Associates, Inc., 2009, pp. 1545–1552.

[63] K. Sun and X. A. Sun, *Algorithms for difference-of-convex programs based on difference-of-moreau-envelopes smoothing*, INFORMS J. Optim., 5 (2022), pp. 321–339, https://api.semanticscholar.org/CorpusID:233025038.

[64] P. D. Tao and L. H. An, *Convex analysis approach to dc programming: theory, algorithms and applications*, Acta mathematica vietnamica, 22 (1997), pp. 289–355.

[65] P. D. Tao and L. T. H. An, *A d.c. optimization algorithm for solving the trust-region subproblem*, SIAM Journal on Optimization, 8 (1998), pp. 476–505.

[66] P. D. Tao and E. B. Souad, *Algorithms for solving a class of nonconvex optimization problems. methods of subgradients*, North-holland Mathematics Studies, 129 (1986), pp. 249–271, https://api.semanticscholar.org/CorpusID:117927490.

[67] H. A. L. Thi, H. M. Le, D. N. Phan, and B. Tran, *Stochastic DCA for the large-sum of non-convex functions problem and its application to group variable selection in classification*, in Proceedings of the 34th International Conference on Machine Learning, D. Precup and Y. W. Teh, eds., vol. 70 of Proceedings of Machine Learning Research, PMLR, 06–11 Aug 2017, pp. 3394–3403, https://proceedings.mlr.press/v70/thi17a.html.

[68] T. van Erven, P. D. Grünwald, N. A. Mehta, M. D. Reid, and R. C. Williamson, *Fast rates in statistical and online learning*, Journal of Machine Learning Research, 16 (2015), pp. 1793–1861.

[69] J. Wang, I. Aravena, and C. G. Petra, *A sequential quadratic programming method for nonsmooth stochastic optimization with upper-C^2 objective*, 2023, https://arxiv.org/abs/2304.04380.

[70] J. Wang and C. G. Petra, *A Sequential Quadratic Programming Algorithm for Nonsmooth Problems with Upper-C^2 Objective*, SIAM Journal on Optimization, 33 (2023), pp. 2379–2405.

[71] Y. Xie, R. Bollapragada, R. Byrd, and J. Nocedal, *Constrained and composite optimization via adaptive sampling methods*, arXiv preprint arXiv:2012.15411, (2020).

[72] H. Xu, *Uniform exponential convergence of sample average random functions under general sampling with applications in stochastic programming*, Journal of Mathematical Analysis and Applications, 368 (2010), pp. 692–710.

[73] H. Xu and D. Zhang, *Smooth sample average approximation of stationary points in nonsmooth stochastic optimization and applications*, Mathematical Programming, 119 (2009), pp. 371–401.

[74] Y. Xu, Q. Qi, Q. Lin, R. Jin, and T. Yang, *Stochastic optimization for DC functions and non-smooth non-convex regularizers with non-asymptotic convergence*, in Proceedings of the 36th International Conference on Machine Learning, ICML 2019, 9-15 June 2019, Long Beach, California, USA, K. Chaudhuri and R. Salakhutdinov, eds., vol. 97 of Proceedings of Machine Learning Research, PMLR, 2019, pp. 6942–6951, http://proceedings.mlr.press/v97/xu19c.html.

[75] Y. Yao, Q. Lin, and T. Yang, *Large-scale optimization of partial auc in a range of false positive rates*, 2022, https://arxiv.org/abs/2203.01505.

[76] M. Zinkevich, *Online convex programming and generalized infinitesimal gradient ascent*, in Proceedings of the Twentieth International Conference on International Conference on Machine Learning, ICML'03, AAAI Press, 2003, pp. 928–935.