



Fossil fuel combustion and PM_{2.5} mass air pollution associations with mortality

George D. Thurston

Department of Environmental Medicine, New York University Grossman School of Medicine, 341 East 25th Street, New York, NY 10010-2598, USA

ARTICLE INFO

Handling Editor: Hanna Boogaard

It is widely acknowledged that outdoor fine particulate matter (PM_{2.5}) air pollution is a major global public health threat: the World Health Organization has ascribed 4.2 million premature cardiovascular, respiratory, and cancer deaths worldwide per year to PM_{2.5}, and recently lowered its PM_{2.5} guidelines to better protect health (World Health Organization (WHO), 2021a; World Health Organization, 2021b). PM_{2.5} is regulated in many countries across the globe, and has been deemed the most consistent and robust predictor of mortality in evaluations of long-term exposure to air pollution (Cohen et al., 2017). Less well appreciated, however, is the fact that not all particles we breathe are the same, having differing sources, compositions, and, therefore, likely also having varying associated health implications per unit mass (National Academy of Sciences, 1999; National Research Council, 1980). As a result, the PM_{2.5} pollution control measures that governments and businesses choose to focus their control efforts upon may well have varying levels of health benefits.

Unfortunately, to date, too few PM_{2.5} health studies have attempted to parse the PM_{2.5} mass health impacts to specific constituents, mixtures of constituents, or particle emission source, mainly due to lack of PM_{2.5} composition data. The earliest such chronic PM_{2.5} mortality study was a cross-sectional analysis I co-authored in 1987, looking at US particulate matter and annual mortality during 1980, based on a brief nationwide Inhalable Particle (IP) monitoring network (Ozkaynak and Thurston, 1987). We found that the mass of smaller sized particles, which we called fine particulate matter (before it was named PM_{2.5}), was more associated with annual mortality than larger particle mass. But, in that work, I also conducted a source apportionment of those effects, concluding that particles from coal combustion were suggested to be more significant contributors to US human mortality than soil-derived particles. The US PM_{2.5} - mortality association was later rigorously confirmed by cohort studies (Dockery et al., 1993; Pope et al., 2002).

The shutdown of the IP Network led to many years passing before the 1999 initiation of the US EPA's Chemical Speciation Network again allowed such nationwide PM_{2.5} source and compositional studies to be conducted to further investigate our earlier findings. The Health Effects Institute (HEI) started the National Particle Component Toxicity (NPACT) initiative, a comprehensive research program in the US that combined epidemiologic and toxicologic approaches to evaluate whether some components and or sources of the PM mixture are of greater public health concern than others (Lippmann et al., 2013; Vedal et al., 2013). Results indicated that PM_{2.5} from coal and oil combustion, and from traffic sources, were most strongly associated across adverse health outcomes, but particles from other sources could not be definitively excluded as having no adverse effects.

This edition of Environment International presents two new studies considering PM_{2.5} source-specific mass, PM_{2.5} constituents, and mortality. Both studies consider the US Medicare population database, although Wang and colleagues focus on the American Southeast (Wang et al., 2021). They conducted a population-based study, comprising Medicare enrollees (aged 65 or older) from 2000 to 2016, in order to explore the all-cause mortality associations with long-term exposures to five PM_{2.5} major components (carbon, nitrate, organic matter, sulfate, and soil particles). While the analyses found that all five PM_{2.5} components were statistically significantly associated with increased risk of mortality, the soil particle mass was noted as having a much smaller effect than the combustion-related PM_{2.5} constituents, with the strongest associations found for sulfates and elemental carbon, proxies of fossil fuel/coal burning and (diesel) traffic PM_{2.5}, respectively, in the US.

Kazemiparkouhi and colleagues (Kazemiparkouhi et al., 2022) analyzed cause-specific mortality among all US Medicare beneficiaries between 2000 and 2008 with trace PM_{2.5} constituents and a full source apportionment of PM_{2.5}. Despite using differing exposure estimation

Abbreviations: PM_{2.5}, fine particulate matter.

E-mail address: george.thurston@nyu.edu.

<https://doi.org/10.1016/j.envint.2021.107066>

Received 19 December 2021; Accepted 22 December 2021

Available online 30 December 2021

0160-4120/© 2021 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

approaches, the two papers concur that the mortality effects of soil particles are much smaller than for other PM_{2.5} components. In addition, the consideration of multiple sources in the nationwide analysis allowed for better discernment of associations by specific combustion source categories, indicating that biomass combustion PM_{2.5} has lower impacts per Interquartile Range (IQR) than PM_{2.5} from fossil fuel combustion sources, especially coal burning and traffic-derived PM_{2.5}. Furthermore, the Kazemiparkouhi et al. source-specific PM_{2.5} findings for ischemic heart disease (IHD) deaths are reassuringly confirmatory to those previously reported by me and my colleagues (Thurston et al., 2016) when we similarly considered PM_{2.5} mass constituent and source contribution impacts in the American Cancer Society (ACS) cohort, also finding the strongest IHD mortality associations with coal combustion and traffic PM_{2.5}, and the least with biomass burning and soil PM_{2.5}.

Taken together, these two new studies compellingly confirm past analyses indicating that the short- and long-term health effects of PM_{2.5} can vary as a function of composition (Achilleos et al., 2017; Wang et al., 2014; Thurston et al., 2021), but also further suggest that analyzing the health effects of PM_{2.5} as a function of source-specific mass may yield more consistent associations than with individual PM_{2.5} constituents. This is likely due to the fact that individual constituents, such as copper or iron or elemental carbon, can derive from multiple sources, each source with its own characteristic aerosol mixture health impacts per PM_{2.5} elemental constituent. Other recent toxicological and epidemiological evidence has indicated, consistent with these new study results, that the co-presence of acidic sulfur with metals, such as emitted by fossil fuel combustion sources, represent an especially toxic aerosol mixture (Fang et al., 2017; Weichenthal et al., 2021; Maciejczyk et al., 2021). A recent HEI sector specific analysis of PM_{2.5} mortality has adopted an assumption of equitoxicity (i.e., every atmospheric particle has the same toxicity per unit mass, regardless of its chemical composition and physical properties) (McDuffie et al., 2021). The above considerations suggest that source sector specific PM_{2.5} concentration–response coefficients are needed in future sector specific health effects analyses, and that the equitoxicity assumption may presently be underestimating the mortality contributions from fossil fuel combustion PM_{2.5}, while overestimating the mortality impacts of windblown dust and biofuel burning PM_{2.5}.

More PM_{2.5} composition and health studies are needed to better quantify PM_{2.5} concentration – health response risks as a function of air pollution source sector. However, these two new studies, combined with other available toxicological and epidemiologic evidence, already indicate to me that, in order to maximize the health benefits of air pollution regulations, a priority should be placed on controlling PM_{2.5} air pollution from fossil-fuel combustion.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Achilleos, S., Kioumourtoglou, M.A., Wu, C.D., Schwartz, J.D., Koutrakis, P., Papathodorou, S.I., 2017. Acute effects of fine particulate matter constituents on mortality: a systematic review and meta-regression analysis. *Environ. Int.* 109, 89–100. <https://doi.org/10.1016/j.envint.2017.09.010>.
- Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbard, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope 3rd, C.A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C.J.L., Forouzanfar, M.H., 2017. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 389 (10082), 1907–1918. [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6). Epub 2017 Apr 10.
- Dockery, D.W., Pope, C.A., Xu, X., Spengler, J.D., Ware, J.H., Fay, M.E., Ferris, B.G., Speizer, F.E., 1993. An association between air pollution and mortality in six US cities. *N. Engl. J. Med.* 329 (24), 1753–1759.
- Fang, T., Guo, H., Zeng, L., Verma, V., Nenes, A., Weber, R.J., 2017. Highly acidic ambient particles, soluble metals, and oxidative potential: a link between sulfate and aerosol toxicity. *Environ. Sci. Technol.* 51 (5), 2611–2620. <https://doi.org/10.1021/acs.est.6b06151>. <https://doi.org/10.1021/acs.est.6b06151.s001>.
- Kazemiparkouhi, F., Honda, T., Eum, K.-D., Wang, B., Manjourides, J., Suh, H.H., 2022. The impact of long-term PM_{2.5} constituents and their sources on specific causes of death in a US medicare cohort. *Environ. Int.* 159, 106988. <https://doi.org/10.1016/j.envint.2021.106988>.
- Lippmann, M., Chen, L.-C., Gordon, T., Ito, K., Thurston, G.D., 2013. National Particle Component Toxicity (NPACT) Initiative: Integrated Epidemiologic and Toxicologic Studies of the Health Effects of Particulate Matter Components. Research Report 177. Health Effects Institute, Boston, MA.
- Maciejczyk, P., Chen, L.-C., Thurston, G., 2021. The role of fossil fuel combustion metals in PM_{2.5} air pollution health associations. *Atmosphere* 12 (9), 1086. <https://doi.org/10.3390/atmos12091086>.
- McDuffie, E., Martin, R.V., Yin, H., Brauer, M., 2021. Global Burden of Disease from Major Air Pollution Sources (GBD MAPS): A Global Approach. Research Report 210. Health Effects Institute, Boston, Massachusetts. <https://www.healtheffects.org/publication/global-burden-disease-major-air-pollution-sources-gbd-maps-global-approach>.
- National Academy of Sciences, 1999. Research Priorities for Airborne Particulate Matter: I. Immediate Priorities and a Long-Range Research. The National Academies Press, Washington, DC. <http://nap.edu/6131>.
- National Research Council, 1980. Controlling Airborne Particles. The National Academies Press, Washington, DC. <https://doi.org/10.17226/19770>.
- Ozkaynak, H., Thurston, G.D., 1987. Associations between 1980 U.S. mortality rates and alternative measures of airborne particle concentration. *Risk Anal.* 7 (4), 449–461.
- Pope III, C.A., Burnett, R.T., Thun, M.J., Calle, E.E., Krewski, D., Ito, K., Thurston, G.D., 2002. Lung cancer, cardiopulmonary mortality and long-term exposure to fine particulate air pollution. *J. Am. Med. Assoc. (JAMA)* 287 (9), 1132–1141. PMID: 11879110.
- Thurston, G., Awe, Y., Ostro, B., Sanchez-Triana, E., 2021. Are All Air Pollution Particles Equal? How Constituents and Sources of Fine Air Pollution Particles (PM_{2.5}) Affect Health. World Bank, Washington, DC.
- Thurston, G.D., Burnett, R.T., Turner, M.C., Shi, Y., Krewski, D., Lall, R., Ito, K., Jerrett, M., Gapstur, S.M., Diver, W.R., Pope, C.A., 2016. Ischemic heart disease mortality and long-term exposure to source-related components of U.S. fine particle air pollution. *Environ. Health Perspect.* 124 (6), 785–794.
- Vedal, S., Campen, M.J., McDonald, J.D., Kaufman, J.D., Larson, T.V., Sampson, P.D., Sheppard, L., Simpson, C.D., Szpiro, A.A., 2013. National Particle Component Toxicity (NPACT) Initiative Report on Cardiovascular Effects. Research Report 178. Health Effects Institute, Boston, MA.
- Wang, M., Beelen, R., Stafoggia, M., Raaschou-Nielsen, O., Andersen, Z.J., Hoffmann, B., Fischer, P., Houthuijs, D., Nieuwenhuijsen, M., Weinmayr, G., Vineis, P., Xun, W.W., Dimakopoulou, K., Samoli, E., Laatikainen, T., Lanki, T., Turunen, A.W., Oftedal, B., Schwarze, P., Aamodt, G., Penell, J., De Faire, U., Korek, M., Leander, K., Pershagen, G., Pedersen, N.L., Östenson, C.-G., Fratiglioni, L., Eriksen, K.T., Sørensen, M., Tjønneland, A., Bueno-de-Mesquita, B., Eeftens, M., Bots, M.L., Meliefste, K., Krämer, U., Heinrich, J., Sugiri, D., Key, T., de Hoogh, K., Wolf, K., Peters, A., Cyrys, J., Jaensch, A., Concin, H., Nagel, G., Tsai, M.-Y., Phuleria, H., Ineichen, A., Künzli, N., Probst-Hensch, N., Schaffner, E., Vilier, A., Clavel-Chapelon, F., Declercq, C., Ricceri, F., Sacerdote, C., Marcon, A., Galassi, C., Migliore, E., Ranzi, A., Cesaroni, G., Badaloni, C., Forastiere, F., Katsoulis, M., Trichopoulos, A., Keuken, M., Jedynska, A., Kooter, I.M., Kukkonen, J., Sokhi, R.S., Brunekreef, B., Katsouyanni, K., Hoek, G., 2014. M. Wang, R. Beelen, S. Massimo Stafoggia, O. Raaschou-Nielsen, Z. Jovanovic Andersen, B. Hoffmann, P. Fischer, D. Houthuijs et al, Long term exposure to elemental constituents of particulate matter and cardiovascular mortality in 19 European cohorts. *Environ. Int.* 66, 97–106. <https://doi.org/10.1016/j.envint.2014.01.026>.
- Wang, Y., Xiao, S., Zhang, Y., Chang, H., Martin, R.V., Van Donkelaar, A., Gaskins, A., Liu, Y., Liu, P., Shi, L., 2021. Long-term exposure to PM_{2.5} major components and mortality in the southeastern United States. *Environ. Int.* 158, 106969. <https://doi.org/10.1016/j.envint.2021.106969>. Epub ahead of print. PMID: 34741960.
- Weichenthal, S., Lavigne, E., Traub, A., Umbrio, D., You, H., Pollitt, K., Shin, T., Kulka, R., Stieb, D.M., Korsiak, J., Jessiman, B., Brook, J.R., Hatzopoulou, M., Evans, G., Burnett, R.T., 2021. Association of sulfur, transition metals, and the oxidative potential of outdoor PM_{2.5} with acute cardiovascular events: a case-crossover study of Canadian adults. *Environ. Health Perspect.* 129 (10) <https://doi.org/10.1289/EHP9449>.
- World Health Organization (WHO), 2021a. Ambient (outdoor) Air Pollution Fact Sheet. 22 September 2021. [https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health).
- World Health Organization, 2021b. WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization. <https://apps.who.int/iris/handle/10665/345329>.