

# Week 1

## Supervised Learning and Regression

# Week 1.1

## Models and Parameters

# Supervised Learning

x: inputs

y: outputs

Training data:

(x = 1, y = 1)

(x = 2, y = 4)

(x = 3, y = 9)

Test data:

(x = 4, y = ?)

*estimate the relationship.*

# Supervised Learning

Training data: *input and output  
could be anything.*

(x = , y = 'cat')

(x = , y = 'dog')

(x = , y = 'cat')


Test Data:

(x = , y = ?)

# Models

In ML, a Model is a function that relates some inputs ( $x$ ) to some outputs ( $y$ )

Most models have parameters ( $\theta$ ), which allows them to represent whole classes of functions.



$$\hat{y} = ax^2 + bx + c$$

$$\hat{y} = a \log(x) + b$$


...

# Linear regression

$$\hat{y} = b_0 + b_1x$$

Diagram illustrating the components of the linear regression equation  $\hat{y} = b_0 + b_1x$ :

- $\hat{y}$  is labeled as the **Output**.
- $b_0$  and  $b_1$  are collectively labeled as **Parameters**.
- $x$  is labeled as the **Input**.



# Fitting a Model

1. Define the model: Choose the “class” of functions that relates the inputs ( $x$ ) to the output ( $y$ )
2. Define your **training loss**
3. **Find the function** in your class/form that gives the **smallest training loss**

*what's a good function/parameter?*

# Week 1.2

## Loss functions




# Fitting a Model

1. Define the model: Choose the “class” of functions that relates the inputs ( $x$ ) to the output ( $y$ )

# Applying the Model to Data

$\% \text{ in poverty} \rightarrow \hat{y} = b_0 + b_1x$   $\swarrow$   $\% \text{ HS Grad.}$

\* But there's no indicies that %HS Grad lead to % in poverty.




which one is the best?  
We need a standard

# Training loss function

- A **training loss function** measures the deviation of the model fits from the observed data
- A large loss indicates a poor fit to the training data
- Different loss functions penalize different deviations differently
- We find the parameters that minimize our loss function given the training data

# Residuals

Residuals are the errors from the model fit:  $\text{Data} = \text{Fit} + \text{Residual}$



# A criterion for the best line

- We want a line that has small residuals
  1. Option 1: Minimize the sum of magnitudes (absolute values) of residuals: The  $L_1$ -norm

$$L(\theta) = \sum_{i=1}^n |y_i - \hat{y}_i| = \sum_{i=1}^n |r_i| = \|\mathbf{r}\|_1$$

LAD: Least Absolute Deviation *(Median regression)*

2. Option 2: Minimize the sum of squared residuals: The squared  $L_2$ -norm

$$L(\theta) = \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n r_i^2 = \|\mathbf{r}\|_2^2$$

OLS: Ordinary Least Squares

*it could be written in vector form.*

- The most commonly used is least squares
  1. Motivated by normal distribution of errors
  2. Solutions can be easily computed
  3. Big errors count relatively more than small errors

Week 1.3


Optimization

Finding the best model fit

# Optimization

- Finding the best *function* is the same problem as finding the best *parameters*.
- Parameter estimation is the process of minimizing the training loss by trying different values of the parameters
- The setting of parameters that gives you the smallest loss is the best estimate of the parameters, and the best function in your class

# Optimization



## Analogy:

Blindfolded hiker in the mountains who is trying to get downhill

Here the parameter was a single number. In linear regression we have 2 parameters, making the loss-function a surface



# Using the derivative of the loss

By providing the derivative of the loss function in respect to the parameters, optimization can be sped up.

$$\text{Fit: } \hat{y}_i = b_0 + b_1 x_i$$


$$\text{Residual: } r_i = y_i - \hat{y}_i$$

$$\frac{\partial \sum f_i(\theta)}{\partial \theta} = \sum \frac{\partial f_i(\theta)}{\partial \theta}$$

$$\text{Loss: } L = \sum_{i=1}^N (y_i - b_0 - b_1 x_i)^2$$

$$\text{Derivative } b_0: \frac{\partial L}{\partial b_0} = -2 \sum_{i=1}^n (y_i - b_0 - b_1 x_i) = -2 \sum_{i=1}^n r_i$$

$$\text{Derivative } b_1: \frac{\partial L}{\partial b_1} = -2 \sum_{i=1}^n (y_i - b_0 - b_1 x_i) x_i = -2 \sum_{i=1}^n r_i x_i$$




$$\frac{\partial f(g(\theta))}{\partial \theta} = \frac{\partial f(g)}{\partial g} \frac{\partial g(\theta)}{\partial \theta}$$

Assuming that all residual are positive, there is, the intercept is too low:

For larger value of intercept ( $b_0$ ), the residual would go down

# Using the derivative of the loss

The vector of partial derivatives is called a gradient or Jacobian.




top is so a vector.


$$\nabla_{\theta} L = \begin{bmatrix} \frac{\partial L}{\partial \theta_0} \\ \frac{\partial L}{\partial \theta_1} \end{bmatrix}$$

partial derivatives  
of  $\theta_0, \theta_1$

# Derivatives of the loss

- Remember: Derivative = slope
- Also remember: want to make loss *small*

• If  $\frac{\partial L}{\partial \theta}$  is positive, should I increase or decrease  $\theta$ ? 

• If  $\frac{\partial L}{\partial \theta}$  is negative, should I increase or decrease  $\theta$ ? 

# Linear regression in matrix notation

$$\hat{y}_i = b_0 + b_1 x_i$$

Fit

Parameters

Input

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_n \end{bmatrix} = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_n \end{bmatrix} \begin{bmatrix} b_0 \\ b_1 \end{bmatrix}$$

$$\hat{\mathbf{y}} = \mathbf{X}\mathbf{b}$$

For each point in the dataset, we get a fit/estimate/prediction from the model. Next, we'll compare those to the actual data.

# Using the derivative (vector notation)

By providing the derivative of the loss function in respect to the parameters, optimization can be sped up.

Prediction:  $\hat{\mathbf{y}} = \mathbf{X}\mathbf{b}$

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_n \end{bmatrix} = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_n \end{bmatrix} \begin{bmatrix} b_0 \\ b_1 \end{bmatrix}$$

Residual:  $\mathbf{r} = \mathbf{y} - \hat{\mathbf{y}}$

Loss:

$$\begin{aligned} L &= (\mathbf{y} - \mathbf{X}\mathbf{b})^\top (\mathbf{y} - \mathbf{X}\mathbf{b}) \\ &= \mathbf{y}^\top \mathbf{y} - 2\mathbf{y}^\top \mathbf{X}\mathbf{b} + \mathbf{b}^\top \mathbf{X}^\top \mathbf{X}\mathbf{b} \end{aligned}$$

Gradient:  $\nabla_{\mathbf{b}} J = -2\mathbf{X}^\top \mathbf{y} + 2\mathbf{X}^\top \mathbf{X}\mathbf{b} = -2\mathbf{X}^\top \mathbf{r}$

## Week 1.4

Implementing OLS regression  
through minimization of L2 loss

# Step 1: Write the model function

Write a function that returns the

*vector*  
First input is parameter  
list or np.array  
(param)

*matrix*  
np.array (2d) of  
explanatory  
variables (input)

```
def linearModelPredict(b,X):  
    # Get Model prediction  
    predY =  
    # return Model prediction  
    return predY
```

# Step 2: Write a loss function

We are modifying the loss function to also return the gradient

*First input is parameter  
list or np.array*

*Explanatory and  
response variable*

```
def linearModelLossRSS(b, X, y):  
    # Get Model prediction  
    predY = linearModelPredict(b, X)  
    # Get the vector of residuals  
    res =  
    # Get the residuals sums of squares  
    rss =  
    # Get the gradient  
    gradient = partial derivative of the loss with given  
    # return rss and gradient param; it should  
    return (rss, gradient) has same dimension  
                                as the input param.  
                                (b).
```



# Step 3: Call the optimizer

```
import scipy.optimize as so
# Set some starting values
bstart=[0,0]
# Call the optimization function
RESULT=so.minimize(linearModelLossRSS,bstart,args=(X, y),jac=True)
```

*if your functions are correct, starting value does not affect the result, but the processing time*

Loss  
function

Starting  
value

Additional  
arguments

Use  
gradient

Remember our definition:

```
def linearModelLossRSS(b,X,y):
```

*return is not only loss, but also gradient. this would speed up the optimization.*

# Step 4: Check the results

## RESULT


*this jacobian here should be close to zero.* *minimal loss value found.*  
**fun:** 14.25 *no zero.* # Final loss function value  
**hess\_inv:** array([[0.33, -0.07], [-0.07, 0.01]])  
*Jacobian* **jac:** array([1.19-06, -1.90-06]) *Jacobian at the place of the best fitting parameter.*  
**message:** 'Optimization terminated successfully.'  
**nfev:** 24 # Number of evaluations  
**nit:** 3 # Number of iterations  
**njev:** 6  
**status:** 0  
**success:** True  
**x:** array([65, -0.6]) # Parameter estimates

- Check if successful
- Get parameter estimates
- Maybe check the loss

# Step 5: Visualize the results

```
b=RESULT.x          # Get the parameters  
x_grid = np.linspace(y.min(), y.max(),10) # get grid  
Xn = np.c_[np.ones(x_grid.size), x_grid] # Make Design  
yp=linearModelPredict(b,X) # get prediction  
ax.plot(x_grid, yp, color = 'red')
```

$$\widehat{\% \text{ in poverty}} = 64.68 - 0.62 \% \text{ HS grad}$$



# Week 1.5

## Evaluating model fit - $R^2$

# Evaluating the fit with $R^2$

- The quality of the fit of a linear regression model is most commonly evaluated using  $R^2$  the **coefficient of determination**.
- $R^2$  is calculated from the ratio of residual sum of squares – total sum of squares.

$$R^2 = 1 - \frac{\text{RSS}}{\text{TSS}} = 1 - \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{\sum_{i=1}^n (y_i - \bar{y})^2}$$

Residual sum of squares (RSS)

Total sum of squares (TSS)


*if it is not 1) data is very noisy.  
perfectly fit: 2) model is overfitted.*

*datapoint - mean of data.*

- It tells us what percent of variability in the response variable is explained by the model. (0=no fit, 1=perfect fit)
- The remainder of the variability is explained by variables not included in the model or by inherent randomness in the data.
- Because OLS is mimicking the RSS, it will always have the highest  $R^2$  value possible for that class of models.

# Summary

- **Models** can be written in general as:

$$\hat{y} = f(x, \theta)$$


Dependent variable      Input      Parameters

- The **training loss function** tells how bad a fit is:

Example: squared error


$$L = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

- **Model fitting** involves selecting the parameters that minimizes the loss function: **Parameter estimation**
- In linear regression, the **proportion of the explained variance** in the response variable is expressed by the **coefficient of determination ( $R^2$ )**

Week 1.6

L1-loss and median regression  
Robust techniques

# The Impact of the Loss Function



Example of best running speeds for a 3km Strava segment.

What do you notice?


Are these old guys for real?

Maybe these data points have other explanations:

- *old guys on bikes*
- *young runners lying about their age*
- *really good doping*
- ....



# The Impact of the Loss Function



Excluding these “outliers” changes the predicted running speed for 80-year olds

Answers that change a lot with exclusion of a few data points are called **non-robust**.

*It is not the most desired try.*

So are we sure that this one is cheater as well?



# Robust statistics

**Robust** statistics seek to provide methods that emulate popular statistical methods, but which are not unduly affected by outliers or other small departures from model assumptions.

- **Mean** is a measure of central tendency that is sensitive to outliers *but it would be greatly affected by so extreme value.*
- **Median** is a measure of central tendency that is robust against outliers

# Robust regression

To develop a robust regression technique, we can think how to change the loss function.



Say we have these 4 data points

If we find the point that minimizes the sum of squared error

$$L = \sum_{i=1}^n (y_i - \underline{b_0})^2$$

$$\frac{\partial L}{\partial b_0} = \sum_{i=1}^n -2(y_i - b_0)$$

$$\frac{\partial L}{\partial b_0} = 0 \Rightarrow b_0 = \frac{\sum_{i=1}^n y_i}{n}$$



*minimize L2 loss.*

The minimum is reached at the **mean**

# Robust regression

To develop a robust regression technique, we can think how to change the cost function.

Say we have these 4 data points



the minimal here is not only one value but a range

If we find the point that minimize the sum of **absolute** errors


$$L = \sum_{i=1}^n |y_i - b_0|$$

$$L = \sum_{i=1}^n \begin{cases} (y_i - b_0) & \text{if } y_i > b_0 \\ -(y_i - b_0) & \text{if } y_i \leq b_0 \end{cases}$$

$$\frac{\partial L}{\partial b_0} = \sum_{i=1}^n \begin{cases} -1 & \text{if } y_i > b_0 \\ 1 & \text{if } y_i \leq b_0 \end{cases}$$

The minimum is reached at the **median**

# Outliers and robustness



Median regression leads to results in a regression line that is closer to the one that excludes the outlier.

	all data	w/o outliers
RSS	————	.....
LAD	————	.....

Median regression is a **robust** regression technique

# Week 1.7

## Implementing Median regression through minimization of L1 loss

# Implementing median regression

```
yp = linearModelPredict(b,x)  
# Computes Prediction
```

↑ *calls*

```
linearModelLossRSS(b,x,y)  
# Computes RSS  
# for linear fit
```

↑ *calls*

```
so.minimize(lossfcn,b0,args=(x,y))  
# Estimates b to  
# minimize lossfcn
```

↑ *calls*

```
LinearModelFit(x,y,lossfcn)
```

← *calls*

```
linearModelLossLAD(b,x,y)  
# Computes summed  
# absolute deviation
```

↗ *calls*

Handing a loss function to your linear regression lets you do normal linear regression (lossfcn = rss) and median regression (lossfcn = lad) with the same code


# Using the derivative

Again, by providing the derivative of the training loss with respect to the parameters, we can make the fit faster.


Prediction:  $\hat{y}_i = b_0 + b_1 x_i$

Residual:  $r_i = y_i - \hat{y}_i$


Loss:  $L = \sum_{i=1}^n |y_i - (b_0 + b_1 x_i)|$



Derivative  $b_0$ :  $\frac{\partial L}{\partial b_0} = - \sum_{i=1}^n \text{sgn}(y_i - (b_0 + b_1 x_i)) = - \sum_{i=1}^n \text{sgn}(r_i)$



Derivative  $b_1$ :  $\frac{\partial L}{\partial b_1} = - \sum_{i=1}^n \text{sgn}(y_i - (b_0 + b_1 x_i)) \cdot x_i = - \sum_{i=1}^n \text{sgn}(r_i) \cdot x_i$





# Parameter estimation

- “Parameter estimation” is the process of minimizing the **loss function** by trying different values of the parameters
- The Gradient can speed up optimization a lot! (But you can get by without it.)
- For nearly every problem there is a more specialized solution that is faster, for example Sklearn-methods usually have a fit (estimation) and predict (prediction) function built in.
- But using general minimization algorithms, such as `scipy.optimize.minimize`, is an incredibly useful and universal tool.

# Summary

- We can fit mathematical models to capture the relationship between  $x$  and  $y$

*line / smooth curve / etc ...*

1. Select a function class/form
2. Select a loss function *← OLS / LAD .*
3. Estimation/ fitting: Find the function that minimizes the loss

- “supervised learning” is a cornerstone of statistics, machine learning, and data science.