

Cross products and Translation

Xin Fu

Western University

xfu82@uwo.ca

Cross product

Definition Let $\vec{u} = (u_1, u_2, u_3)$ and v_1, \vec{v}_2, v_3 in \mathbb{R}^3 . The *cross product* of \vec{u} and \vec{v} is the vector

$$\vec{u} \times \vec{v} = (u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1).$$

Definition Let $\vec{u} = (u_1, u_2, u_3)$ and v_1, \vec{v}_2, v_3 in \mathbb{R}^3 . The *cross product* of \vec{u} and \vec{v} is the vector

$$\vec{u} \times \vec{v} = (u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1).$$

Theorem The cross product $\vec{u} \times \vec{v}$ is a vector in \mathbb{R}^3 which is both orthogonal (or perpendicular) to \vec{u} and \vec{v} . Namely, we have $\vec{u} \cdot (\vec{u} \times \vec{v}) = \vec{v} \cdot (\vec{u} \times \vec{v}) = 0$.

Definition Let $\vec{u} = (u_1, u_2, u_3)$ and v_1, \vec{v}_2, v_3 in \mathbb{R}^3 . The *cross product* of \vec{u} and \vec{v} is the vector

$$\vec{u} \times \vec{v} = (u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1).$$

Theorem The cross product $\vec{u} \times \vec{v}$ is a vector in \mathbb{R}^3 which is both orthogonal (or perpendicular) to \vec{u} and \vec{v} . Namely, we have $\vec{u} \cdot (\vec{u} \times \vec{v}) = \vec{v} \cdot (\vec{u} \times \vec{v}) = 0$.

Example Find a nonzero vector that is both orthogonal to $\vec{u} = (1, 2, 1)$ and $\vec{v} = (2, 0, -3)$.

Definition Let $\vec{u} = (u_1, u_2, u_3)$ and v_1, \vec{v}_2, v_3 in \mathbb{R}^3 . The *cross product* of \vec{u} and \vec{v} is the vector

$$\vec{u} \times \vec{v} = (u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1).$$

Theorem The cross product $\vec{u} \times \vec{v}$ is a vector in \mathbb{R}^3 which is both orthogonal (or perpendicular) to \vec{u} and \vec{v} . Namely, we have $\vec{u} \cdot (\vec{u} \times \vec{v}) = \vec{v} \cdot (\vec{u} \times \vec{v}) = 0$.

Example Find a nonzero vector that is both orthogonal to $\vec{u} = (1, 2, 1)$ and $\vec{v} = (2, 0, -3)$.

$$\begin{aligned}\vec{u} \times \vec{v} &= (u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1) \\ &= (2 \times (-3) - 1 \times 0, 1 \times 2 - 1 \times (-3), 1 \times 0 - 2 \times 2) \\ &= (-6, 5, -4).\end{aligned}$$

A (2×2) -matrix is given by the following

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ where } a, b, c, d \text{ are real numbers.}$$

A (2×2) -matrix is given by the following

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ where } a, b, c, d \text{ are real numbers.}$$

The *determinant* of A is defined by letting

$$\det A = ad - bc.$$

A (2×2) -matrix is given by the following

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ where } a, b, c, d \text{ are real numbers.}$$

The *determinant* of A is defined by letting

$$\det A = ad - bc.$$

By the notation of determinant, the cross product of $\vec{u} \times \vec{v}$ is given by

$$\left(\det \begin{bmatrix} u_2 & u_3 \\ v_2 & v_3 \end{bmatrix}, -\det \begin{bmatrix} u_1 & u_3 \\ v_1 & v_3 \end{bmatrix}, \det \begin{bmatrix} u_1 & u_2 \\ v_1 & v_2 \end{bmatrix} \right)$$

where $\begin{bmatrix} u_2 & u_3 \\ v_2 & v_3 \end{bmatrix}$, $\begin{bmatrix} u_1 & u_3 \\ v_1 & v_3 \end{bmatrix}$ and $\begin{bmatrix} u_1 & u_2 \\ v_1 & v_2 \end{bmatrix}$ are obtained by deleting the first, second and third columns of

$$\begin{bmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{bmatrix}.$$

Theorem Let \vec{u}, \vec{v} and \vec{w} be vectors in \mathbb{R}^3 . Let c be a scalar. Then

$$\vec{u} \times \vec{v} = -\vec{v} \times \vec{u}$$

$$\begin{cases} \vec{u} \times (\vec{v} + \vec{w}) = \vec{u} \times \vec{v} + \vec{u} \times \vec{w} \\ (\vec{u} + \vec{v}) \times \vec{w} = \vec{u} \times \vec{w} + \vec{v} \times \vec{w} \end{cases} \quad (\text{distributive law})$$

$$c(\vec{u} \times \vec{v}) = (c\vec{u}) \times \vec{v} = \vec{u} \times (c\vec{v}) \quad (\text{scalars factor out})$$


$$\vec{u} \times \vec{0} = \vec{0} \times \vec{u} = \vec{0}$$

$$\vec{u} \times \vec{u} = \vec{0}$$


$$\|\vec{u} \times \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin\theta = \sqrt{\|\vec{u}\|^2 \|\vec{v}\|^2 - (\vec{u} \cdot \vec{v})^2}$$

where θ is the angle determined by \vec{u} and \vec{v} .

Theorem The area of the parallelogram determined by \vec{u} and \vec{v} is given by $\|\vec{u} \times \vec{v}\|$.




Theorem The area of the parallelogram determined by \vec{u} and \vec{v} is given by $\|\vec{u} \times \vec{v}\|$.




Example Consider the vectors $\vec{u} = (1, 2, 3)$ and $\vec{v} = (2, 1, 0)$. Find the area of the parallelogram determined by these vectors.

Theorem The area of the triangle determined by \vec{u} and \vec{v} is given by $\frac{1}{2} \|\vec{u} \times \vec{v}\|$.




Theorem The area of the triangle determined by \vec{u} and \vec{v} is given by $\frac{1}{2} \|\vec{u} \times \vec{v}\|$.



Example Find the area of the triangle OAB , where O is the origin, A is the point $(2, -3, 1)$ and B is the point $(4, 6, 2)$.

Theorem The area of the triangle determined by \vec{u} and \vec{v} is given by $\frac{1}{2} \|\vec{u} \times \vec{v}\|$.




Example Find the area of the triangle OAB , where O is the origin, A is the point $(2, -3, 1)$ and B is the point $(4, 6, 2)$.

How about the area of the triangle $O'AB$, where O' is the point $(1, 4, -2)$ and A, B are the same as above?

Translation


Directed line segment \overrightarrow{PQ}

For points P and Q in \mathbb{R}^2 or \mathbb{R}^3 , we denote the directed line segment from P to Q by \overrightarrow{PQ} .




Directed line segment \overrightarrow{PQ}

For points P and Q in \mathbb{R}^2 or \mathbb{R}^3 , we denote the directed line segment from P to Q by \overrightarrow{PQ} .




Definition Two directed line segments \overrightarrow{PQ} and \overrightarrow{ST} are *equivalent* if they have the same direction and length.




Theorem Let \overrightarrow{PQ} be a directed line segment from P to Q , where P and Q are two distinct points in \mathbb{R}^2 or \mathbb{R}^3 . Then \overrightarrow{PQ} is equivalent to the vector $\vec{u} - \vec{v}$, where $\vec{u} = \overrightarrow{OQ}$ and $\vec{v} = \overrightarrow{OP}$ and O denotes the origin.

Theorem Let \overrightarrow{PQ} be a directed line segment from P to Q , where P and Q are two distinct points in \mathbb{R}^2 or \mathbb{R}^3 . Then \overrightarrow{PQ} is equivalent to the vector $\vec{u} - \vec{v}$, where $\vec{u} = \overrightarrow{OQ}$ and $\vec{v} = \overrightarrow{OP}$ and O denotes the origin.




Theorem Let \overrightarrow{PQ} be a directed line segment from P to Q , where P and Q are two distinct points in \mathbb{R}^2 or \mathbb{R}^3 . Then \overrightarrow{PQ} is equivalent to the vector $\vec{u} - \vec{v}$, where $\vec{u} = \overrightarrow{OQ}$ and $\vec{v} = \overrightarrow{OP}$ and O denotes the origin.



Remark The process of replacing \overrightarrow{PQ} by the vector $\vec{u} - \vec{v}$ is called *translating P to the origin*.

Definition The process of replacing a directed line segment \overrightarrow{PQ} with the equivalent vector is called *translating* \overrightarrow{PQ} to the origin. Similarly, the process of replacing the vector \vec{v} with an equivalent directed line segment which starts at some point P is called translating \vec{v} to P .

Definition The process of replacing a directed line segment \overrightarrow{PQ} with the equivalent vector is called *translating* \overrightarrow{PQ} to the origin. Similarly, the process of replacing the vector \vec{v} with an equivalent directed line segment which starts at some point P is called translating \vec{v} to P.




*Pic is from the online note.

Example

Find the area of the triangle $O'AB$, where O' is the point $(1, 4, -2)$, A is the point $(2, -3, 1)$ and B is the point $(4, 6, 2)$.

Definition The **parallelepiped** determined by vectors \vec{u} , \vec{v} and \vec{w} is the 6-faced solid whose faces are the parallelograms determined by \vec{u} and \vec{v} , by \vec{u} and \vec{w} , and by \vec{v} and \vec{w} .



Theorem The volume of the parallelepiped determined by the vectors \vec{u} , \vec{v} and \vec{w} is given by

$$\text{Volume} = |(\vec{u} \times \vec{v}) \cdot \vec{w}|.$$

Theorem The volume of the parallelepiped determined by the vectors \vec{u} , \vec{v} and \vec{w} is given by

$$\text{Volume} = |(\vec{u} \times \vec{v}) \cdot \vec{w}|.$$

Example Find the volume of the parallelepiped determined by the directed line segments \overrightarrow{AB} , \overrightarrow{AC} and \overrightarrow{AD} for the points $A(1, 0, 1)$, $B(2, 1, 1)$, $C(2, 2, 1)$ and $D(1, 1, 2)$.