
Systems of Linear Equations


In \mathbb{R}^2 , we have seen that the standard form of a line is ax + by = c.

Of course, we also have other equations, for instance $2x - y^2 = 0$, $x^2 + y^2 = 1$.

In \mathbb{R}^2 , we have seen that the standard form of a line is ax + by = c.

Of course, we also have other equations, for instance $2x - y^2 = 0$, $x^2 + y^2 = 1$.





They are NOT straight lines in $\mathbb{R}^2!$

Definition A *linear equation* in the variables $x_1, x_2, \dots x_n$ is an equation which can be put into the form

$$a_1x_1+a_2x_2+\ldots+a_nx_n=b$$

for some constants a_1, a_2, \ldots, a_n and b, where not all of the a_i values are 0.

Definition A *linear equation* in the variables $x_1, x_2, ... x_n$ is an equation which can be put into the form

$$a_1x_1+a_2x_2+\ldots+a_nx_n=b$$

for some constants a_1, a_2, \ldots, a_n and b, where not all of the a_i values are 0.

- You could think that x_i are unknown.
- An *n*-vector $\vec{v} = (v_1, v_2, \dots, v_n)$ is a solution to a linear equation if it satisfies the linear equation.
- If n=3, all the solutions to a linear equation ax+by+cz=d form a **PLANE** in \mathbb{R}^3 .
- For any positive integer $n \ge 2$, all the solutions to a linear equation $a_1x_1 + a_2x_2 + \ldots + a_nx_n = b$ forms a **HYPERPLANE**.

Examples

Are these equations linear or not?

1.
$$x_1x_2 + x_3 = \sqrt{7}$$
:

2.
$$x_1 + x_2 + \sqrt{x_4} = 3$$
;

3.
$$sinx+y=0$$
;

4.
$$2x_3 + \frac{2}{3}x_6 = \cos\frac{\pi}{3}$$
;

5.
$$x_1 + \frac{2}{x_4} - x_6 = 0$$
;

6.
$$\sqrt[3]{12}x_1 + 2x_2 - x_3 = -3$$
;

7.
$$\sqrt[3]{12x_1} + 2x_2 - x_5 = -3$$
;

System of linear equations

A collection of a finite number of linear equations involving the same variables is called a *system of linear equations* or ($linear\ system$), abbreviated as **SLE**. For example, we have m linear equations and each equation has n variables.

System of linear equations

A collection of a finite number of linear equations involving the same variables is called a *system of linear equations* or (*linear system*), abbreviated as **SLE**.

For example, we have m linear equations and each equation has n variables.

- A system of m linear equations in n variables is called an $m \times n$ SLE (pronounced "m by n").
- An *n*-vector (v_1, v_2, \dots, v_n) is called a solution to a particular SLE with *n* variables if it satisfies all of the equations in the system.

For example, we did the question to find the intersection of the line 2x+y=1 and the line x-2y=4.

Examples

Is each system of equations linear?

1.
$$x_1 + x_4 = 1$$

 $x_2 - x_3 + 2x_4 = 2$
 $x_1 - x_2x_3 + x_4 = 0$
2. $(sin2)x_1 + x_2 + 9x_3 = cos3$
 $2x_1 + cosx_2 - x_3 = \sqrt{2}$
 $\frac{1}{2}x + (cos4)y - z = 1$

3.
$$\frac{1}{2}x + \cos 4y - z = 1$$

 $x + y + z = 1$

Standard form

A SLE is in standard form if

- all of the equations have all of the variables appearing on the left hand side of the equation;
- all the variables appear in the same order for all equations, with spaces left for any variables missing in that equation;
- the constant term appears on the right hand side of each equation.

Standard form

A SLE is in standard form if

- all of the equations have all of the variables appearing on the left hand side of the equation;
- all the variables appear in the same order for all equations, with spaces left for any variables missing in that equation;
- the constant term appears on the right hand side of each equation.

$$x + 2y + z = 5$$
$$-2 - x + z = 0$$

Standard form

A SLE is in standard form if

- all of the equations have all of the variables appearing on the left hand side of the equation;
- all the variables appear in the same order for all equations, with spaces left for any variables missing in that equation;
- the constant term appears on the right hand side of each equation.

$$x + 2y + z = 5$$
$$-2 - x + z = 0$$

A standard-form SLE is

$$x + 2y + z = 5$$
$$-x + z = 2$$

If a SLE has at least one solution, then it is called consistent.

If a SLE has at least one solution, then it is called consistent.

1. Consider the SLE

$$x + y - z = 0$$

$$x + y - z = 1$$

If a SLE has at least one solution, then it is called *consistent*.

1. Consider the SLE

$$x + y - z = 0$$
$$x + y - z = 1$$

It has no solution. They are two parallel planes in \mathbb{R}^3 . So it is inconsistent.

If a SLE has at least one solution, then it is called *consistent*.

1. Consider the SLE

$$x + y - z = 0$$
$$x + y - z = 1$$

It has no solution. They are two parallel planes in $\mathbb{R}^3.$ So it is inconsistent. 2.

$$x + 3y - z = 0$$
$$2y - z = 1$$

If a SLE has at least one solution, then it is called *consistent*.

1. Consider the SLE

$$x + y - z = 0$$
$$x + y - z = 1$$

It has no solution. They are two parallel planes in $\mathbb{R}^3.$ So it is inconsistent. 2.

$$x + 3y - z = 0$$
$$2y - z = 1$$

It has "many" solutions. Any vectors in a form of (-t-1,t,2t-1) are solutions, where t can be any arbitrary real numbers. It is consistent.

Examples

Put the following SLEs into a standard form. Are they consistent or inconsistent?

1)
$$2 - x + 3y = -1$$
$$x = z + 2$$

2)
$$x_1 - 2x_2 = 0$$
$$x_2 + 3x_3 = 2$$
$$x_1 + x_3 = -1$$

3)
$$1 + 4y - x = 0$$
$$8y + 2 = 2x$$

r-parameter family of solutions

Definition When a solution to a system of equations can be stated with one or more parameters, such that any value of the parameter(s) gives a solution to the system, the system is said to have an r-parameter family of solutions, where r is the number of parameters in the solution.

For example, the SLE

$$x + 3y - z = 0$$
$$2y - z = 1$$

has 1-parameter family of solutions (-t-1,t,2t-1) for any $t\in\mathbb{R}$.

r-parameter family of solutions

Definition When a solution to a system of equations can be stated with one or more parameters, such that any value of the parameter(s) gives a solution to the system, the system is said to have an r-parameter family of solutions, where r is the number of parameters in the solution.

For example, the SLE

$$x + 3y - z = 0$$
$$2y - z = 1$$

has 1-parameter family of solutions (-t-1,t,2t-1) for any $t\in\mathbb{R}$.

Definition Two SLEs are *equivalent* if they have the same solutions.