

# Basic Structures: Sets, Functions, Sequences, Sums, and Matrices

## Chapter 2

© Peter Valovcik 2021

UWO – February 2, 2021

# Basic Structures: Sets, Functions, Sequences, Sums, and Matrices

## Chapter 2

© Peter Valovcik 2021

UWO – February 2, 2021

# Introduction

- ① Sets are the basic building blocks for the types of objects considered in discrete mathematics and in mathematics, in general.
- ② Set theory is an important branch of mathematics:
  - ⓐ Many different systems of axioms have been used to develop set theory.
  - ⓑ Here, we are not concerned with a formal set of axioms for set theory.
  - ⓒ Instead, we will use what is called naïve set theory.

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Plan for Part I

## 1. Sets

### 1.1 Defining sets

### 1.2 Venn Diagram

### 1.3 Set Equality

### 1.4 Subsets

### 1.5 Venn Diagrams and Truth Sets

### 1.6 Set Cardinality

### 1.7 Power Sets

### 1.8 Cartesian Products

## 2. Set Operations

### 2.1 Boolean Algebra

### 2.2 Union

### 2.3 Intersection

### 2.4 Complement

### 2.5 Difference

### 2.6 The Cardinality of the Union of Two Sets

### 2.7 Set Identities

### 2.8 Generalized Unions and Intersections

# Sets

- ① A set is an unordered collection of “objects”, e.g. intuitively described by some common property or properties (in *naïve set theory*):
  - a the students in this class,
  - b the chairs in this room.
- ② The objects in a set are called the elements, or members of the set. A set is said to *contain* its elements.
- ③ The notation  $a \in A$  denotes that  $a$  is an element of set  $A$ .
- ④ If  $a$  is not a member of  $A$ , write  $a \notin A$

# Describing a set: the roster method

The roster method is defined as a way to show the members of a set by listing the members inside of brackets.

- ① Example:  $S = \{a, b, c, d\}$   *$a \in S, e \notin S$ .*
- ② The order of the elements in that list is not important.
  - Ⓐ For instance,  $\{a, b, c, d\} = \{b, c, a, d\}$
- ③ Each object in the universe is either a member or not.
- ④ Listing a member more than once does not change the set.
  - Ⓐ For instance,  $\{a, b, c, d\} = \{a, b, c, b, c, d\}$
- ⑤ *Ellipses* (...) may be used to describe a set without listing all of the members when the pattern is clear: *arguable.*
  - Ⓐ For instance,  $S = \{a, b, c, d, \dots, z\}$ .

# The roster method: more examples

- ① The set of all vowels in the English alphabet:

$$V = \{a, e, i, o, u\}$$

- ② The set of all odd positive integers less than 10:

$$O = \{1, 3, 5, 7, 9\}$$

- ③ The set of all positive integers less than 100:

$$S = \{1, 2, 3, \dots, 99\}$$

- ④ The set of all integers less than 0:

$$S = \{\dots, -3, -2, -1\}$$



# Some important sets

$\mathbb{N}$	= <i>natural numbers</i>	= $\{0, 1, 2, 3, \dots\}$
$\mathbb{Z}$	= <i>integers</i>	= $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
$\mathbb{Z}^+$	= <i>positive integers</i>	= $\{1, 2, 3, \dots\}$
$\mathbb{Z}^-$	= <i>negative integers.</i>	
$\mathbb{Q}$	= set of <i>rational numbers</i>	
$\mathbb{R}$	= set of <i>real numbers</i>	
$\mathbb{R}^+$	= set of <i>positive real numbers</i>	
$\mathbb{C}$	= set of <i>complex numbers.</i>	

# The set-builder notation

- 1 It is used to specify the property or properties that all members must satisfy. *Examples:*

- a  $S = \{x \mid x \text{ is a positive integer less than } 100\}$

- b  $O = \{x \mid x \text{ is an odd positive integer less than } 10\}$

- c  $O = \{x \in \mathbb{Z}^+ \mid x \text{ is odd and } x < 10\}$

- d the set of positive rational numbers:

$$\mathbb{Q}^+ = \{x \in \mathbb{R} \mid x = \frac{p}{q}, \text{ for some positive integers } p, q\}$$

- 2 A predicate may be used, as in  $S = \{x \mid P(x)\}$

- a *Example:*  $S = \{x \mid \text{Prime}(x)\}$

# The interval notation

- ① It is used to specify a range of real numbers.
- ② Consider two real numbers  $a, b$  with  $a \leq b$ . The following notations are commonly used:
  - ▶  $[a, b] = \{x \mid a \leq x \leq b\}$
  - ▶  $[a, b) = \{x \mid a \leq x < b\}$
  - ▶  $(a, b] = \{x \mid a < x \leq b\}$
  - ▶  $(a, b) = \{x \mid a < x < b\}$
  - ▶ *closed interval*  $[a, b]$
  - ▶ *open interval*  $(a, b)$

# Truth sets of quantifiers

- ① Given a predicate  $P$  and a domain  $D$ , we define the *truth set* of  $P$  to be the set of the elements in  $D$  for which  $P(x)$  is true.
- ② The truth set of  $P(x)$  is denoted by:

$$\{x \in D \mid P(x)\}$$

- ③ **Example:** The truth set of  $P(x)$  where the domain is the integers and  $P(x)$  is “ $|x| = 1$ ” is the set  $\{-1, 1\}$

# Sets can be elements of sets

Examples:

①  $\{\{1,2,3\}, a, \{b, c\}\}$

②  $\{\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}\}$

# Russell's paradox

- 1 Let  $S$  be the set of all sets which are not members of themselves.
- 2 A paradox results from trying to answer the question “Is  $S$  a member of itself?”

Related simple example:

- a Henry is a barber who shaves all people who do not shave themselves. A paradox results from trying to answer the question “Does Henry shave himself?”



Bertrand Russell

(1872 - 1970)

Nobel Prize

Winner,

Cambridge, UK

- 1 To avoid this and other paradoxes, *sets* can be (formally) defined via appropriate axioms more carefully than just *an unordered collection of “objects”*
- 2 where *objects* are intuitively described by any given property *in naïve set theory*

# The universal set $U$ and the empty set $\emptyset$

- 1 The *universal set* is the set containing all the “objects” currently under consideration:
  - a often symbolized by  $U$ ,
  - b sometimes implicitly stated,
  - c sometimes explicitly stated,
  - d its contents depend on the context.
- 2 The *empty set* is the set with no elements.
  - a symbolized by  $\emptyset$ , but  $\{\}$  is also used.
  - b **Important:** the empty set is different from a set containing the empty set:

$$\emptyset \neq \{ \emptyset \}$$

# Plan for Part I

## 1. Sets

1.1 Defining sets

1.2 Venn Diagram

1.3 Set Equality

1.4 Subsets

1.5 Venn Diagrams and Truth Sets

1.6 Set Cardinality

1.7 Power Sets

1.8 Cartesian Products

## 2. Set Operations

2.1 Boolean Algebra

2.2 Union

2.3 Intersection

2.4 Complement

2.5 Difference


2.6 The Cardinality of the Union of Two Sets

2.7 Set Identities

2.8 Generalized Unions and Intersections




# Venn Diagram



John Venn (1834 -

1923) Cambridge, UK

Sets and their elements can be represented via Venn diagrams




□ – Universal set  $U$

● – elements


○ – Some set  $V$

# Venn Diagram



John Venn (1834 -


1923) Cambridge, UK




□ – Universal set  
 $U$ : all letters in the  
Latin alphabet

**letters** – elements  
○ – Set  $V$ : all vowels

# Venn Diagram



John Venn (1834 -  
1923) Cambridge, UK



- – Universal set  $U$
- – Set  $A$
- – Set  $B$

- 1 Venn diagrams are often drawn to abstractly illustrate **relations between multiple sets**. Elements are implicit/omitted (shown as dots only when an explicit element is needed)
- 2 *Example*: shaded area illustrates a set of elements that are in both sets  $A$  and  $B$  (i.e. *intersection* of two sets, see later).  
E.g consider  $A = \{a, b, c, f, z\}$  and  $B = \{c, d, e, f, x, y\}$ .

# Plan for Part I

## 1. Sets

1.1 Defining sets

1.2 Venn Diagram

**1.3 Set Equality**

1.4 Subsets

1.5 Venn Diagrams and Truth Sets

1.6 Set Cardinality

1.7 Power Sets

1.8 Cartesian Products

## 2. Set Operations

2.1 Boolean Algebra

2.2 Union

2.3 Intersection

2.4 Complement

2.5 Difference

2.6 The Cardinality of the Union of Two Sets

2.7 Set Identities

2.8 Generalized Unions and Intersections

# Set equality

**Definition:** Two sets are *equal* if and only if they have the same elements.

- 1 If  $A$  and  $B$  are sets, then  $A$  and  $B$  are equal iff:

$$\forall x \ (x \in A \leftrightarrow x \in B)$$

- 2 We write  $A = B$  if  $A$  and  $B$  are equal sets.

$$\{1, 3, 5\} = \{3, 5, 1\}$$

$$\{1, 5, 5, 5, 3, 3, 1\} = \{1, 3, 5\}$$

# Plan for Part I

## 1. Sets

1.1 Defining sets

1.2 Venn Diagram

1.3 Set Equality

### 1.4 Subsets

1.5 Venn Diagrams and Truth Sets

1.6 Set Cardinality

1.7 Power Sets

1.8 Cartesian Products

## 2. Set Operations

2.1 Boolean Algebra

2.2 Union

2.3 Intersection

2.4 Complement

2.5 Difference

2.6 The Cardinality of the Union of Two Sets

2.7 Set Identities

2.8 Generalized Unions and Intersections

# Subsets

**Definition:** The set  $A$  is a *subset* of  $B$ , if and only if every element of  $A$  is also an element of  $B$ .

- ① The notation  $A \subseteq B$  is used to indicate that  $A$  is a subset of the set  $B$
- ②  $A \subseteq B$  holds if and only if  $\forall x \ (x \in A \rightarrow x \in B)$  is true.
- ③ Observe that:
  - a Because  $a \in \emptyset$  is always false, for every set  $S$  we have:
$$\emptyset \subseteq S.$$
  - b Because  $a \in S \rightarrow a \in S$ , for every set  $S$ , we have:
$$S \subseteq S.$$

# Showing that a set is or is not a subset of another set

- ① **Showing that A is a Subset of B:** To show that  $A \subseteq B$ , show that if  $x$  belongs to  $A$ , then  $x$  also belongs to  $B$ .
- ② **Showing that A is not a Subset of B:** To show that  $A$  is not a subset of  $B$ , that is  $A \not\subseteq B$ , find an element  $x \in A$  with  $x \notin B$ . (Such an  $x$  is a counterexample to the claim that  $x \in A$  implies  $x \in B$ .)
- ③ **Examples:**
  - a The set of all computer science majors at your school is a subset of all students at your school.
  - b The set of integers with squares less than 100 is not a subset of the set of all non-negative integers.

$$-1^2 < 100.$$



## Another look at equality of sets

- ① Recall that two sets  $A$  and  $B$  are equal (denoted by  $A = B$ ) iff:

$$\forall x \quad (\underline{x \in A \leftrightarrow x \in B})$$

- ② That is, using logical equivalences we have that  $A = B$  iff:

$$\forall x \quad ((x \in A \rightarrow x \in B) \wedge (x \in B \rightarrow x \in A))$$

- ③ This is also equivalent to:

$$A \subseteq B \quad \text{and} \quad B \subseteq A$$


# Proper subsets

**Definition:** If  $A \subseteq B$ , but  $A \neq B$ , then we say  $A$  is a proper subset of  $B$ , denoted by  $A \subset B$ . If  $A \subset B$ , then the following is true:

$$\forall x \quad (x \in A \rightarrow x \in B) \quad \wedge \quad \exists x \quad (x \in B \wedge x \notin A)$$

**Example:**  $A = \{c, f, z\}$  and  $B = \{a, b, c, d, e, f, t, x, z\}$ .

Venn Diagram for a proper subset  $A \subset B$



# Plan for Part I

## 1. Sets


- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets**
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products


## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Venn diagrams and truth sets

Consider any predicate  $P(x)$  for elements  $x$  in  $U$  and its truth set  $P = \{x \mid P(x)\}$ .




 – truth set of  
 $P = \{x \mid P(x)\}$

that is, all elements  $x$  where  
 $P(x)$  is true

**Note that:**  $x \in P \equiv P(x)$

# Venn diagrams and logical connectives: negations

Consider any predicate  $P(x)$  for elements  $x$  in  $U$  and its truth set  $P = \{x \mid P(x)\}$ .



■ – truth set of  $\{x \mid \neg P(x)\}$


all elements  $x$  where  $\neg P(x)$  is true, i.e. where  $P(x)$  is false


Same as complement of set  $P$  (see section 2.4)

**Note that:**  $x \notin P \equiv \neg P(x)$

# Venn Diagrams and logical connectives: conjunctions

Consider two arbitrary predicates  $P(x)$  and  $Q(x)$  defined for elements  $x$  in  $U$  together with their corresponding truth sets  $P = \{x \mid P(x)\}$  and  $Q = \{x \mid Q(x)\}$ .




 – truth set of  $\{x \mid P(x) \wedge Q(x)\}$  that is, all elements  $x$  where both  $P(x)$  and  $Q(x)$  is true

Same as intersection of sets  $P$  and  $Q$  (see section 2.3)

# Venn Diagrams and logical connectives: disjunctions

Consider two arbitrary predicates  $P(x)$  and  $Q(x)$  defined for elements  $x$  in  $U$  together with their corresponding truth sets  $P = \{x \mid P(x)\}$  and  $Q = \{x \mid Q(x)\}$ .





■ – truth set  
 $\{x \mid P(x) \vee Q(x)\}$  that is, all  
elements  $x$  where  $P(x)$  or  
 $Q(x)$  is true

Same as *union* of sets  $P$  and  $Q$  (see section 2.2)


# Venn diagrams and logical connectives: implications

Consider two arbitrary predicates  $P(x)$  and  $Q(x)$  defined for elements  $x$  in  $U$  together with their corresponding truth sets  $P = \{x \mid P(x)\}$  and  $Q = \{x \mid Q(x)\}$ .



 – truth set  $\{x \mid P(x) \rightarrow Q(x)\}$

(all  $x$  where implication  $P(x) \rightarrow Q(x)$  is true)

 – set where implication  $P(x) \rightarrow Q(x)$  is false:

$$\begin{aligned}\{x \mid \neg(\neg P(x) \vee Q(x))\} &= \{x \mid P(x) \wedge \neg Q(x)\} \\ &= \{x \mid x \in P \wedge x \notin Q\}\end{aligned}$$

① Remember:

$$p \rightarrow q \equiv \neg p \vee q$$

② Thus, we have:


$$\{x \mid P(x) \rightarrow Q(x)\} = \{x \mid \neg P(x) \vee Q(x)\} = \{x \mid x \notin P \vee x \in Q\}$$





# Venn diagram and implication: special case

- 1 Assume that implication  $P(x) \rightarrow Q(x)$  is true for all  $x$ .
- 2 That is, assume  $\{x \mid P(x) \rightarrow Q(x)\} \equiv U$  is true.
- 3 Note that, from the definition of subsets:

$$\forall x(P(x) \rightarrow Q(x)) \quad \equiv \quad \forall x(x \in P \rightarrow x \in Q) \quad \equiv \quad P \subseteq Q$$



 — truth set  
 $\{x \mid P(x) \rightarrow Q(x)\}$   
(all  $x$  where implication  $P(x) \rightarrow Q(x)$  is true)

 — set  
where implication  
 $P(x) \rightarrow Q(x)$  is false:


Venn diagram for  $P \subseteq Q$  often shows  $P$  as a proper subset of  $Q$ , thus assuming  $P \neq Q$


$$\begin{aligned} \forall x(P(x) \rightarrow Q(x)) &\equiv \forall x(\neg P(x) \vee Q(x)) \\ &\equiv \neg \exists x(P(x) \wedge \neg Q(x)) \\ &\equiv \neg \exists x(x \in P \wedge x \notin Q) \end{aligned}$$


# Venn diagram and implication: special case

- 1 Assume that implication  $P(x) \rightarrow Q(x)$  is true for all  $x$ .
- 2 That is, assume  $\{x \mid P(x) \rightarrow Q(x)\} \equiv U$  is true.
- 3 Note that, from the definition of subsets:

$$\forall x(P(x) \rightarrow Q(x)) \quad \equiv \quad \forall x(x \in P \rightarrow x \in Q) \quad \equiv \quad P \subseteq Q$$



 – truth set  
 $\{x \mid P(x) \rightarrow Q(x)\}$   
(all  $x$  where implication  $P(x) \rightarrow Q(x)$  is true)

 – set  
where implication  
 $P(x) \rightarrow Q(x)$  is false:

Venn diagram for  $P \subseteq Q$  often shows  $P$  as a empty in this case:  $\{x \mid x \in P \wedge x \notin Q\} = \emptyset$   
proper subset of  $Q$ , thus assuming  $P \neq Q$

- 1 This gives intuitive interpretation for logical “implications”:
- 2 Proving theorems of the form  $\forall x(P(x) \rightarrow Q(x))$  is equivalent to proving the subset relationship for the truth sets  $P \subseteq Q$ .


# Venn diagram and logical connectives: biconditional


- 1 Similarly one can show that

$$\forall x P(x) \leftrightarrow Q(x) \equiv P = Q$$

- 2 that is,

$$\forall x (P(x) \rightarrow Q(x) \wedge Q(x) \rightarrow P(x)) \equiv P \subseteq Q \wedge Q \subseteq P.$$



 — truth set of  
 $\{x \mid P(x) \leftrightarrow Q(x)\}$

assuming  $\forall x (P(x) \leftrightarrow Q(x))$  is true

- 1 This gives intuitive interpretation for “biconditional”:
- 2 Proving theorems of the form  $\forall x (P(x) \leftrightarrow Q(x))$  is equivalent to proving the subset relationship for the truth sets  $P = Q$ .

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality**
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Set cardinality

**Definition:** If there are exactly  $n$  (distinct) elements in  $S$  where  $n$  is a non-negative integer, we say that  $S$  is *finite*. Otherwise, the set  $S$  is said *infinite*.

**Definition:** The *cardinality* of a finite set  $A$ , denoted by  $|A|$ , is the number of (distinct) elements of  $A$ .

## Examples:

- ①  $|\emptyset| = 0$
- ② Let  $S$  be the letters of the English alphabet. Then  $|S| = 26$
- ③  $|\{1,2,3\}| = 3$
- ④  $|\{\emptyset\}| = 1$
- ⑤ The set of integers is infinite.

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets**
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Power sets

**Definition:** The set of all subsets of a set  $A$ , denoted  $\mathcal{P}(A)$ , is called the *power set* of  $A$ .

**Example:** If  $A = \{a, b\}$  then:

$$\mathcal{P}(A) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$$

- 1 If a set has  $n$  elements, then the cardinality of the power set is  $2^n$ .
- 2 In Chapters 5 and 6, we will discuss different ways to show this.

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products**

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections



# Tuples

- ① The *ordered  $n$ -tuple*  $(a_1, a_2, \dots, a_n)$  is the ordered collection that has  $a_1$  as its first element and  $a_2$  as its second element and so on until  $a_n$  as its last element.
- ② Two  $n$ -tuples are equal if and only if their corresponding elements are equal.
- ③ 2-tuples are called *ordered pairs*, e.g.  $(a_1, a_2)$
- ④ The ordered pairs  $(a, b)$  and  $(c, d)$  are equal if and only if  $a = c$  and  $b = d$ .

# Cartesian Product



René

Descartes

(1596-1650)

**Definition:** The *Cartesian Product* of two sets  $A$  and  $B$ , denoted by  $A \times B$  is the set of ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$ .

**Example:**

$$A = \{a, b\}, B = \{1, 2, 3\}$$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

**Definition:** A subset  $R$  of the Cartesian product  $A \times B$  is called a *relation* from the set  $A$  to the set  $B$ . Relations will be covered in depth in Chapter 9.

# Cartesian product

**Definition:** The Cartesian products of the sets  $A_1, A_2, \dots, A_n$  denoted by  $A_1 \times A_2 \times \dots \times A_n$  is the set of ordered n-tuples  $(a_1, a_2, \dots, a_n)$  where  $a_i$  belongs to  $A_i$  for  $i = 1, 2, \dots, n$ .

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) \mid a_i \in A_i \text{ for } i = 1, 2, \dots, n\}$$

**Question:** What is  $A \times B \times C$  where  $A = \{0,1\}$ ,  $B = \{1,2\}$  and  $C = \{0,1,2\}$ ?

**Solution:**  $A \times B \times C = \{(0,1,0), (0,1,1), (0,1,2), (0,2,0), (0,2,1), (0,2,2), (1,1,0), (1,1,1), (1,1,2), (1,2,0), (1,2,1), (1,2,2)\}$

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Boolean Algebra

- ① Propositional calculus and set theory are both instances of an algebraic system called a *Boolean Algebra*. This is discussed in CS2209.
- ② The operators in set theory are analogous to corresponding operators in propositional calculus.
- ③ As always there must be a universal set  $U$ .
- ④ All sets  $A, B, \dots$  shown in the next slides are assumed to be subsets of  $U$ .

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union**
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections


# Union

- ① **Definition:** Let  $A$  and  $B$  be sets. The *union* of the sets  $A$  and  $B$ , denoted by  $A \cup B$ , is the set:

$$\{x \mid x \in A \vee x \in B\}$$

- ② **Example:** What is  $\{1,2,3\} \cup \{3, 4, 5\}$ ?

**Solution:**  $\{1,2,3,4,5\}$



*Union* is analogous to *disjunction*, see earlier slides.



# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection**
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Intersection

- ① **Definition:** The *intersection* of sets  $A$  and  $B$ , denoted by  $A \cap B$ , is:

$$\{x \mid x \in A \wedge x \in B\}$$

- ② If the intersection is empty, then  $A$  and  $B$  are said to be *disjoint*.


- ① **Example:** What is  $\{1,2,3\} \cap \{3,4,5\}$  ?

**Solution:**  $\{3\}$

- ① **Example:** What is  $\{1,2,3\} \cap \{4,5,6\}$  ?

**Solution:**  $\emptyset$

Venn Diagram for  $A \cap B$



*Intersection* is analogous to *conjunction*, see earlier slides.

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement**
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Complement

**Definition:** If  $A$  is a set, then the complement of the  $A$  (with respect to  $U$ ), denoted by  $\bar{A}$  is the set:


$$\bar{A} = \{x \in U \mid x \notin A\}$$

(The complement of  $A$  is sometimes denoted by  $A^c$ .)

**Example:** If  $U$  is the positive integers less than 100, what is the complement of  $\{x \mid x > 70\}$ ?

**Solution:**  $\{x \mid x \leq 70\}$

Venn Diagram for complement



Complement is analogous to [negation](#), see earlier.

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations


- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference**
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Difference

**Definition:** Let  $A$  and  $B$  be sets. The *difference* of  $A$  and  $B$ , denoted by  $A - B$ , is the set containing the elements of  $A$  that are not in  $B$ . The difference of  $A$  and  $B$  is also called the complement of  $B$  with respect to  $A$ .

$$A - B = \{x \mid x \in A \wedge x \notin B\} = A \cap \overline{B}$$

Venn Diagram for  $A - B$



**Note:**  $\overline{A} = U - A$

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products


## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# The cardinality of the union of two sets

## 1 Inclusion-Exclusion:

$$|A \cup B| = |A| + |B| - |A \cap B|$$



Venn Diagram for  $A$ ,  $B$ ,  $A \cap B$ ,  $A \cup B$

## 1 Example:

- a Let  $A$  be the math majors in your class and  $B$  be the CS majors in your class.
- b To count the number of students in your class who are either math majors or CS majors, add the number of math majors and the number of CS majors, and subtract the number of joint CS/math majors.
- c We will return to this principle in Chapter 6 and Chapter 8, where we will derive a formula for the cardinality of the union of  $n$  sets, where  $n$  is a positive integer.



# Review questions

**Example:** Given  $U = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ ,  
 $A = \{1, 2, 3, 4, 5\}$ ,  $B = \{4, 5, 6, 7, 8\}$  solve the following:

①  $A \cup B$

**Solution:**

$$\{1, 2, 3, 4, 5, 6, 7, 8\}$$

②  $A \cap B$

**Solution:**  $\{4, 5\}$

③  $\overline{A}$

**Solution:**

$$\{0, 6, 7, 8, 9, 10\}$$

④  $\overline{B}$

**Solution:**

$$\{0, 1, 2, 3, 9, 10\}$$

⑤  $A - B$

**Solution:**  $\{1, 2, 3\}$

⑥  $B - A$

**Solution:**  $\{6, 7, 8\}$

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Set identities

## ① Identity laws

$$A \cup \emptyset = A$$

$$A \cap U = A$$

## ② Domination laws

$$A \cup U = U$$

$$A \cap \emptyset = \emptyset$$

## ③ Idempotent laws

$$A \cup A = A$$

$$A \cap A = A$$

## ④ Complementation law

$$\overline{(\overline{A})} = A$$

*Continued on next slide ↪*

# Set identities

## ① Commutative laws

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

## ② Associative laws

$$A \cup (B \cup C) = (A \cup B) \cup C \quad A \cap (B \cap C) = (A \cap B) \cap C$$

## ③ Distributive laws

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

*Continued on next slide ↪*

# Set identities

## ① De Morgan's laws

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

## ② Absorption laws

$$A \cup (A \cap B) = A$$

$$A \cap (A \cup B) = A$$

## ③ Complement laws

$$A \cup \overline{A} = U$$

$$A \cap \overline{A} = \emptyset$$

# Proving set identities

- ① Different ways to prove set identities:
  - a Prove that each set (i.e. each side of the identity) is a subset of the other.
  - b Use set builder notation and propositional logic.
  - c Membership tables

(to be explained)

# Proof of second De Morgan law

**Example:** Prove that  $\overline{A \cap B} = \overline{A} \cup \overline{B}$

**Solution:** We prove this identity by showing that:

$$① \quad \overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$$

$$② \quad \overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$$

*Continued on next slide ↪*

# Proof of second De Morgan law

These steps show that:  $\overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$

$$x \in \overline{A \cap B}$$

by assumption

$$x \notin A \cap B$$

definition of complement

$$\neg((x \in A) \wedge (x \in B))$$

definition of intersection

$$\neg(x \in A) \vee \neg(x \in B)$$

De Morgan's 1<sup>st</sup> Law

$$(x \notin A) \vee (x \notin B)$$

definition of negation

$$(x \in \overline{A}) \vee (x \in \overline{B})$$

definition of complement

$$x \in \overline{A} \cup \overline{B}$$

definition of union

*Continued on next slide ↪*



# Proof of second De Morgan law

These steps show that:  $\overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$

$$x \in \overline{A} \cup \overline{B}$$

by assumption

$$(x \in \overline{A}) \vee (x \in \overline{B})$$

definition of union

$$(x \notin A) \vee (x \notin B)$$

definition of complement

$$\neg(x \in A) \vee \neg(x \in B)$$

definition of negation

$$\neg((x \in A) \wedge (x \in B))$$

De Morgan's 1<sup>st</sup> Law

$$\neg x \in (A \cap B)$$

definition of intersection

$$x \in \overline{A \cap B}$$

definition of complement



## Set-builder notation: second De Morgan law

$\overline{A \cap B}$	$= \{x \mid x \notin A \cap B\}$	by definition of complement
	$= \{x \mid \neg x \in (A \cap B)\}$	by definition of 'not in'
	$= \{x \mid \neg((x \in A) \wedge (x \in B))\}$	by definition of intersection
	$= \{x \mid \neg(x \in A) \vee \neg(x \in B)\}$	by De Morgan's 1 <sup>st</sup> Law
	$= \{x \mid (x \notin A) \vee (x \notin B)\}$	by definition of 'not'
	$= \{x \mid (x \in \overline{A}) \vee (x \in \overline{B})\}$	by definition of complement
	$= \{x \mid x \in \overline{A} \cup \overline{B}\}$	by definition of union
	$= \overline{A} \cup \overline{B}$	by definition of notation



# Membership table

**Example:** Construct a membership table to show that the distributive law holds:

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

**Solution:**

$A$	$B$	$C$	$B \cap C$	$A \cup (B \cap C)$	$A \cup B$	$A \cup C$	$(A \cup B) \cap (A \cup C)$
1	1	1	1	1	1	1	1
1	1	0	0	1	1	1	1
1	0	1	0	1	1	1	1
1	0	0	0	1	1	1	1
0	1	1	1	1	1	1	1
0	1	0	0	0	1	0	0
0	0	1	0	0	0	1	0
0	0	0	0	0	0	0	0

# Plan for Part I

## 1. Sets

- 1.1 Defining sets
- 1.2 Venn Diagram
- 1.3 Set Equality
- 1.4 Subsets
- 1.5 Venn Diagrams and Truth Sets
- 1.6 Set Cardinality
- 1.7 Power Sets
- 1.8 Cartesian Products

## 2. Set Operations

- 2.1 Boolean Algebra
- 2.2 Union
- 2.3 Intersection
- 2.4 Complement
- 2.5 Difference
- 2.6 The Cardinality of the Union of Two Sets
- 2.7 Set Identities
- 2.8 Generalized Unions and Intersections

# Generalized unions and intersections

- ① Let  $A_1, A_2, \dots, A_n$  be an indexed collection of sets.

We define:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n$$

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \dots \cap A_n$$

These are well defined, since union and intersection are associative.

- ② *Example:* for  $(i = 1, 2, \dots)$  let  $A_i = \{i, i + 1, i + 2, \dots\}$ . Then,

$$\bigcup_{i=1}^n A_i = \bigcup_{i=1}^n \{i, i + 1, i + 2, \dots\} = \{1, 2, 3, \dots\}$$

$$\bigcap_{i=1}^n A_i = \bigcap_{i=1}^n \{i, i + 1, i + 2, \dots\} = \{n, n + 1, n + 2, \dots\} = A_n$$