


Part 2

CHAPTER 2

Computer Arithmetic and Digital Logic



1

These slides are provided with permission from the copyright for CS2208 use only. The slides must not be reproduced or provided to anyone outside the class.

All downloaded copies of the slides are for personal use only.

Students must destroy these copies within 30 days after receiving the course's final assessment.

Signed Integers

- Signed numbers can be represented in many ways, including
 - *sign and magnitude*,
 - *biased representation*, and
 - *two's complement*.

Sign and Magnitude Representation

- ❑ An n -bit word has 2^n possible values
 - for example, an eight-bit word can represent 256 different values.
- ❑ One way of representing a negative number is to take the *most-significant bit* and *reserve it* to indicate *the sign of the number*.
 - 0 represents **positive** numbers and
 - 1 represents **negative** numbers.
- ❑ *In an eight-bit word, using **sign and magnitude representation***
 - How many positive numbers can be represented?* $2^7 = 128$.
 - How many negative numbers can be represented?* 128 positive #s, 128 negative #s

issue:
 +0, -0
 2 representations for the same #
- ❑ The value of a *sign and magnitude number* can be expressed as $(-1)^S \times M$, where **S** is the **sign bit** and **M** is the **magnitude** of the number.
 - If **S** = 0, $(-1)^0 = +1$ and the number is positive.
 - If **S** = 1, $(-1)^1 = -1$ and the number is negative.
- ❑ for example, in 8 bits we can **interpret** the numbers 00001101_2 and 10001101_2 as $+13_{10}$ and -13_{10} , respectively.
- ❑ **Sign and magnitude** representation is
 - **not generally used in integer arithmetic,**
 - however, it is **used in floating-point arithmetic.**

Biased Representation

- ❑ Assume that we have values ranging from -7 to $+7$ \rightarrow 15 integers.
- ❑ To go around the negative sign, we can *shift the scale* (by adding a bias = 7 to all values) to have only non-negative values.
 - The original number = -7 \rightarrow The biased number = 0
 - The original number = -6 \rightarrow The biased number = 1
 - The original number = 0 \rightarrow The biased number = 7
 - The original number = $+6$ \rightarrow The biased number = 13
 - The original number = $+7$ \rightarrow The biased number = 14
- ❑ To convert a biased number to its original value, you need to *shift back the scale* (by subtracting the bias from the biased number).
 - The biased number = 0 \rightarrow The original number = -7
 - The biased number = 1 \rightarrow The original number = -6
 - The biased number = 7 \rightarrow The original number = 0
 - The biased number = 13 \rightarrow The original number = $+6$
 - The biased number = 14 \rightarrow The original number = $+7$
- ❑ In this representation, *biased numbers are unsigned integers*; yet, they *represent both positive and negative values*.
- ❑ The biased representation is also called excess- K , where K is a pre-specified biasing value

Excess-7
code

\hookrightarrow excess -7 in this case

Complementary Arithmetic

- A number and its **complement** add up to a constant;
 - In *n-bit decimal ten's complement* (*radix complement*) arithmetic, if P is a number, then its *ten's* complement is Q , where $P + Q = 10^n$.
 - the *ten's* complement of 12 is 88 because $12 + 88 = 100$.
 - the *ten's* complement of 88 is 12 because $88 + 12 = 100$.
 - In *n-bit binary two's complement* (*radix complement*) arithmetic, if P is a number, then its *two's* complement is Q , where $P + Q = 2^n$.
 - the *two's* complement of 101_2 is 011_2 because $101_2 + 011_2 = 1000_2$.
 - the *two's* complement of 011_2 is 101_2 because $011_2 + 101_2 = 1000_2$.

Two's Complement Arithmetic

□ In binary arithmetic, the *two's complement* of a number is formed by

Method
①

- *Subtracting the number from 2^n .* ↖ 8 digits

The *two's* complement of 01100101_2 is
 $100000000_2 - 01100101_2 = 10011011_2$
↖ 9 digits

- ② ○ *Flipping (inverting) all the bits of the number and adding 1.*

The *two's* complement of 01100101_2 is
 $10011010_2 + 1_2 = 10011011_2$.

→ change 0 to 1 & vice versa,
add 1

- ③ ○ *Processing all the bits of the number from the least significant bit (LSB) towards the most significant bit (MSB)*

- copying all the zeros until the first 1 is reached,
- copying that 1,
- flipping (inverting) all the remaining bits.

seems easiest.

The *two's* complement of 01100100_2 is 10011100_2 .

The *two's* complement of 01100101_2 is 10011011_2 .

Two's Complement Arithmetic

*the complement of a value
is the negative of that value*

- ❑ The *two's* complement of a *positive* number is the corresponding *negative* value of the *positive* number when it is represented in *two's* complement.
- ❑ The *two's* complement of a *negative* number is the corresponding *positive* value of the *negative* number when it is represented in *two's* complement.
- ❑ We are interested in *complementary* arithmetic because *subtracting* a number is the same as *adding its complement*.
 - To subtract 001100101_2 from a binary number, we just need to add its *two's* complement (i.e., 110011011_2) to the number.

Two's Complement Arithmetic

- The *two's* complement of an n -bit binary value, N , is defined as $2^n - N$.
- If $N = 5_{10} = 00000101_2$ (8-bit arithmetic), the *two's* complement of N is given by $2^8 - 00000101_2 = 100000000_2 - 00000101_2 = \underbrace{11111011}_{123}_2$.
- 11111011_2 represents -00000101_2 (i.e., -5_{10}) or -123_{10} depending only on whether we interpret 11111011_2 as a *two's complement integer* or as a *sign-magnitude integer*, respectively.

(-123)

This is sign-magnitude not unsigned

This is
-123 not +123

Two's Complement Arithmetic

- This example demonstrates 8-bit *two's* complement arithmetic.

We begin by writing down the representations of +5, -5, +7 and -7.

$$+5_{10} = 00000101_2 \quad -5_{10} = 11111011_2 \quad +7_{10} = 00000111_2 \quad -7_{10} = 11111001_2$$

- Now $7_{10} - 5_{10}$ can be calculated as $7_{10} + (-5_{10})$ by adding the binary value for 7_{10} to the *two's* complement of 5_{10} .

| | |
|------------------|----------|
| 11111111 | |
| 00000111 | 7 |
| +11111011 | -5 |
| <u>100000010</u> | <u>2</u> |

→ 8-bit system.
ignore carry out

The result is correct if the left-hand carry-out is ignored.

Two's Complement Arithmetic

□ Now consider the addition of -7_{10} to $+5_{10}$, i.e., $+5_{10} + (-7_{10})$

$$\begin{array}{r}
 1 \\
 \textcolor{blue}{0}0000101 \qquad 5 \\
 + \textcolor{blue}{1}1111001 \qquad -7 \\
 \hline
 \textcolor{blue}{1}1111110 \qquad -2
 \end{array}$$

The result is $\textcolor{blue}{1}1111110_2$ (the carry-out bit is $\textcolor{red}{0}$).

The expected answer is -2_{10} ;

that is, $2^8 - 2_{10} = 100000000_2 - \textcolor{blue}{0}0000010_2 = \textcolor{blue}{1}1111110_2$.

Two's Complement Arithmetic

- ❑ *Two's* complement arithmetic is not magic.
- ❑ Consider the calculation $Z = X - Y$ in n -bit arithmetic
 - we do it by *adding* the *two's* complement of Y to X .
- ❑ The *two's* complement of Y is defined as $2^n - Y$.

We get

$$Z = X - Y = X + (2^n - Y) = 2^n + (X - Y).$$

- ❑ This is
 - *the desired result*, $X - Y$
 - If the result of $X - Y$ is negative, the *two's* complement will be automatically calculated, i.e., $2^n + (X - Y)$, where $(X - Y)$ is negative.
 - Otherwise, we discard the *unwanted carry-out digit* (i.e., 2^n) at the leftmost position

Two's Complement Arithmetic

Time examples using radix complement

In the 24H system (*i.e.*, $\text{radix} = 24$), assume that

- ❑ the time right now is 2 am and we want to subtract 5 hours from 2 am
 - The radix complement of 5 is $24 - 5 = 19$
 - To subtract 5 from 2 in the 24H system, it is equivalent to adding the 2 to the complement of 5, *i.e.*, $2 + 19 = 21$ (*correct answer*)
 - The 21 can also be read as -3 in the radix complement system ^(9 pm)
- ❑ the time right now is 5 am and we want to subtract 2 hours from 5 am
 - The radix complement of 2 is $24 - 2 = 22$
 - To subtract 2 from 5 in the 24H system, it is equivalent to adding the 5 to the complement of 2, *i.e.*, $5 + 22 = 27$
 - As all the time values must be between 0 and 23, we need to subtract a full 24H from the result, *i.e.*, $27 - 24 = 3$ (*correct answer*)

Two's Complement Arithmetic

Let $X = 9_{10} = 00001001_2$ and $Y = 6_{10} = 00000110_2$

$$-X = -9_{10} = 100000000_2 - 00001001_2 = 11110111_2$$

$$-Y = -6_{10} = 100000000_2 - 00000110_2 = 11111010_2$$

$$\begin{array}{r} 1. \quad +X \quad +9 \quad 00001001 \\ \quad +Y \quad +6 \quad +00000110 \\ \hline \quad \quad +15 \quad 00001111 \end{array}$$

$$\begin{array}{r} 2. \quad +X \quad +9 \quad 00001001 \\ \quad -Y \quad -6 \quad +11111010 \\ \hline \quad \quad +3 \quad 100000011 \end{array}$$

$$\begin{array}{r} 3. \quad -X \quad -9 \quad 11110111 \\ \quad +Y \quad +6 \quad +00000110 \\ \hline \quad \quad -3 \quad 11111101 \end{array}$$

$$\begin{array}{r} 4. \quad -X \quad -9 \quad 11110111 \\ \quad -Y \quad -6 \quad +11111010 \\ \hline \quad \quad -15 \quad 111110001 \end{array}$$

11111101 is a negative *two's* complement number

The two's complement of **11111101** is **00000011**₂

Hence, **11111101** = **-00000011**₂ = **-3**₁₀

11110001 is a negative *two's* complement number

The two's complement of **11110001** is **00001111**₂

Hence, **11110001** = **-00001111**₂ = **-15**₁₀

All four examples give the result we would expect when the result is interpreted as a *two's* complement number.

Properties of Two's complement Numbers

- ❑ The *two's* complement system is a true complement system in that $+X + (-X) = 0$.
- ❑ There is *one* unique zero 00...0.
- ❑ This is *not the case with the signed magnitude* number system
- ❑ The *most-significant bit* of a *two's* complement number is a *sign bit*.
 - The number is *positive* if the most-significant bit is *0*, and
 - *negative* if it is *1*.

This is the same as the signed magnitude number system
- ❑ The range of an *n-bit two's* complement number is from -2^{n-1} to $+2^{n-1} - 1$.
 - For $n = 4$, the range is -8 to $+7$.

The total number of different values is $2^n = 16$

 - 8 negative,
 - zero and
 - 7 positive.
 - For $n = 8$, the range is -128 to $+127$.

The total number of different values is $2^n = 256$

 - 128 negative,
 - zero and
 - 127 positive.

❑ *How about the signed magnitude number system?*

Arithmetic Overflow

- ❑ The *range* of *two's* complement numbers in *n-bits* is from -2^{n-1} to $+2^{n-1} - 1$.
- ❑ What happens if we *violate* this rule by carrying out an operation whose *result falls outside the range* of values that can be represented by *two's* complement numbers?
 - In a *five-bit* representation, the range of *valid two's complement numbers* is -16 to $+15$. <-16, >15 : overflow

Case 1

$$\begin{array}{r}
 5 = 00101 \\
 +7 = 00111 \\
 \hline
 12 = 01100 = 12_{10}
 \end{array}$$

Case 2

$$\begin{array}{r}
 12 = 01100 \\
 +13 = 01101 \\
 \hline
 25 = 11001 = -7_{10} \text{ (as a two's complement value)}
 \end{array}$$

That's why sometimes when you add 2 big #s (of same sign), you get a negative.

In *Case 1* we get the *expected* answer of $+12_{10}$, but

In *Case 2* we get a *negative* result because the *sign bit is 1*.

- If the answer were regarded as an *unsigned binary number*, it would be $+25$, which is, of course, the correct answer.
- However, as the *two's* complement system has been chosen to represent signed numbers, all answers must be interpreted in this light.

Arithmetic Overflow

- If we add together two negative numbers whose total is less than -16 , we will go out of range.

Case 3

$$\begin{array}{rcl} -9 & = & 10111 \\ -12 & = & +10100 \\ \hline -21 & & 101011 = +11_{10} \text{ (as a } \textit{two's} \text{ complement value)} \end{array}$$

In *Case 3* we get a **positive** result because the *sign bit is 0*.

Arithmetic Overflow

- ❑ The last two examples demonstrate *arithmetic overflow*
- ❑ *Arithmetic overflow* occurs during a *two's complement addition* when
 - the result of *adding two positive* numbers *yields a negative* result,
 - or
 - the result of *adding two negative* numbers *yields a positive* result.
- ❑ *If the sign bits of A and B are the same but the sign bit of the result is different, it means arithmetic overflow has occurred.*
- ❑ *Overflow never occurs when adding two numbers with opposite signs*
- ❑ If a_{n-1} is the sign bit of A,
 b_{n-1} is the sign bit of B, and
 s_{n-1} is the sign bit of the sum of A and B,
 then

overflow is defined by the *logical expression* as follow:

$$V = (a_{n-1})^c \cdot (b_{n-1})^c \cdot s_{n-1} + a_{n-1} \cdot b_{n-1} \cdot (s_{n-1})^c$$

where $()^c$ is the complement binary operation

← 2209.

Arithmetic Overflow

- In practice, the overflow can be detected from the *carry bits* *in to* (i.e., C_{in}) and *out of* (i.e., C_{out}) the *most-significant bit* of an adder;

overflow occurs when $C_{in} \neq C_{out}$

same: no overflow

same: no o.f.

8-bit examples in slide#30

5-bit examples in slides#32,33

35

Overflow occurred

Overflow occurred

$$\begin{array}{r}
 C_{out} \quad C_{in} \\
 00000000 \\
 +X \quad +9 \quad 00001001 \\
 +Y \quad +6 \quad +0000110 \\
 \hline
 +15 \quad 00001111 \\
 C_{out} \quad C_{in} \\
 00000110
 \end{array}$$

$$\begin{array}{r}
 C_{out} \quad C_{in} \\
 00000110 \\
 -X \quad -9 \quad 11110111 \\
 +Y \quad +6 \quad +0000110 \\
 \hline
 -3 \quad 11111101 \\
 C_{out} \quad C_{in} \\
 00111
 \end{array}$$

$$\begin{array}{r}
 5 = 00101 \\
 +7 = 00111 \\
 \hline
 12 \quad 01100 = +12_{10}
 \end{array}$$

$$\begin{array}{r}
 C_{out} \quad C_{in} \\
 01100 \\
 12 = 01100 \\
 +13 = 01101 \\
 \hline
 25 \quad 11001 = -7_{10}
 \end{array}$$

$$\begin{array}{r}
 C_{out} \quad C_{in} \\
 10100 \\
 -9 = 10111 \\
 -12 = +10100 \\
 \hline
 -21 \quad 101011 = +11_{10}
 \end{array}$$


Shifting Operations

- ❑ In a shift operation, the *bits of a word* are *shifted* one or more places to the *left* or *right*.
- ❑ The *arithmetic* shift is just one type of shift operation.
- ❑ In Chapter 3, we will cover the other types of shift operations.

Shifting Operations (left)

- Figure 2.2(a) describes the arithmetic *shift left*.
 - A *zero* enters into the vacated *least-significant bit* position and
 - the bit *shifted out* of the *most-significant bit* position is recorded in the computer's *carry flag*.
 - In *two's* complement and unsigned numbers:
 - Arithmetic *shift left* means *multiplying* by 2.
- If 00100111_2 (39_{10}) is shifted one place left it becomes 01001110_2 (78_{10}).
- If 11100011_2 (-29_{10}) is shifted one place left it becomes 11000110_2 (-58_{10}).

FIGURE 2.2 Arithmetic shift operations



(a) Arithmetic shift left:


A zero enters the least-significant bit position and the most-significant bit is copied into the carry flag. For example: 11000101 becomes 10001010 after shifting one place left.

Shifting Operations (**right**)

- ❑ Figure 2.2(b) describes the arithmetic *shift right*.
 - the **sign bit** (the **MSB**) is *replicated* into the vacated *most-significant bit* position and
 - the bit *shifted out* of the *least-significant bit* position is recorded in the computer's *carry flag*.
- In *two's* complement and unsigned numbers:
 - Arithmetic *shift right* means *dividing* by 2.
- ❑ Shifting 00001100_2 $(12)_{10}$ one place right produces 00000110_2 $(6)_{10}$.
- ❑ Shifting 00001101_2 $(13)_{10}$ one place right produces 00000110_2 $(6)_{10}$ as well.
- ❑ Shifting 11100010_2 $(-30)_{10}$ one place right produces 11110001_2 $(-15)_{10}$.
- ❑ Shifting 11100011_2 (-29) one place right produces 11110001_2 (-15) as well.

FIGURE 2.2

Arithmetic shift operations



(b) Arithmetic shift right:

A copy of the most-significant bit enters the most-significant bit position. All other bits are shifted one place right. The least-significant bit is copied into the carry flag.

For example; 00100101 becomes 00010010 after shifting one place right, and 11100101 becomes 11110010 after shifting one place right.

© Cengage Learning 2014

Binary Multiplication

- Multiplication can be implemented using arithmetic *shift left* and *addition* operations to add up the partial products as they are formed.

| Multiplicand | Multiplier | Step | Partial products | | | | | | | | | | | | | |
|---------------|-------------------|------|------------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| 01101001 | 0100100 1 | 1 | | | | | | | | | | | | | | |
| 01101001 | 0100100 1 | 2 | | | | | | | | | | | | | | |
| 01101001 | 01001 0 01 | 3 | | | | | | | | | | | | | | |
| 01101001 | 0100 1 001 | 4 | | | | | | | | | | | | | | |
| 01101001 | 010 0 1001 | 5 | | | | | | | | | | | | | | |
| 01101001 | 01 0 01001 | 6 | | | | | | | | | | | | | | |
| 01101001 | 0 1 001001 | 7 | | | | | | | | | | | | | | |
| 01101001 | 0 1001001 | 8 | | | | | | | | | | | | | | |
| Result | | | 0 | 0 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | 1 |