

These slides are being provided with permission from the copyright for CS2208 use only. The slides must not be reproduced or provided to anyone outside of the class.

All download copies of the slides and/or lecture recordings are for personal use only. Students must destroy these copies within 30 days after receipt of final course evaluations.

# Tutorial 01: Number Systems

*Computer Science Department*

*CS2208: Introduction to Computer Organization and Architecture*

*Fall 2022-2023*

*Instructor: Mahmoud R. El-Sakka*

*Office: MC-419*

*Email: [elsakka@csd.uwo.ca](mailto:elsakka@csd.uwo.ca)*

*Phone: 519-661-2111 x86996*

# Number Systems

- In positional notation number systems
  - Numbers are *represented* (*encoded*) using digits
  - Each digit has a *value* and a *place*
  - Each *place* has a *weight*
    - for example, in decimal, we have the *ones place*, *tens place*, *hundreds place*, ...

# Number Systems

- A radix or base is
  - the number of unique digits, *including zero*, used to represent numbers in a positional numeral system.
- With the use of a *radix point*, the notation can be extended to include *fractions* and *real numbers*.

# Number Systems

## ■ Examples of positional numeral systems

- Decimal is base-10 → {0, 1, 2, 3, 4, 5, 6, 7, 8, and 9}
- Binary is base-2 → {0, and 1}
- Quaternary is base-4 → {0, 1, 2, and 3}
- Octal is base-8 → {0, 1, 2, 3, 4, 5, 6, and 7}
- Hexadecimal is base-16 → {0, 1, 2, 3, 4, 5, 6, 7, 8, 9, **A**, **B**, **C**, **D**, **E** and **F**}
- Trinary is base-3 → {0, 1, and 2}
- Quinary is base-5 → {0, 1, 2, 3, and 4}
- Senary is base-6 → {0, 1, 2, 3, 4, and 5}
- Septenary is base-7 → {0, 1, 2, 3, 4, 5, and 6}
- Nonary is base-9 → {0, 1, 2, 3, 4, 5, 6, 7, and 8}
- Duodecimal is base-12 → {0, 1, 2, 3, 4, 5, 6, 7, 8, 9, **A**, and **B**}
- Sexagesimal is base-60 → {0, 1, 2, 3, 4, 5, ..., **58** and **59**}

You need to know how  
to convert values from  
one system to  
the other

# Conversion from any other base system to decimal

□ If the original number in base  $b$  is:

$$(a_{n-1} a_{n-2} \dots a_i \dots a_1 a_0 \cdot a_{-1} a_{-2} \dots a_{-m})_b$$

□ The *decimal* value of this number is defined as

$$N_{10} = (a_{n-1}b^{n-1} + \dots + a_i b^i + \dots + a_1 b^1 + a_0 b^0 + a_{-1} b^{-1} + a_{-2} b^{-2} + \dots + a_{-m} b^{-m})_{10}$$

# Conversion from any other base system to decimal

■ Example 1: Convert  $2E8_{16}$  to *decimal*

$$\begin{aligned}
 2E8_{16} &= 2 \times 16^2 + E \times 16^1 + 8 \times 16^0 \\
 &= 2 \times 256 + 14 \times 16 + 8 \times 1 \\
 &= 512 + 224 + 8 \\
 &= 744_{10}
 \end{aligned}$$

This question can be asked  
as follow:  
Convert the hexadecimal  
value  $2E8$  to *decimal*

Calculators are not allowed during exams.  
You need to improve your mental math skills.

During exams, calculations will be simplified.

Yet, when you answer the assignment/quiz questions, you  
may want to use calculators, as simplifying the  
calculations are not considered in the assignment/quiz.

|        |
|--------|
| 10 = A |
| 11 = B |
| 12 = C |
| 13 = D |
| 14 = E |
| 15 = F |

# Conversion from any other base system to decimal

■ Example 2: Convert  $361_8$  to *decimal*

$$\begin{aligned} 361_8 &= 3 \times 8^2 + 6 \times 8^1 + 1 \times 8^0 \\ &= 3 \times 64 + 6 \times 8 + 1 \times 1 \\ &= 192 + 48 + 1 \\ &= 241_{10} \end{aligned}$$

This question can be asked  
as follow:  
Convert the octal value 361  
to *decimal*

# Conversion from any other base system to decimal

■ Example 3: Convert  $0.361_8$  to *decimal*

$$\begin{aligned}
 0.361_8 &= 3 \times 8^{-1} + 6 \times 8^{-2} + 1 \times 8^{-3} \\
 &= 3 \times 0.125 + 6 \times 0.015625 + 1 \times 0.001953125 \\
 &= 0.375 + 0.09375 + 0.001953125 \\
 &= 0.470703125_{10}
 \end{aligned}$$

Another method:

$$\begin{aligned}
 0.361_8 &= 361_8 / 1000_8 \\
 &= (3 \times 8^2 + 6 \times 8^1 + 1 \times 8^0) / (1 \times 8^3) \\
 &= (3 \times 64 + 6 \times 8 + 1 \times 1) / (1 \times 512) \\
 &= (192 + 48 + 1) / (512) \\
 &= 241 / 512 \\
 &= 0.470703125_{10}
 \end{aligned}$$



# Conversion from any other base system to decimal

## ■ Example 4: $12.112_3$ to *decimal*

$$12.112_3$$

$$= 1 \times 3^1 + 2 \times 3^0 + 1 \times 3^{-1} + 1 \times 3^{-2} + 2 \times 3^{-3}$$

$$= 1 \times 3 + 2 \times 1 + 1 \times 0.33333 + 1 \times 0.11111 + 2 \times 0.03703$$

$$= 3 + 2 + 0.33333 + 0.11111 + 0.07406 = 5.5185_{10}$$

## Another method:

$$12.112_3 = 12112_3 / 1000_3$$

$$= (1 \times 3^4 + 2 \times 3^3 + 1 \times 3^2 + 1 \times 3^1 + 2 \times 3^0) / (1 \times 3^3)$$

$$= (81 + 54 + 9 + 3 + 2) / 27$$

$$= 149 / 27 = 5.5185_{10}$$

# Conversion from decimal to any other base system

## ■ Division Method (for integer numbers)

- Initialize the **quotient** by the value of the *decimal number*
- *Repeat*:
  - **Divide** the **quotient** from the previous stage **by the new base** to get
    - A new **quotient** (the whole number)
    - A *remainder*
  - The *remainder* here is *the next least significant digit* in the new number
- Until* the new **quotient** becomes 0.

# Conversion from decimal to any other base system

- Example 5: Convert  $14_{10}$  to binary

Binary means the new base is 2

- $14/2 = 7$       Remainder: 0 → This is the least significant binary digit  
Quotient =  $7 \neq 0$  → continue
- $7/2 = 3$       Remainder: 1 → This is the 2<sup>nd</sup> least significant binary digit  
Quotient =  $3 \neq 0$  → continue
- $3/2 = 1$       Remainder: 1 → This is the 3<sup>rd</sup> least significant binary digit  
Quotient =  $1 \neq 0$  → continue
- $1/2 = 0$       Remainder: 1 → This is the 4<sup>th</sup> least significant binary digit  
Quotient = 0 → exit the *repeat-until* control structure

□  $14_{10} = 1110_2 \cdot \bullet \bullet \bullet$

Note that, it is  $1110_2$   
It is NOT  $0111_2$

# Conversion from decimal to any other base system

■ Example 6: Convert  $2477_{10}$  to hexadecimal:

Hexadecimal means the new base is 16

□  $2477/16 = 154$  Remainder: 13 → This is the least significant Hex digit

Quotient =  $154 \neq 0$  → continue

□  $154/16 = 9$  Remainder: 10 → This is the 2<sup>nd</sup> least significant Hex digit

Quotient =  $9 \neq 0$  → continue

□  $9/16 = 0$  Remainder: 9 → This is the 3<sup>rd</sup> least significant Hex digit

Quotient = 0 → exit the *repeat-until* control structure

□  $2477_{10} = 9AD_{16}$


Note that, it is  $9AD_{16}$   
It is NOT  $DA9_{16}$

|        |        |
|--------|--------|
| 10 = A | 13 = D |
| 11 = B | 14 = E |
| 12 = C | 15 = F |

# Conversion from decimal to any other base system

## ■ Multiplication Method (for fraction numbers)

- Initialize the **fraction** by the value of the *fractional decimal number*
  - *Repeat*:
    - **Multiply** the **fraction** from the previous stage **by the new base** to get
      - A *whole number*
      - A new *fraction*
    - The *whole number* here is *the next digit to the right after the radix point* in the new number
- Until* the new **fraction** becomes 0.



This condition might be *reached* or not.  
*If reached*,  
then we will have 100% accurate result.  
*If not*,  
then we have to approximate the result.

# Conversion from decimal to any other base system

- Example 7: Convert  $0.017578125_{10}$  to hexadecimal

Hexadecimal means the new base is 16

$$\square 0.01757812 \times 16 = 0.28125$$

whole number: 0  $\rightarrow$  the next digit to the right after the radix point

fraction = 0.28125  $\neq 0 \rightarrow$  continue

$$\square 0.28125 \times 16 = 4.5$$

whole number: 4  $\rightarrow$  the next digit to the right after the radix point

fraction = 0.5  $\neq 0 \rightarrow$  continue

$$\square 0.5 \times 16 = 8.0$$

whole number: 8  $\rightarrow$  the next digit to the right after the radix point

fraction = 0.0  $\rightarrow$  exit the *repeat-until* control structure

$$\square 0.017578125_{10} = 0.048_{16}$$

# Conversion from decimal to any other base system

- Example 8: Convert  $255.017578125_{10}$  to hexadecimal:

Hexadecimal means the new base is 16

$$255.017578125_{10} = 255_{10} + 0.017578125_{10}$$

Using the *division method*:  $255_{10} \rightarrow FF_{16}$

Using the *multiplication method*:  $0.017578125_{10} \rightarrow 0.048_{16}$

$$255.017578125_{10} = FF.048_{16}$$

# Conversion from decimal to any other base system

- Example 9: Convert  $0.85_{10}$  to hexadecimal

Hexadecimal means the new base is 16

□  $0.85 \times 16 = 13.6$

whole number: 13 → *the next digit to the right after the radix point*

fraction =  $0.6 \neq 0$  → continue

□  $0.6 \times 16 = 9.6$

whole number: 9 → *the next digit to the right after the radix point*

fraction =  $0.6 \neq 0$  → continue

□  $0.6 \times 16 = 9.6$

whole number: 9 → *the next digit to the right after the radix point*

fraction =  $0.6 \neq 0$  → continue

□ ...

□  $0.85_{10} = 0.D99999...9_{16}$

□ Can be approximated in 4 digits after the radix point, for example, as

- $0.D999_{16}$  (using truncation) or as

- $0.D99A_{16}$  (using rounding)



## **Conversion between any two bases, other than decimal**

- This task can be done in **two steps**:
  - Convert from the **source base** to the **decimal**
  - Convert from the **decimal** to the **destination base**

## Conversion between any two bases, other than decimal

- **Example 10**: Convert  $2E8_{16}$  to *octal*

$$\begin{aligned} 2E8_{16} &= 2 \times 16^2 + E \times 16^1 + 8 \times 16^0 \\ &= 2 \times 256 + 14 \times 16 + 8 \times 1 \\ &= 512 + 224 + 8 = 744_{10} \end{aligned}$$

|        |        |
|--------|--------|
| 10 = A | 13 = D |
| 11 = B | 14 = E |
| 12 = C | 15 = F |

$744/8 = 93$  Remainder: 0 → This is the least significant octal digit

Quotient = 93  $\neq 0$  → continue

$93/8 = 11$  Remainder: 5 → This is the 2<sup>nd</sup> least significant octal digit

Quotient = 11  $\neq 0$  → continue

$11/8 = 1$  Remainder: 3 → This is the 3<sup>rd</sup> least significant octal digit

Quotient = 1  $\neq 0$  → continue

$1/8 = 0$  Remainder: 1 → This is the 4<sup>th</sup> least significant octal digit

Quotient = 0  $\neq 0$  → exit the *repeat-until* control structure

$$2E8_{16} = 744_{10} = 1350_8$$

## Conversion between any two bases, other than decimal (Special cases)

### ■ Binary to octal or hexadecimal:

#### □ *Binary to octal conversion*

- Group bits in three's, starting from the binary point  
(pad the last group with 0's, if needed)
- Convert each of these three-bit group into an octal digit

#### □ *Binary to hexadecimal conversion*

- Group bits in four's, starting from the binary point  
(pad the last group with 0's, if needed)
- Convert each of these four-bit group into a hexadecimal digit

## Conversion between any two bases, other than decimal (Special cases)

- Example 11: Convert  $11001111_2$  to *octal*

$11001111_2$

→  $011\ 001\ 111_2$

→  $317_8$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

## Conversion between any two bases, other than decimal (Special cases)

- Example 12: Convert  $1111010101_2$  to hexadecimal

$1111010101_2$

→  $0011\ 1101\ 0101_2$

→  $3D5_{16}$

|          |          |
|----------|----------|
| 0 = 0000 | 8 = 1000 |
| 1 = 0001 | 9 = 1001 |
| 2 = 0010 | A = 1010 |
| 3 = 0011 | B = 1011 |
| 4 = 0100 | C = 1100 |
| 5 = 0101 | D = 1101 |
| 6 = 0110 | E = 1110 |
| 7 = 0111 | F = 1111 |

## Conversion between any two bases, other than decimal (Special cases)

### ■ Octal or hexadecimal to binary:

#### □ *Octal to binary conversion*

- Expanding each octal digit into three bits

#### □ *Hexadecimal to binary conversion*

- Expanding each hexadecimal digit into four bits

## Conversion between any two bases, other than decimal (Special cases)

- Example 13: Convert  $743_8$  to binary

$743_8$

→  $111\ 100\ 011_2$

→  $111100011_2$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

## Conversion between any two bases, other than decimal (Special cases)

- Example 14: Convert  $FA9_{16}$  to binary

$FA9_{16}$

→  $1111\ 1010\ 1001_2$

→  $111110101001_2$

|          |          |
|----------|----------|
| 0 = 0000 | 8 = 1000 |
| 1 = 0001 | 9 = 1001 |
| 2 = 0010 | A = 1010 |
| 3 = 0011 | B = 1011 |
| 4 = 0100 | C = 1100 |
| 5 = 0101 | D = 1101 |
| 6 = 0110 | E = 1110 |
| 7 = 0111 | F = 1111 |



## Conversion between any two bases, other than decimal (Special cases)

### ■ Octal to hexadecimal or hexadecimal to octal:

- Convert from the **source** base to the **binary**
  - Expanding each digit into three bits (in case of octal) or four bits (in case of hexadecimal)
- Convert from the **binary** to the **destination** base
  - Group bits in three's (in case of octal) or four's (in case of hexadecimal), starting from the binary point (*pad the last group from both sides with 0's, if needed*)

## Conversion between any two bases, other than decimal (Special cases)

- Example 15: Convert  $ABC_{16}$  to octal

$ABC_{16}$

→  $1010\ 1011\ 1100_2$

→  $101010111100_2$

→  $101\ 010\ 111\ 100_2$

→  $5274_8$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

0 = 0000

1 = 0001

2 = 0010

3 = 0011

4 = 0100

5 = 0101

6 = 0110

7 = 0111

8 = 1000

9 = 1001

A = 1010

B = 1011

C = 1100

D = 1101

E = 1110

F = 1111

## Conversion between any two bases, other than decimal (Special cases)

- Example 16: Convert  $0.AB_{16}$  to octal

$0.AB_{16}$

→  $0.1010\ 1011_2$

→  $0.10101011_2$

→  $00.101\ 010\ 110_2$

→  $0.526_8$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

0 = 0000

1 = 0001

2 = 0010

3 = 0011

4 = 0100

5 = 0101

6 = 0110

7 = 0111

8 = 1000

9 = 1001

A = 1010

B = 1011

C = 1100

D = 1101

E = 1110

F = 1111

## Conversion between any two bases, other than decimal (Special cases)

- Example 17: Convert  $AB.BA_{16}$  to octal

$AB.BA_{16}$

→  $1010\ 1011.1011\ 1010_2$

→  $10101011.1011101_2$

→  $010\ 101\ 011.101\ 110\ 100_2$

→  $253.564_8$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

0 = 0000

1 = 0001

2 = 0010

3 = 0011

4 = 0100

5 = 0101

6 = 0110

7 = 0111

8 = 1000

9 = 1001

A = 1010

B = 1011

C = 1100

D = 1101

E = 1110

F = 1111

## Conversion between any two bases, other than decimal (Special cases)

- Example 18: Convert  $123_8$  to hexadecimal

$123_8$

→  $001\ 010\ 011_2$

→  $\quad\quad 1010011_2$

→  $0101\ 0\ 011_2$

→  $53_{16}$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

0 = 0000

1 = 0001

2 = 0010

3 = 0011

4 = 0100

5 = 0101

6 = 0110

7 = 0111

8 = 1000

9 = 1001

A = 1010

B = 1011

C = 1100

D = 1101

E = 1110

F = 1111

## Conversion between any two bases, other than decimal (Special cases)

- Example 19: Convert  $0.123_8$  to hexadecimal

$0.123_8$

→  $0.001\ 010\ 011_2$

→  $0.001010011_2$

→  $0000.0010\ 1001\ 1000_2$

→  $0.298_{16}$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

0 = 0000

1 = 0001

2 = 0010

3 = 0011

4 = 0100

5 = 0101

6 = 0110

7 = 0111

8 = 1000

9 = 1001

A = 1010

B = 1011

C = 1100

D = 1101

E = 1110

F = 1111

## Conversion between any two bases, other than decimal (Special cases)

- Example 20: Convert  $321.123_8$  to hexadecimal

$321.123_8$

→  $011\ 010\ 001.001\ 010\ 011_2$

→  $11010001.001010011_2$

→  $1101\ 0001.0010\ 1001\ 1000_2$

→  $D1.298_{16}$

0 = 000

1 = 001

2 = 010

3 = 011

4 = 100

5 = 101

6 = 110

7 = 111

0 = 0000

1 = 0001

2 = 0010

3 = 0011

4 = 0100

5 = 0101

6 = 0110

7 = 0111

8 = 1000

9 = 1001

A = 1010

B = 1011

C = 1100

D = 1101

E = 1110

F = 1111