


# Part 2

## CHAPTER 2

### Computer Arithmetic and Digital Logic



1

These slides are provided with permission from the copyright for CS2208 use only. The slides must not be reproduced or provided to anyone outside the class.

All downloaded copies of the slides are for personal use only.

Students must destroy these copies within 30 days after receiving the course's final assessment.

# Signed Integers

- Signed numbers can be represented in many ways, including
  - *sign and magnitude*,
  - *biased representation*, and
  - *two's complement*.

# Sign and Magnitude Representation

- ❑ An  $n$ -bit word has  $2^n$  possible values
  - for example, an eight-bit word can represent 256 different values.
- ❑ One way of representing a negative number is to take the *most-significant bit* and *reserve it* to indicate *the sign of the number*.
  - 0 represents **positive** numbers and
  - 1 represents **negative** numbers.
- ❑ *In an eight-bit word, using sign and magnitude representation*  
*How many positive numbers can be represented?*  
*How many negative numbers can be represented?*
- ❑ The value of a *sign and magnitude number* can be expressed as  $(-1)^S \times M$ , where **S** is the **sign bit** and **M** is the **magnitude** of the number.
  - If **S** = 0,  $(-1)^0 = +1$  and the number is positive.
  - If **S** = 1,  $(-1)^1 = -1$  and the number is negative.
- ❑ for example, in 8 bits we can *interpret* the numbers  $00001101_2$  and  $10001101_2$  as  $+13_{10}$  and  $-13_{10}$ , respectively.
- ❑ **Sign and magnitude** representation is
  - **not** generally **used in integer arithmetic**,
  - however, it is **used in floating-point arithmetic**.

## Biased Representation

- ❑ Assume that we have values ranging from  $-7$  to  $+7$
- ❑ To go around the negative sign, we can *shift the scale* (by adding a bias = 7 to all values) to have only non-negative values.
  - The original number =  $-7$  → The biased number = 0
  - The original number =  $-6$  → The biased number = 1
  - The original number = 0 → The biased number = 7
  - The original number =  $+6$  → The biased number = 13
  - The original number =  $+7$  → The biased number = 14
- ❑ To convert a biased number to its original value, you need to *shift back the scale* (by subtracting the bias from the biased number).
  - The biased number = 0 → The original number =  $-7$
  - The biased number = 1 → The original number =  $-6$
  - The biased number = 7 → The original number = 0
  - The biased number = 13 → The original number =  $+6$
  - The biased number = 14 → The original number =  $+7$
- ❑ In this representation, *biased numbers are unsigned integers*; yet, they *represent both positive and negative values*.
- ❑ The biased representation is also called excess- $K$ , where  $K$  is a pre-specified biasing value

Excess-7  
code

## Complementary Arithmetic

- A number and its **complement** add up to a constant;
  - In *n-digit decimal ten's complement* (*radix complement*) arithmetic, if  $P$  is a number, then its *ten's* complement is  $Q$ , where  $P + Q = 10^n$ .
    - the *ten's* complement of 12 is 88 because  $12 + 88 = 100$ .
    - the *ten's* complement of 88 is 12 because  $88 + 12 = 100$ .
  - In *n-bit binary two's complement* (*radix complement*) arithmetic, if  $P$  is a number, then its *two's* complement is  $Q$ , where  $P + Q = 2^n$ .
    - the *two's* complement of  $101_2$  is  $011_2$  because  $101_2 + 011_2 = 1000_2$ .
    - the *two's* complement of  $011_2$  is  $101_2$  because  $011_2 + 101_2 = 1000_2$ .

# Two's Complement Arithmetic

□ In binary arithmetic, the *two's complement* of a number is formed by

- *Subtracting the number from  $2^n$ .*

The *two's* complement of  $01100101_2$  is  
 $\overset{1}{\underset{1}{1}}00000000_2 - 01100101_2 = 10011011_2$

- *Flipping (inverting) all the bits of the number and adding 1.*

The *two's* complement of  $01100101_2$  is  
 $10011010_2 + 1_2 = 10011011_2$ .

- *Processing all the bits of the number from the least significant bit (LSB) towards the most significant bit (MSB)*
  - *copying all the zeros until the first 1 is reached,*
  - *copying that 1,*
  - *flipping (inverting) all the remaining bits.*

The *two's* complement of  $01100100_2$  is  $10011100_2$ .

The *two's* complement of  $01100101_2$  is  $10011011_2$ .

## Two's Complement Arithmetic


- ❑ The *two's* complement of a *positive* number is the corresponding *negative* value of the *positive* number when it is represented in *two's* complement.
- ❑ The *two's* complement of a *negative* number is the corresponding *positive* value of the *negative* number when it is represented in *two's* complement.
- ❑ We are interested in *complementary* arithmetic because *subtracting* a number is the same as *adding its complement*.
  - To subtract  $001100101_2$  from a binary number, we just need to add its *two's* complement (i.e.,  $110011011_2$ ) to the number.

## Two's Complement Arithmetic

- ❑ The *two's* complement of an  $n$ -bit binary value,  $N$ , is defined as  $2^n - N$ .
- ❑ If  $N = 5_{10} = 00000101_2$  (8-bit arithmetic), the *two's* complement of  $N$  is given by  $2^8 - 00000101_2 = 100000000_2 - 00000101_2 = 11111011_2$ .
- ❑  $11111011_2$  represents  $-00000101_2$  (i.e.,  $-5_{10}$ ) or  $-123_{10}$  depending only on whether we interpret  $11111011_2$  as a *two's complement integer* or as a *sign-magnitude integer*, respectively.



This is sign-magnitude not unsigned



This is  
-123 not +123



## Two's Complement Arithmetic

- This example demonstrates 8-bit *two's* complement arithmetic.

We begin by writing down the representations of +5, -5, +7 and -7.

$$+5_{10} = 00000101_2 \quad -5_{10} = 11111011_2 \quad +7_{10} = 00000111_2 \quad -7_{10} = 11111001_2$$

- Now  $7_{10} - 5_{10}$  can be calculated as  $7_{10} + (-5_{10})$  by adding the binary value for  $7_{10}$  to the *two's* complement of  $5_{10}$ .

$$\begin{array}{r}
 11111111 \\
 00000111 \quad 7 \\
 + 11111011 \quad -5 \\
 \hline
 100000010 \quad 2
 \end{array}$$

*The result is correct if the left-hand carry-out is ignored.*

## Two's Complement Arithmetic

□ Now consider the addition of  $-7_{10}$  to  $+5_{10}$ , i.e.,  $+5_{10} + (-7_{10})$

$$\begin{array}{r}
 \phantom{0}1 \\
 \textcolor{blue}{0}0000101 \qquad 5 \\
 + \textcolor{blue}{1}1111001 \qquad -7 \\
 \hline
 \textcolor{blue}{1}1111110 \qquad -2
 \end{array}$$

The result is  $\textcolor{blue}{1}1111110_2$  (the carry-out bit is  $\textcolor{red}{0}$ ).

The expected answer is  $-2_{10}$ ;

that is,  $2^8 - 2_{10} = 100000000_2 - \textcolor{blue}{0}0000010_2 = \textcolor{blue}{1}1111110_2$ .

## Two's Complement Arithmetic

- ❑ *Two's* complement arithmetic is not magic.
- ❑ Consider the calculation  $Z = X - Y$  in  $n$ -bit arithmetic
  - we do it by *adding* the *two's* complement of  $Y$  to  $X$ .
- ❑ The *two's* complement of  $Y$  is defined as  $2^n - Y$ .

We get

$$Z = X - Y = X + (2^n - Y) = 2^n + (X - Y).$$

- ❑ This is
  - *the desired result*,  $X - Y$
  - If the result of  $X - Y$  is negative, the *two's* complement will be automatically calculated, i.e.,  $2^n + (X - Y)$ , where  $(X - Y)$  is negative.
  - Otherwise, we discard the *unwanted carry-out digit* (i.e.,  $2^n$ ) at the leftmost position

# Two's Complement Arithmetic

## *Time examples using radix complement*

In the 24H system (*i.e.*,  $\text{radix} = 24$ ), assume that

- ❑ the time right now is 2 am and we want to subtract 5 hours from 2 am
  - The radix complement of 5 is  $24 - 5 = 19$
  - To subtract 5 from 2 in the 24H system, it is equivalent to adding the 2 to the complement of 5, *i.e.*,  $2 + 19 = 21$  (*correct answer*)
  - The 21 can also be read as  $-3$  in the radix complement system
- ❑ the time right now is 5 am and we want to subtract 2 hours from 5 am
  - The radix complement of 2 is  $24 - 2 = 22$
  - To subtract 2 from 5 in the 24H system, it is equivalent to adding the 5 to the complement of 2, *i.e.*,  $5 + 22 = 27$
  - As all the time values must be between 0 and 23, we need to subtract a full 24H from the result, *i.e.*,  $27 - 24 = 3$  (*correct answer*)

## Two's Complement Arithmetic

Let  $X = 9_{10} = 00001001_2$  and  $Y = 6_{10} = 00000110_2$

$$-X = -9_{10} = 100000000_2 - 00001001_2 = 11110111_2$$

$$-Y = -6_{10} = 100000000_2 - 00000110_2 = 11111010_2$$

$$\begin{array}{r} 1. \quad +X \quad +9 \quad 00001001 \\ \quad +Y \quad +6 \quad +00000110 \\ \hline \quad \quad +15 \quad 00001111 \end{array}$$

$$\begin{array}{r} 2. \quad +X \quad +9 \quad 00001001 \\ \quad -Y \quad -6 \quad +11111010 \\ \hline \quad \quad +3 \quad 100000011 \end{array}$$

$$\begin{array}{r} 3. \quad -X \quad -9 \quad 11110111 \\ \quad +Y \quad +6 \quad +00000110 \\ \hline \quad \quad -3 \quad 11111101 \end{array}$$

$$\begin{array}{r} 4. \quad -X \quad -9 \quad 11110111 \\ \quad -Y \quad -6 \quad +11111010 \\ \hline \quad \quad -15 \quad 111110001 \end{array}$$

**11111101** is a negative *two's* complement number

The two's complement of **11111101** is **00000011**<sub>2</sub>

Hence, **11111101** = **-00000011**<sub>2</sub> = **-3**<sub>10</sub>

**11110001** is a negative *two's* complement number

The two's complement of **11110001** is **00001111**<sub>2</sub>

Hence, **11110001** = **-00001111**<sub>2</sub> = **-15**<sub>10</sub>

All four examples give the result we would expect when the result is interpreted as a *two's* complement number.

## Properties of Two's complement Numbers

- ❑ The *two's* complement system is a true complement system in that  $+X + (-X) = 0$ .
- ❑ There is *one* unique zero 00...0.
- ❑ This is *not the case with the signed magnitude* number system
- ❑ The *most-significant bit* of a *two's* complement number is a *sign bit*.
  - The number is *positive* if the most-significant bit is 0, and
  - *negative* if it is 1.

This is the same as the signed magnitude number system

- ❑ The range of an *n-bit two's* complement number is from  $-2^{n-1}$  to  $+2^{n-1} - 1$ .

- For  $n = 4$ , the range is  $-8$  to  $+7$ .

The total number of different values is  $2^n = 16$

- 8 negative,
- zero and
- 7 positive.

*these two  
are same  
number.*

*000001 => 011111*

*000000*

*011111 + 100001 = 0*

*100000*

- For  $n = 8$ , the range is  $-128$  to  $+127$ .

The total number of different values is  $2^n = 256$

- 128 negative,
- zero and
- 127 positive.

*2's complement:*

*$2^{n-1}$  negative 10*

*$2^{n-1} - 1$  positive.*

- ❑ *How about the signed magnitude number system?*

## Arithmetic Overflow

- ❑ The *range* of *two's* complement numbers in *n-bits* is from  $-2^{n-1}$  to  $+2^{n-1} - 1$ .
- ❑ What happens if we *violate* this rule by carrying out an operation whose *result falls outside the range* of values that can be represented by *two's* complement numbers?
  - In a *five-bit* representation, the range of *valid two's complement numbers* is  $-16$  to  $+15$ .

### Case 1

$$\begin{array}{r} 5 = 00101 \\ +7 = 00111 \\ \hline 12 = 01100 = 12_{10} \end{array}$$

### Case 2

$$\begin{array}{r} 12 = 01100 \\ +13 = 01101 \\ \hline 25 = 11001 = -7_{10} \text{ (as a two's complement value)} \end{array}$$

In *Case 1* we get the *expected* answer of  $+12_{10}$ , but

In *Case 2* we get a *negative* result because the *sign bit is 1*.

- If the answer were regarded as an *unsigned binary number*, it would be  $+25$ , which is, of course, the correct answer.
- However, as the *two's* complement system has been chosen to represent signed numbers, all answers must be interpreted in this light.

## Arithmetic Overflow

- If we add together two negative numbers whose total is less than  $-16$ , we will go out of range.

### Case 3

$$\begin{array}{rcl} -9 & = & 10111 \\ -12 & = & +10100 \\ \hline -21 & & 101011 = +11_{10} \text{ (as a } \textit{two's} \text{ complement value)} \end{array}$$

In *Case 3* we get a **positive** result because the *sign bit is 0*.



## Arithmetic Overflow

- ❑ The last two examples demonstrate *arithmetic overflow*
- ❑ *Arithmetic overflow* occurs during a *two's complement addition* when
  - the result of *adding two positive* numbers *yields a negative* result,
  - or
  - the result of *adding two negative* numbers *yields a positive* result.
- ❑ *If the sign bits of A and B are the same but the sign bit of the result is different, it means arithmetic overflow has occurred.*
- ❑ *Overflow never occurs when adding two numbers with opposite signs*
- ❑ If  $a_{n-1}$  is the sign bit of A,  
 $b_{n-1}$  is the sign bit of B, and  
 $s_{n-1}$  is the sign bit of the sum of A and B,  
 then  
 overflow is defined by the *logical expression* as follow:  

$$V = (a_{n-1})^c \cdot (b_{n-1})^c \cdot s_{n-1} + a_{n-1} \cdot b_{n-1} \cdot (s_{n-1})^c$$
 where  $()^c$  is the complement binary operation

# Arithmetic Overflow

- In practice, the overflow can be detected from the *carry bits* *in to* (i.e.,  $C_{in}$ ) and *out of* (i.e.,  $C_{out}$ ) the *most-significant bit* of an adder;

*overflow occurs when  $C_{in} \neq C_{out}$*

**8-bit examples**

*8-bit examples in slide#30*

$C_{out}$   $C_{in}$

$00000000$

$+X \quad +9 \quad 00001001$

$+Y \quad +6 \quad +00000110$

$+15 \quad 00001111$

$C_{out}$   $C_{in}$

$00000110$

$-X \quad -9 \quad 11110111$

$+Y \quad +6 \quad +00000110$

$-3 \quad 11111101$

$C_{out}$   $C_{in}$

$00111$

$5 = 00101$

$+7 = 00111$

$12 \quad 01100 = +12_{10}$

$C_{out}$   $C_{in}$

$11111000$

$+X \quad +9 \quad 00001001$

$-Y \quad -6 \quad +11111010$

$+3 \quad 100000011$

$C_{out}$   $C_{in}$

$11111110$

$-X \quad -9 \quad 11110111$

$-Y \quad -6 \quad +11111010$

$-15 \quad 111110001$

$C_{out}$   $C_{in}$

$10100$

$-9 = 10111$

$-12 = +10100$

$-21 \quad 101011 = +11_{10}$

$C_{out}$   $C_{in}$

$01100$

$12 = 01100$

$+13 = 01101$

$25 \quad 11001 = -7_{10}$

**Overflow occurred**

$C_{out}$   $C_{in}$

$10100$

$-9 = 10111$

$-12 = +10100$

$-21 \quad 101011 = +11_{10}$

**Overflow occurred**

*5-bit examples in slides#32,33*

# Shifting Operations


- ❑ In a shift operation, the *bits of a word* are *shifted* one or more places to the *left* or *right*.
- ❑ The *arithmetic* shift is just one type of shift operation.
- ❑ In Chapter 3, we will cover the other types of shift operations.

## Shifting Operations (left)

- Figure 2.2(a) describes the arithmetic *shift left*.
  - A *zero* enters into the vacated *least-significant bit* position and
  - the bit *shifted out* of the *most-significant bit* position is recorded in the computer's *carry flag*.
  - In *two's* complement and unsigned numbers:
    - Arithmetic *shift left* means *multiplying* by 2.
- If  $00100111_2$  ( $39_{10}$ ) is shifted one place left it becomes  $01001110_2$  ( $78_{10}$ ).
- If  $11100011_2$  ( $-29_{10}$ ) is shifted one place left it becomes  $11000110_2$  ( $-58_{10}$ ).

FIGURE 2.2

Arithmetic shift operations



(a) Arithmetic shift left:


A zero enters the least-significant bit position and the most-significant bit is copied into the carry flag. For example:  $11000101$  becomes  $10001010$  after shifting one place left.

## Shifting Operations (**right**)

- ❑ Figure 2.2(b) describes the arithmetic *shift right*.
  - the **sign bit** (the **MSB**) is *replicated* into the vacated *most-significant bit* position and
  - the bit *shifted out* of the *least-significant bit* position is recorded in the computer's *carry flag*.
- In *two's* complement and unsigned numbers:
  - Arithmetic *shift right* means *dividing* by 2.
- ❑ Shifting  $00001100_2$   $(12)_{10}$  one place right produces  $00000110_2$   $(6)_{10}$ .
- ❑ Shifting  $00001101_2$   $(13)_{10}$  one place right produces  $00000110_2$   $(6)_{10}$  as well.
- ❑ Shifting  $11100010_2$   $(-30)_{10}$  one place right produces  $11110001_2$   $(-15)_{10}$ .
- ❑ Shifting  $11100011_2$   $(-29)$  one place right produces  $11110001_2$   $(-15)$  as well.

FIGURE 2.2

Arithmetic shift operations



(b) Arithmetic shift right:

A copy of the most-significant bit enters the most-significant bit position. All other bits are shifted one place right. The least-significant bit is copied into the carry flag.

For example; 00100101 becomes 00010010 after shifting one place right, and 11100101 becomes 11110010 after shifting one place right.

© Cengage Learning 2014

# Binary Multiplication

- Multiplication can be implemented using arithmetic *shift left* and *addition* operations to add up the partial products as they are formed.

Multiplicand	Multiplier	Step	Partial products
01101001	0100100 <b>1</b>	1	0 1 1 0 1 0 0 1
01101001	010010 <b>0</b> 1	2	0 0 0 0 0 0 0 0
01101001	01001 <b>0</b> 01	3	0 0 0 0 0 0 0 0
01101001	0100 <b>1</b> 001	4	0 1 1 0 1 0 0 1
01101001	010 <b>0</b> 1001	5	0 0 0 0 0 0 0 0
01101001	01 <b>0</b> 01001	6	0 0 0 0 0 0 0 0
01101001	0 <b>1</b> 001001	7	0 1 1 0 1 0 0 1
01101001	<b>0</b> 1001001	8	0 0 0 0 0 0 0 0
<b>Result</b>			<b>0 0 1 1 1 0 1 1 1 1 0 0 0 1</b>