

Multivariate Analysis
MATH 3109
UCL

Based on lectures by Dr. Iannis Petridis
Typed by John Sylvester

Autum 2011

Contents

1	Multivariable Calculus	2
1.1	Notation	2
1.2	Functions & Continuity	3
1.3	Partial Derivatives	5
1.4	Total Derivative	6
1.5	The Chain Rule	9
1.6	Mixed Derivatives	11
1.7	Jacobian	13
1.8	Inverse Function Theorem	15
1.9	Implicit Function Theorem	20
2	Integration	25
2.1	Multiple integrals	25
2.2	Fubini's theorem	28
2.3	Change of variables	29
3	Calculus on Manifolds	29
3.1	Manifolds	29
3.2	Dual Space	35
3.3	Multilinear Algebra	37
3.4	Alternating Tensors	40
3.5	Differential Forms	46
3.6	Closed and Exact forms	52
3.7	Stoke's Theorem	60
3.8	Classical Stoke's Theorem in \mathbb{R}^2	65

1 Multivariable Calculus

1.1 Notation

$X \in \mathbb{R}^n$, $X = \{x_1, x_2, \dots, x_n\}$ where $x_i \in \mathbb{R}$ \mathbb{R}^n is a vector space
length norm:

$$|x| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$$

If $Y, X \in \mathbb{R}^n$ and $Y = \{y_1, y_2, \dots, y_n\}$ then

$$X \cdot Y = x_1 y_1 + x_2 y_2 + \dots + x_n y_n$$

Standard Basis:

$$e_j = (0, \dots, 0, 1, 0, \dots)$$

j-1, j, j+1

Properties of norm

$$|x| \geq 0$$

$$|x| = 0 \Leftrightarrow x = \vec{0}$$

$$|\lambda x| = |\lambda| \cdot |x|, \quad x \in \mathbb{R}^n, \quad \lambda \in \mathbb{R}$$

linear Transformation

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$(i) \quad T(x + y) = T(x) + T(y)$$

$$(ii) \quad T(\lambda x) = \lambda T(x)$$

Matrix Representation of T with respect to the standard basis:

$$T(e_i) = \sum_{j=1}^m a_{i,j} e_j \quad \text{where} \quad [T]_{\epsilon}^{\epsilon} = A = (a_{i,j})_{\substack{i=1,\dots,m \\ j=1,\dots,n}}$$

Given: $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $U : \mathbb{R}^m \rightarrow \mathbb{R}^k$

$$(i) \quad [UT]_{kxm} = [U]_{kxm} [T]_{m \times n}$$

$$(ii) \quad [T + S] = [T] + [S]$$

$$(iii) \quad \lambda [T] = [\lambda T]$$

$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $X \in \mathbb{R}^n$, $Y \in \mathbb{R}^m$, $X = (x^1, \dots, x^n)$, $Y = (y^1, \dots, y^m)$

$$\begin{pmatrix} y^1 \\ y^2 \\ \vdots \\ y^m \end{pmatrix} = [T] \begin{pmatrix} x^1 \\ x^2 \\ \vdots \\ x^n \end{pmatrix}$$

1.2 Functions & Continuity

$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ vector valued function

$f : A \rightarrow \mathbb{R}^m$ where $A \subset \mathbb{R}^n$

then f has components which are scalar fields.

$f^i : A \rightarrow \mathbb{R}$

$$f(x) = (f^1(x), \dots, f^m(x))$$

$\Pi^i : \mathbb{R}^m \rightarrow \mathbb{R}$

$$\Pi^i((x)^1, \dots, (x)^m)$$

Π^i is a linear transformation for $i=1, \dots, m$

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{f} & \mathbb{R}^m \\ & \searrow f^i & \downarrow \Pi^i \\ & & \mathbb{R} \end{array}$$

Definition 1.1. $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ then $\lim_{x \rightarrow a}(f(x)) = b$ means:

$$\forall \epsilon > 0, \exists \delta > 0 \text{ st, } 0 < |x - a| < \delta \Rightarrow |f(x) - b| < \epsilon$$

Definition 1.2. f is called continuous at a if:

$$\lim_{x \rightarrow a}(f(x)) = f(a)$$

f is called continuous on the set A if it is continuous at a , $\forall a \in A$

Theorem 1.1 (Combination Theorem). Assume

$$\lim_{x \rightarrow a}(f(x)) = b, \quad \lim_{x \rightarrow a}(g(x)) = c$$

then:

$$(i) \lim_{x \rightarrow a}(f(x) + g(x)) = b + c$$

$$(ii) \lim_{x \rightarrow a}(\lambda f(x)) = \lambda b$$

$$(iii) \lim_{x \rightarrow a}(f(x) \cdot g(x)) = b \cdot c$$

$$(iv) \lim_{x \rightarrow a} |f(x)| = |b|$$

Proof. of (iii)

$$\begin{aligned} f(x) \cdot g(x) - b \cdot c &= f(x) \cdot g(x) - b \cdot g(x) + b \cdot g(x) - b \cdot c \\ &= g(x)(f(x) - b) + b \cdot (g(x) - c) \\ |f(x) \cdot g(x) - b \cdot c| &= |g(x)(f(x) - b) + b \cdot (g(x) - c)| \\ &\leq |g(x)(f(x) - b)| + |b \cdot (g(x) - c)| \end{aligned}$$

Cauchy-Schwartz: $|x^1y^1 + \dots + x^ny^n| \leq \sqrt{(x^1)^2 + \dots + (x^n)^2} \cdot \sqrt{(y^1)^2 + \dots + (y^n)^2}$

$$|f(x) \cdot g(x) - b \cdot c| \leq |g(x)(f(x) - b)| + |b \cdot (g(x) - c)| \leq |g(x)| \cdot |f(x) - b| + |b| \cdot |g(x) - c|$$

Since $\lim_{x \rightarrow a}(g(x)) = c$, g is a bounded neighbourhood of a , i.e:

$$\exists M \geq 0, \exists \delta > 0 \text{ st, } |g(x)| \leq M \text{ for } |x - a| < \delta$$

□

Remark. We have:

(i) $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuous iff: $f^i : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous for $i = 1, \dots, m$

(ii) Polynomial functions in n -variables, $f(x^1, \dots, x^n)$, are continuous

(iii) Rational functions, $R(x) = \frac{P(x)}{Q(x)}$, are continuous where defined, ie: $Q(x) \neq 0$ and P, Q are polynomials in n -variables.

Theorem 1.2. Linear transformations are continuous.

Proof. $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ let $a \in \mathbb{R}^n$ to show:

$$\lim_{h \rightarrow 0} T(a + h) = T(a)$$

where $h = (h^1, \dots, h^n)$

$$\begin{aligned} |T(a + h) - T(a)| &= |T(h)| = |T(h^1e_1 + \dots + h^ne_n)| = |h^1T(e_1) + \dots + h^nT(e_n)| \\ &\leq |h^1||T(e_1)| + \dots + |h^n||T(e_n)| \leq |h|(T(e_1) + \dots + T(e_n)) \end{aligned}$$

$$\text{So : } |T(a + h) - T(a)| \leq M|h| \quad \text{where} \quad M = \sum_{i=1}^n |T(e_i)|$$

$$\text{So given } \epsilon > 0, \quad \text{choose } \delta = \frac{\epsilon}{M} \quad \text{such that} \quad |h| < \delta \Rightarrow |T(a + h) - T(a)| < \epsilon$$

□

Example 1.1. $f(x, y) = \frac{x^2 - y^2}{x^2 + y^2}$, $(x, y) = (0, 0)$ assume $\lim_{(x, y) \rightarrow (0, 0)} f(x, y) = L$

$$\forall \epsilon > 0, \quad \exists \delta > 0 \quad \text{such that} \quad 0 < |(x, y)| < \delta \Rightarrow |f(x, y) - L| < \epsilon$$

Plug $(x, 0)$ into f :

$$f(x, 0) = \frac{x^2 - 0}{x^2 + 0} = 1$$

Plug $(0, y)$ into f :

$$f(0, y) = \frac{0 - y^2}{0 + y^2} = -1$$

$$\begin{aligned}
\text{If } |x| < \delta \quad |f(x, 0)| < \delta &\Rightarrow |f(x, 0) - L| < \epsilon \quad \text{ie} \quad |1 - L| < \epsilon \\
\text{If } |y| < \delta \quad |f(0, y)| < \delta &\Rightarrow |f(0, y) - L| < \epsilon \quad \text{ie} \quad |-1 - L| < \epsilon \\
&\Rightarrow \epsilon = \frac{1}{2} \quad \text{contradiction!}
\end{aligned}$$

Now consider $y = mx, m \in \mathbb{R}$

$$\begin{aligned}
f(x, mx) &= \frac{x^2 - (mx)^2}{x^2 + (mx)^2} = \frac{1 - m^2}{1 + m^2} \\
\lim_{x \rightarrow 0} (\lim_{y \rightarrow 0} f(x, y)) &= \lim_{x \rightarrow 0} 1 = 1 \\
\lim_{y \rightarrow 0} (\lim_{x \rightarrow 0} f(x, y)) &= \lim_{y \rightarrow 0} -1 = -1
\end{aligned}$$

However checking along straight lines is not enough to prove continuity.

Example 1.2.


$$f(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2 + y^2}} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Show f is continuous at $(0, 0)$

$$\forall \epsilon > 0, \quad \exists \delta > 0$$

$$\left| \frac{xy}{\sqrt{x^2 + y^2}} \right| \leq \frac{|x| \cdot |y|}{\sqrt{x^2 + y^2}} \leq \frac{\sqrt{x^2 + y^2} \cdot \sqrt{x^2 + y^2}}{\sqrt{x^2 + y^2}} = \sqrt{x^2 + y^2} = |(x, y)|$$

Since:



Note. if the total degree of the numerator is higher than the denominator in a rational function, then the limit should be 0.

Theorem 1.3. If f is continuous at a and g is continuous at $f(a)$ then $g \circ f$ is continuous at a .

1.3 Partial Derivatives

Definition 1.3. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $a \in \mathbb{R}$ Define:

$$D_i f(a) = \lim_{h \rightarrow 0} \frac{f(a^1, \dots, a^{i-1}, a^i + h, a^{i+1}, \dots, a^n) - f(a)}{h}$$

Example 1.3. if $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\left. \frac{df}{dx} \right|_{(a,b)} = D_1 f(a, b)$$

$$\left. \frac{df}{dy} \right|_{(a,b)} = D_2 f(a, b)$$

and in \mathbb{R}^3 we use $\frac{df}{dx}$, $\frac{df}{dy}$ and $\frac{df}{dz}$ etc.

Example 1.4.

$$f(x, y) = \begin{cases} \frac{x^2 - y^2}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 1 & \text{if } (x, y) = (0, 0) \end{cases}$$

$$D_1 f(0, 0) = \left. \frac{df}{dx} \right|_{(0,0)} = \lim_{x \rightarrow 0} \frac{f(x, 0) - f(0, 0)}{x} = \lim_{x \rightarrow 0} \frac{\frac{x^2 - 0}{x^2 + 0} - 1}{x} = 0$$

$$D_2 f(0, 0) = \left. \frac{df}{dy} \right|_{(0,0)} = \lim_{y \rightarrow 0} \frac{f(0, y) - f(0, 0)}{y} = \lim_{y \rightarrow 0} \frac{\frac{0 - y^2}{0 + y^2} - 1}{y} = \lim_{y \rightarrow 0} \frac{-2}{y} = \pm\infty$$

1.4 Total Derivative

In 1 dimension we write the following for the derivative of $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

we try to write it in higher dimensions $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ in this form

$$\begin{aligned} \lim_{h \rightarrow 0} \left[\frac{f(a + h) - f(a)}{h} - f'(a) \right] &= \lim_{h \rightarrow 0} \left[\frac{f(a + h) - f(a) - h \cdot f'(a)}{h} \right] \\ &= \lim_{h \rightarrow 0} \frac{|f(a + h) - f(a) - h \cdot f'(a)|}{|h|} = 0 \end{aligned}$$

For $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ consider the tangent line at a : $y = f(a) + f'(a)(x - a)$
call $x - a = h$ then we have:

$$y = f(a) + f'(a)(h)$$

this is an Affine transformation, not a linear map.

Look at the map:

$$\lambda : h \rightarrow hf'(a), \quad h \in \mathbb{R}$$

This is a linear map.

$$\lambda(h_1 + h_2) = (h_1 + h_2)f'(a) = h_1f'(a) + h_2f'(a) = \lambda(h_1) + \lambda(h_2)$$

$$\lambda(\alpha \cdot h) = (\alpha h)f'(a) = \alpha(hf'(a)) = \alpha \cdot \lambda(h)$$

$$\lim_{h \rightarrow 0} \frac{|f(a + h) - f(a) - \lambda(h)|}{|h|} = 0$$

Definition 1.4 (Total Derivative). $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ or $(f : A \rightarrow \mathbb{R}^m, A \subset \mathbb{R}^n, A \text{ is open})$ is differentiable at a ($a \in A$) if we can find a linear transformation $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$ st:

$$\lim_{h \rightarrow 0} \frac{|f(a + h) - f(a) - \lambda(h)|}{|h|} = 0$$

The linear transformation λ is called the total derivative of f at a and denoted $Df(a)$ st

$$Df(a) = \lambda(h)$$

Example 1.5. $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $f(x) = k$, $k \in \mathbb{R}^m$ is differentiable at $a \in \mathbb{R}^n$ with the 0 linear transformation $0 : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $0(h) = 0$

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - 0(h)|}{|h|} = \lim_{h \rightarrow 0} \frac{|k - k - 0|}{|h|} = 0$$

Example 1.6. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, it is differentiable at $a \in \mathbb{R}^n$ with linear transformation $Df(a) = f$

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - f(h)|}{|h|} = \lim_{h \rightarrow 0} \frac{|f(a+h-a-h)|}{|h|} = 0$$

Theorem 1.4 (Uniqueness of Total Derivative). If f is differentiable at a then there exists a unique linear transformation, $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$, such that

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} = 0$$

Proof. suppose $\mu : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is another linear transformation such that:

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \mu(h)|}{|h|} = 0$$

deduce that $\lambda = \mu$ i.e. $\forall h \in \mathbb{R}^n$

$$\lambda(h) = \mu(h)$$

$$\begin{aligned} \frac{|\lambda(h) - \mu(h)|}{|h|} &= \frac{|\lambda(h) + f(a) - f(a+h) + f(a+h) - f(a) - \mu(h)|}{|h|} \\ &\leq \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} + \frac{|f(a+h) - f(a) - \mu(h)|}{|h|} \end{aligned}$$

Conclude that:

$$\lim_{h \rightarrow 0} \frac{|\lambda(h) - \mu(h)|}{|h|} \leq 0 + 0 = 0 \quad (*)$$

Let $h=0$ $\lambda = 0 = \mu$ since λ, μ are linear. Now fix $h \in \mathbb{R}^n$, $h \neq 0$ and let $t \in \mathbb{R}$ such that $th \in \mathbb{R}^n$ then replace h with th in (*):

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{|\lambda(th) - \mu(th)|}{|th|} &= \lim_{t \rightarrow 0} \frac{|t\lambda(h) - t\mu(h)|}{|t||h|} \\ &= \lim_{t \rightarrow 0} \frac{|t|}{|t|} \frac{|\lambda(h) - \mu(h)|}{|h|} = \frac{|\lambda(h) - \mu(h)|}{|h|} = 0 \end{aligned}$$

$$\Rightarrow |\lambda(h) - \mu(h)| = 0 \Rightarrow \lambda(h) = \mu(h)$$

deduce that $\lambda = \mu$ i.e. $\forall h \in \mathbb{R}^n$

$$\lambda(h) = \mu(h)$$

□

Definition 1.5 (Jacobian Matrix). $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at $a \in \mathbb{R}^n$ and it is derivative at a $Df(a) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear map. Then the matrix representation of $Df(a)$ is $f'(a) \in \mathbb{M}_{m \times n}$ and is called the Jacobian Matrix of f at a .

Example 1.7. $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $f(x, y) = (x^2y, x + 5)$ $x, y \in \mathbb{R}$
Show that $Df(1, 2)(h^1, h^2) = (4h^1 + h^2, h^1)$:

$$\begin{aligned} & f((1, 2) + (h^1, h^2)) - f(1, 2) - Df(1, 2)(h^1, h^2) \\ &= f(1 + h^1, 2 + h^2) - f(1, 2) - (4h^1 + h^2, h^1) \\ &= ((1 + h^1)^2(2 + h^2), (1 + h^1 + 5)) - (2, 6) - (4h^1 + h^2, h^1) \\ &= (2 + h^2 + 2(h^1)^2 + (h^1)^2h^2 + 2h^1h^2 + 4h^1 - 2 - 4h^1 - h^2, 6 + h^1 - 6 - h^1) \end{aligned}$$

Take length:

$$|(2(h^1)^2 + (h^1)^2h^2 + 2h^1h^2, 0)| \leq 2|h|^2 + |h|^2|h| + 2|h||h| = 4|h|^2 + |h|^3$$

So:

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{|f((1, 2) + (h^1, h^2)) - f(1, 2) - Df(1, 2)(h^1, h^2)|}{|h|} \\ & \leq \lim_{h \rightarrow 0} \frac{4|h|^2 + |h|^3}{|h|} = \lim_{h \rightarrow 0} 4|h| + |h|^2 = 0 \end{aligned}$$

Definition 1.6. $f'(a)$ is the matrix representation of $Df(a)$

$$\begin{aligned} Df(a)(h)^t &= \begin{pmatrix} y^1 \\ y^2 \\ \vdots \\ y^m \end{pmatrix} = f'(a) \begin{pmatrix} h^1 \\ h^2 \\ \vdots \\ h^n \end{pmatrix} \\ f'(a) &= \begin{pmatrix} D_1f^1(a) & D_2f^1(a) & \cdots & D_nf^1(a) \\ D_1f^2(a) & D_2f^2(a) & \cdots & D_nf^2(a) \\ \vdots & \vdots & \ddots & \vdots \\ D_1f^m(a) & D_2f^m(a) & \cdots & D_nf^m(a) \end{pmatrix} \end{aligned}$$

Example 1.8. With this new information we can tackle example 1.7:

$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $f(x, y) = (x^2y, x + 5)$ $x, y \in \mathbb{R}$
Show that $Df(1, 2)(h^1, h^2) = (4h^1 + h^2, h^1)$:

$$\frac{df^1}{dx} = 2xy, \quad \frac{df^1}{dy} = x^2, \quad \frac{df^2}{dx} = 1, \quad \frac{df^2}{dy} = 0$$

$$f'(1, 2) = \begin{pmatrix} 4 & 1 \\ 1 & 0 \end{pmatrix}$$

$$f'(1, 2) \begin{pmatrix} h^1 \\ h^2 \end{pmatrix} = \begin{pmatrix} 4 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} h^1 \\ h^2 \end{pmatrix} = \begin{pmatrix} 4h^1 + h^2 \\ h^1 \end{pmatrix}$$

Remark. Having directional derivatives in all directions $u \neq 0$ is not enough to guarantee $df(a)$ exists.

Theorem 1.5. *If f is differentiable at a then f is continuous at a .*

Proof.

$$\begin{aligned}\lim_{h \rightarrow 0} |f(a+h) - f(a)| &= \lim_{h \rightarrow 0} |f(a+h) - f(a) - Df(a)h + Df(a)h| \\ &\leq \lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - Df(a)h|}{|h|} \cdot |h| + \lim_{h \rightarrow 0} |Df(a)h| \\ &= 0\end{aligned}$$

since $Df(a)$ is a linear transformation $Df(a)$ is continuous so:

$$\lim_{h \rightarrow 0} |Df(a)h| = |Df(a)0| = 0$$

□

1.5 The Chain Rule

Theorem 1.6 (Chain Rule). *if $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at a and $g : \mathbb{R}^m \rightarrow \mathbb{R}^k$ is differentiable at $f(a)$ then $g \circ f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is differentiable at a and*

$$D(g \circ f)(a) = Dg(f(a)) \circ Df(a)$$

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{Df(a)} & \mathbb{R}^m \\ & \searrow D(g \circ f)(a) & \downarrow Dg(f(a)) \\ & & \mathbb{R}^k \end{array}$$

$$(g \circ f)'(a) = g'(f(a)) \cdot f'(a), \quad \text{where } \cdot \text{ represents matrix multiplication}$$

Proof. if $b = f(a)$ and we let $Df(a) = \lambda$ and $Dg(f(a)) = \mu$ then if we define:

$$\varphi(x) = f(x) - f(a) - \lambda(x - a) \tag{1}$$

$$\psi(y) = g(y) - g(b) - \mu(y - b) \tag{2}$$

$$\rho(x) = g \circ f(x) - g \circ f(a) - \mu \circ \lambda(x - a) \tag{3}$$

Then:

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - Df(a)h|}{|h|} = \lim_{x \rightarrow a} \frac{|\varphi(x)|}{|x - a|} = 0 \tag{4}$$

$$\lim_{h \rightarrow 0} \frac{|g(b+h) - g(b) - Dg(b)h|}{|h|} = \lim_{y \rightarrow b} \frac{|\psi(y)|}{|y - b|} = 0 \tag{5}$$

We must show:

$$\lim_{h \rightarrow 0} \frac{|g \circ f(x) - g \circ f(a) - \mu \circ \lambda(x - a)|}{|h|} = \lim_{x \rightarrow b} \frac{|\rho(x)|}{|x - b|} = 0$$

Now:

$$\begin{aligned}\rho(x) &= g(f(x)) - g(b) - \mu(\lambda(x - a)) \\ &= g(f(x)) - g(b) - \mu(f(x) - f(a) - \varphi(x)) && \text{by (1)} \\ &= [g(f(x)) - g(b) - \mu(f(x) - f(a))] + \mu(\varphi(x)) \\ &= \psi(f(x)) + \mu(\varphi(x)) && \text{by (2)}\end{aligned}$$

Thus we must Prove

$$\lim_{x \rightarrow a} \frac{|\psi(f(x))|}{|x - a|} = 0 \quad (6)$$

$$\lim_{x \rightarrow a} \frac{|\mu(\varphi(x))|}{|x - a|} = 0 \quad (7)$$

It follows from (4) that for some $\delta > 0$ we have

$$|\psi(f(x))| < \epsilon |f(x) - b| \quad \text{if } |f(x) - b| < \delta$$

which is true if $|x - a| < \delta_1$ for a suitable δ_1 . We also have that if T is a linear transformation then $\exists M \geq 0$ such that $|T(x)| < M|x|$. So then:

$$\begin{aligned} |\psi(f(x))| &< \epsilon |f(x) - b| \\ &= \epsilon |\varphi(x) + \lambda(x - a)| \\ &\leq \epsilon |\varphi(x)| + \epsilon M |x - a| \end{aligned}$$

So

$$\lim_{x \rightarrow a} \frac{|\psi(f(x))|}{|x - a|} \leq \lim_{x \rightarrow a} \frac{\epsilon |\varphi(x)|}{|x - a|} + \lim_{x \rightarrow a} \frac{\epsilon M |x - a|}{|x - a|} = \epsilon M \rightarrow 0$$

Also

$$\lim_{x \rightarrow a} \frac{|\mu\varphi(x)|}{|x - a|} \leq \lim_{x \rightarrow a} \frac{M|\varphi(x)|}{|x - a|} = 0$$

□

Theorem 1.7. Define $s : \mathbb{R}^2 \rightarrow \mathbb{R}$ $s(x, y) = x + y$ then s is differentiable and $Ds = s$

Proof. S is linear i.e

$$\begin{aligned} s((x, y) + (x', y')) &= s(x + x', y + y') = s(x, y) + s(x', y') \\ s(\lambda(x, y)) &= \lambda s(x, y) \end{aligned}$$

So

$$\lim_{h \rightarrow 0} \frac{|s(a + h) - s(a) - s(h)|}{|h|} = 0$$

□

Theorem 1.8. Define $p : \mathbb{R}^2 \rightarrow \mathbb{R}$, $p(x, y) = xy$, then p is differentiable and:
 $Dp(a, b) : \mathbb{R}^2 \rightarrow \mathbb{R}$ is linear with $Dp(a, b)(h, k) = ak + bh$ and $p' = (b, a)$

Proof. use of derivative

$$\begin{aligned} p((a, b) + (h, k)) - p(a, b) - Dp(a, b)(h, k) &= p(a + h, b + k) - p(a, b) - (ak + bh) \\ &= (a + h)(b + k) - ab - (ak + bh) = hk \\ \frac{|p((a, b) + (h, k)) - p(a, b) - Dp(a, b)(h, k)|}{|(h, k)|} &= \frac{|hk|}{\sqrt{h^2 + k^2}} \leq \frac{\sqrt{h^2 + k^2} \sqrt{h^2 + k^2}}{\sqrt{h^2 + k^2}} = \sqrt{h^2 + k^2} \rightarrow 0 \end{aligned}$$

□

Remark. To check some $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear we listed two properties:

$$\begin{aligned} T(x + y) &= T(x) + T(y) \\ T(\lambda x) &= \lambda T(x) \end{aligned}$$

we can instead just check:

$$T(\lambda x + y) = \lambda T(x) + T(y)$$

Corollary 1.1. $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ differentiable at $a \in \mathbb{R}^n$

$$(i) \quad D(f + g)(a) = Df(a) + Dg(a)$$

$$(ii) \quad \text{Product rule: } D(f \cdot g)(a) = g(a) \cdot Df(a) + f(a) \cdot Dg(a)$$

$$(iii) \quad \text{Quotient rule: if } g(a) \neq 0, \quad D\left(\frac{f}{g}\right)(a) = \frac{1}{g(a)^2} \cdot (g(a) \cdot Df(a) - f(a) \cdot Dg(a))$$

Proof. For (i):

We can consider the function s from theorem 1.7, $s : \mathbb{R}^2 \rightarrow \mathbb{R}$ $s(x, y) = x + y$, but acting on f and g ie $s(f, g) = f + g$ and $Ds = s$

$$D(f + g)(a) = Ds(f(a), g(a)) \circ D(f, g)(a) = s \circ (Df(a), Dg(a)) = Df(a) + Dg(a)$$

For (ii):

We can consider the function p from theorem 1.8, $p : \mathbb{R}^2 \rightarrow \mathbb{R}$ $p(x, y) = xy$, but acting on f and g ie $p(f, g) = fg$ with $Dp(f, g)(h, k) = fk + gh$

$$D(f \cdot g)(a) = Dp(f, g)(a) \cdot D(f, g)(a) = Dp(f(a), g(a)) \cdot (Df(a), Dg(a)) = f(a) \cdot Dg(a) + g(a) \cdot Df(a)$$

(iii) follows from (ii) □

1.6 Mixed Derivatives

$f : \mathbb{R}^n \rightarrow \mathbb{R}, a \in \mathbb{R}$

$$D_i = \lim_{h \rightarrow 0} \frac{f(a^1, \dots, a^{i-1}, a^i + h, a^{i+1}, \dots, a^n) - f(a)}{h}$$

if $D_i f(x)$ exists for all a in some open set U then we get a function $U \xrightarrow{D_i} \mathbb{R}, x \rightarrow D_i f(x)$ then we can talk about partial derivatives of $D_i f$ eg $D_j(D_i f(x)) = D_{ij} f(x)$

If $D_i f(x)$ exists $\forall x \in U$ this is a function of x and we can consider $D_j(D_i f(x)) = D_{ji} f(x)$

In general $i \neq j$ eg $f(x, y) = x^3 y^5$:

$$\begin{aligned} D_1 f(x, y) &= 3x^2 y^5 & D_2 f(x, y) &= 5x^3 y^4 \\ D_{2,1} f(x, y) &= 15x^2 y^4 & D_{1,2} f(x, y) &= 15x^3 y^4 \end{aligned}$$

Theorem 1.9. If $D_{i,j}$ and $D_{j,i}$ are continuous on an open set containing a then

$$D_{i,j} = D_{j,i}$$

Proof. from homework 5:

First we repeat the well-known proof that, if $g : U \rightarrow \mathbb{R}$ is continuous and $g(p) > 0$, then there exists a neighborhood V of p ($p \in V \subset U$, V open) with

$$q \in V \Rightarrow g(q) > 0$$

Take $\epsilon = g(p)$ in the definition of continuity of g . There there exists a V open with $p \in V$ and

$$q \in V \Rightarrow |g(q) - g(p)| < g(p)$$

Since

$$g(p) - g(q) \leq |g(q) - g(p)| < g(p) \Rightarrow -g(q) < 0 \Leftrightarrow g(q) > 0$$

we get the result. The set V can be taken to contain a closed rectangle $[a, b] \times [c, d]$.

We apply the result to $g = D_{1,2}f - D_{2,1}f$. Assume (by contradiction) that $g(p)$ is not always 0. Then there exists a point p with $g(p) \neq 0$. We can assume that $g(p) > 0$, otherwise consider $-g$. The function g is given to be continuous. We have (using Fubini twice)

$$\begin{aligned} 0 &< \int_{[a,b] \times [c,d]} (D_{1,2}f(x, y) - D_{2,1}f(x, y)) dA \\ &= \int_a^b \left(\int_c^d D_{1,2}f(x, y) dy \right) dx - \int_a^b \left(\int_c^d D_{2,1}f(x, y) dx \right) dy \\ &= \int_a^b (D_1f(x, d) - D_1f(x, c)) dx - \int_c^d (D_2f(b, y) - D_2f(a, y)) dy \\ &= (f(b, d) - f(a, d) - f(b, c) + f(a, c)) - (f(b, d) - f(b, c) - f(a, d) + f(a, c)) = 0 \end{aligned}$$

using the fundamental theorem of calculus 6 times. This is a contradiction, so the mixed partial derivatives are equal on the rectangle. \square

Theorem 1.10. $A \subset \mathbb{R}$ If the max or min of $f : A \rightarrow \mathbb{R}$ occur at a point a in the interior of A and $D_i f(x)$ exists then $Df(a) = 0$

Proof. Consider $h(x) = f(a^1, \dots, a^{i-1}, x^i, a^{i+1}, \dots, a^n)$ x in an open interval around a^i . Since f has a max or min at a , h has a max or min at a^i

$$\frac{dh}{dx}(a^i) = D_i f(a)$$

By analysis 2:

$$\frac{dh}{dx}(a^i) = 0 \Rightarrow Df(a) = 0$$

\square

Note. The converse of Theorem 1.10 is not true, even in one dimension.

1.7 Jacobian

For $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with total derivative $Df(a) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ a linear map. Then the Jacobian $f'(a) \in \mathbb{M}_{m \times n}$ is the unique representation of $Df(a)$ in the standard basis.

Theorem 1.11. *If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at a then $D_j f^i(a)$ exists $\forall i = 1, \dots, m \forall j = 1, \dots, n$ and the jacobian matrix is*

$$f'(a) = (D_j f^i(a))_{j=1, \dots, n}^{i=1, \dots, m}$$

$$f'(a) = \begin{pmatrix} D_1 f^1(a) & D_2 f^1(a) & \cdots & D_n f^1(a) \\ D_1 f^2(a) & D_2 f^2(a) & \cdots & D_n f^2(a) \\ \vdots & \vdots & \ddots & \vdots \\ D_1 f^m(a) & D_2 f^m(a) & \cdots & D_n f^m(a) \end{pmatrix}$$

Where $f(x) = (f^1(x), \dots, f^m(x))$, $f^i : \mathbb{R}^n \rightarrow \mathbb{R}$

Proof. Case $m = 1$

$$\begin{array}{ccc} \mathbb{R} & \xrightarrow{h} & \mathbb{R}^n \\ & \searrow f \circ h & \downarrow f \\ & & \mathbb{R} \end{array}$$

$$h(t) = (a^1, \dots, a^{i-1}, t, a^{i+1}, \dots, a^n) \quad \frac{d(f \circ h)}{dt} \Big|_{t=a^i} = D_i f(a)$$

$$\lim_{t \rightarrow a^i} \frac{(f \circ h)(t) - (f \circ h)(a^i)}{t - a^i} = \lim_{t \rightarrow a^i} \frac{f(a^1, \dots, a^{i-1}, t, a^{i+1}, \dots, a^n) - f(a^1, \dots, a^n)}{t - a^i}$$

h is differentiable because its components are differentiable ie component h^i is either constant a^j where $j \neq i$ or t when $j = i$

$$\begin{aligned} Dh(t) &= (Dh^1(t), \dots, Dh^n(t)) \\ &= (0, \dots, 1, \dots, 0) \end{aligned}$$

$$h'(a^i) = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}_{(m \times 1)}$$

Case $m > 1$

$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$

$$f(x) = (f^1(x), \dots, f^m(x))$$

$$Df(a) = (Df^1(a), \dots, Df^m(a))$$

$$f'(a) = \begin{pmatrix} (f^1)'(a) \\ \vdots \\ (f^m)'(a) \end{pmatrix}_{(m \times n)}$$

$$f'(a) = \begin{pmatrix} D_1 f^1(a) & D_2 f^1(a) & \cdots & D_n f^1(a) \\ D_1 f^2(a) & D_2 f^2(a) & \cdots & D_n f^2(a) \\ \vdots & \vdots & \ddots & \vdots \\ D_1 f^m(a) & D_2 f^m(a) & \cdots & D_n f^m(a) \end{pmatrix}$$

□

Remark. Abuse of notation since if $f : \mathbb{R} \rightarrow \mathbb{R}$

this is a number $\rightarrow \frac{dg(t_0)}{dt} = g'(t_0) \leftarrow$ this is the 1×1 jacobian matrix

Example 1.9.

$$G(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Fix a vector $u \in \mathbb{R}^2$, $u = (u^1, u^2) \neq (0, 0)$, $u^2 \neq 0$ then the directional derivative D_u with $h \in \mathbb{R}$ is:

$$\begin{aligned} D_u G(0, 0) &= \lim_{h \rightarrow 0} \frac{G((0, 0) + hu) - G(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{G(hu^1, hu^2) - 0}{h} \\ &= \lim_{h \rightarrow 0} \frac{(hu^1)^2 (hu^2)}{(hu^1)^4 + (hu^2)^2} \cdot \frac{1}{h} = \lim_{h \rightarrow 0} \frac{h^3 (u^1)^2 u^2}{h(h^4 (u^1)^4 + h^2 (u^2)^2)} \\ &= \lim_{h \rightarrow 0} \frac{(u^1)^2 u^2}{h^2 (u^1)^4 + (u^2)^2} = \frac{(u^1)^2}{u^2} \end{aligned}$$

$u^2 = 0$

$$D_u G(0, 0) = \lim_{h \rightarrow 0} \frac{G(hu^1, h \cdot 0)}{h} = \lim_{h \rightarrow 0} \frac{(\frac{(hu^1)^2 0}{(hu^1)^4 + 0^2})}{h} = 0$$

Theorem 1.12. $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ if $D_j f^i(x)$ exist $\forall x \in U$, U open, $a \in U$, $\forall i = 1, \dots, m$, and $j = 1, \dots, n$ and if $D_j f^i(x)$ continuous at a ie

$$\lim_{x \rightarrow a} (D_j f^i(x)) = D_j f^i(a)$$

then $Df(a)$ exists and f is differentiable at a

Proof. As in the proof of theorem 1.11 It suffices to consider the case $m = 1$, so that $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Then

$$\begin{aligned} f(a + h) - f(a) &= f(a^1 + h^1, a^2, \dots, a^n) && - f(a^1, \dots, a^n) \\ &+ f(a^1 + h^1, a^2 + h^2, a^3, \dots, a^n) && - f(a^1 + h^1, a^2, \dots, a^n) \\ &+ \dots && - \dots \\ &+ f(a^1 + h^1, \dots, a^n + h^n) && - f(a^1 + h^1, \dots, a^{n-1} + h^{n-1}, a^n) \end{aligned}$$

Recal from theorem 1.11 that $D_1 f$ is the derivative of the function h defined by $h(x) = (x, a^2, \dots, a^n)$. Applying the mean-value theorem to h we obtain

$$f(a^1 + h^1, a^2, \dots, a^n) - f(a^1, \dots, a^n) = h^1 \cdot D_1 f(b_1, a^2, \dots, a^n)$$

for some b_1 between a^1 and $a^1 + h^1$. Similarly the i th term in the sum equals

$$h^i \cdot D_i f(a^1 + h^1, \dots, a^{i-1} + h^{i-1}, b_i, a^{i+1}, \dots, a^n) = h^i D_i f(c_i) \quad \text{for some } c_i$$

Then

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \sum_{i=1}^n D_i f(a) \cdot h^i|}{|h|} &= \lim_{h \rightarrow 0} \frac{|\sum_{i=1}^n [D_i f(c_i) - D_i f(a)] \cdot h^i|}{|h|} \\ &\leq \lim_{h \rightarrow 0} \left| \sum_{i=1}^n [D_i f(c_i) - D_i f(a)] \right| \cdot \frac{|h^i|}{|h|} \\ &\leq \lim_{h \rightarrow 0} \left| \sum_{i=1}^n [D_i f(c_i) - D_i f(a)] \right| \\ &= 0 \end{aligned}$$

Since $D_i f$ is continuous at a and as $h \rightarrow 0$, $c^i \rightarrow a^i$. □

Definition 1.7. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ has partial derivatives $D_j f^i \quad \forall x \in U$, U open, $a \in U$ and $D_j f^i$ is continuous at a then we say f is continuously differentiable at a .

Example 1.10. $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, with $x : \mathbb{R} \rightarrow \mathbb{R}, y : \mathbb{R} \rightarrow \mathbb{R}$.

$$\text{Define } g : \mathbb{R} \rightarrow \mathbb{R} \quad g(t) = f(x(t), y(t))$$

$$\begin{aligned} \frac{dg(t_0)}{dt} &= (g'(t_0)) = f'(x(t_0), y(t_0)) \cdot \begin{pmatrix} x'(t_0) \\ y'(t_0) \end{pmatrix} \\ &= \frac{df}{dx}(x(t_0), y(t_0)) \cdot \frac{dx}{dt}(t_0) + \frac{df}{dy}(x(t_0), y(t_0)) \cdot \frac{dy}{dt}(t_0) \\ &= \frac{df}{dx} \cdot \frac{dx}{dt} + \frac{df}{dy} \cdot \frac{dy}{dt} \end{aligned}$$

1.8 Inverse Function Theorem

Lemma 1.13. Let $A \subset \mathbb{R}^n$ be a rectangle with interior A^0 and let $g : A \rightarrow \mathbb{R}^n$ be continuously differentiable. If there exist a constant $M > 0$ such that

$$|D_j g^i(x)| \leq M, \quad x \in A^0, \quad i, j = 1, \dots, n.$$

then

$$|g(x) - g(y)| \leq n^2 M |x - y|, \quad x, y \in A.$$

Proof. Fix $i = 1, \dots, n$. Then

$$\begin{aligned} g^i(y) - g^i(x) &= g^i(y^1, y^2, \dots, y^n) - g^i(x^1, x^2, \dots, x^n) \\ &= g^i(y^1, y^2, \dots, y^n) - g^i(x^1, y^2, \dots, y^n) + g^i(x^1, y^2, \dots, y^n) - g^i(x^1, x^2, \dots, y^n) \\ &\quad + g^i(x^1, x^2, \dots, y^n) - \dots + g^i(x^1, x^2, \dots, y^n) - g^i(x^1, x^2, \dots, x^n) \\ &= \sum_{j=1}^n (g^i(x^1, x^2, \dots, x^{j-1}, y^j, \dots, y^n) - g^i(x^1, x^2, \dots, x^{j-1}, x^j, y^{j+1}, \dots, y^n)) \\ &= \sum_{j=1}^n (y^j - x^j) D_j g^i(z_j^i) \end{aligned}$$

where z_j^i is between y^j and x^j , and we used the mean-value theorem in the interval between y_j and x_j and in the j variable. Using the triangle inequality and $|z^j| \leq |z|$, we get

$$|g^i(y) - g^i(x)| \leq \sum_{j=1}^n |y^j - x^j| M \leq \sum_{j=1}^n |y - x| M = nM|y - x|.$$

Since $|z| \leq \sum_i |z^i|$, finally we get

$$|g(x) - g(y)| \leq \sum_{i=1}^n |g^i(y) - g^i(x)| \leq \sum_{i=1}^n nM|y - x| = n^2 M|y - x|.$$

□

Remark. It is clear that the dimension of the target space enters only in the last line of the calculation. If $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then we get as upper bound $nmM|x - y|$. The inequality is actually not optimal: one can use the Cauchy-Schwarz inequality twice to get a bound $n^{1/2}m^{1/2}M|x - y|$ for $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

Theorem 1.14 (Inverse Function Theorem). *Theorem Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be continuously differentiable on an open set containing a and assume $\det f'(a) \neq 0$. Then there exists an open set V containing a and an open set W containing $f(a)$ such that $f : V \rightarrow W$ is bijective with $f^{-1} : W \rightarrow V$ continuously differentiable and which satisfies:*

$$(f^{-1})'(y) = [f'(f^{-1}(y))]^{-1}, \quad y \in W.$$

Proof. Step 1:

We reduce proving the theorem to the case where actually $f'(a) = I_{nn}$. Call $\lambda = Df(a)$. This is a linear transformation with nonsingular matrix representation $f'(a)$, as $\det f'(a) \neq 0$. Therefore, λ is invertible. The inverse λ^{-1} is also a linear transformation, so $D(\lambda^{-1})(y) = \lambda^{-1}$ for $y \in \mathbb{R}^n$. Both λ and its inverse are continuous as linear transformations. Consider the function $h = \lambda^{-1} \circ f$ defined on an open set containing a .

Then:

$$Dh(a) = D\lambda^{-1}(f(a)) \circ Df(a) = \lambda^{-1} \circ Df(a) = \lambda^{-1} \circ \lambda = Id,$$

by using the chain rule. Here Id is the identity transformation. This gives $h'(a) = I_{n \times n}$, which has determinant 1 $\neq 0$. Let A be the matrix representation of λ^{-1} . (which gives that A^{-1} is the matrix representation of $\lambda = D\lambda$). This is an $n \times n$ matrix with constant entries, i.e. not depending on y . Moreover, h is continuously differentiable, as

$$(D_j h^i(x)) = h'(x) = [\lambda^{-1} \circ Df(x)] = A \cdot f'(x) = A(D_j f^i(x)),$$

with entries depending continuously on x . Therefore, h satisfies the conditions of the inverse function theorem. Suppose that we can prove the conclusion of it for h , i.e. that there exists an open set V containing a and \tilde{W} open containing $h(a) = \lambda^{-1}(f(a))$ such that $h : V \rightarrow \tilde{W}$ is bijective with continuously differentiable inverse h^{-1} . Even more, assume that we have prove the formula for the derivative of the inverse of h :

$$(h^{-1})'(z) = [h'(h^{-1}(z))]^{-1}.$$

Define $W = \lambda(\tilde{W}) = (\lambda^{-1})^{-1}(\tilde{W})$. This is the inverse image of \tilde{W} by λ^{-1} , which is continuous, so it is an open set. Since λ is bijective, $f = \lambda \circ h$ is bijective on V with image $\lambda(\tilde{W}) = W$. Moreover,

$$f^{-1} = h^{-1} \circ \lambda^{-1},$$

which is continuously differentiable as the composition of two such maps. By the chain rule for Jacobian

$$\begin{aligned}(f^{-1})'(y) &= (h^{-1})'(\lambda^{-1}(y)) \cdot (\lambda^{-1})'(y) = [h'(h^{-1}(\lambda^{-1}(y)))]^{-1}A = [h'((\lambda \circ h)^{-1}(y))]^{-1}A = [h'(f^{-1}(y))]^{-1}A \\ &= [A^{-1}h'(f^{-1}(y))]^{-1} = [\lambda'h'(f^{-1}(y))]^{-1} = [(\lambda \circ h)'(f^{-1}(y))]^{-1} = [f'(f^{-1}(y))]^{-1}.\end{aligned}$$

All these imply that it is enough to work with $h = \lambda^{-1} \circ f$. The main property we will use is that $h'(a) = I_{n \times n}$. For simplicity in our notation we call this function f so we can assume that

$$f'(a) = I_{n \times n}.$$

This also means that $\lambda = Df(a) = Id$.

Step 2: The function f cannot take the value $f(a)$ arbitrarily close to a . Suppose that there is a sequence $h_n \in \mathbb{R}^n$ such that $h_n \rightarrow 0$ and $f(a + h_n) = f(a)$. We plug the sequence into the definition of the derivative at a and use that $Df(a) = Id$ to get

$$0 = \lim_{h_n \rightarrow 0} \frac{|f(a + h_n) - f(a) - Df(a)(h_n)|}{|h_n|} = \lim_{h_n \rightarrow 0} \frac{|h_n|}{|h_n|} = 1$$

So this is a contradiction. Therefore, we can find a closed rectangle U containing a such that

$$f(x) \neq f(a), \quad \forall x \in U \setminus \{a\}.$$

Step 3: The determinant is a polynomial expression in the entries of a matrix. If the matrix entries depend continuously on x , the same is true for the determinant of the matrix. So $\det f'(x)$ is a continuous function on an open set containing a . Since $\det f'(a) \neq 0$, by the inertia principle, there exists a small enough (rectangular) neighbourhood of a , which we call U again, such that

$$\det f'(x) \neq 0, \quad x \in U \tag{1}$$

Moreover the partial derivatives $D_j f^i(x)$ are continuous and $D_j f^i(a) = \delta_{ij}$, as $Df(a) = Id$. So, for x close enough to a we have

$$|D_j f^i(x) - \delta_{ij}| < \frac{1}{2n^2}, \quad i, j = 1, \dots, n, \quad x \in U \tag{2}$$

We assumed again that the neighbourhood is U

Step 4: Constructing a contraction map and showing that f is injective in appropriate small neighbourhood. Now we define the function

$$g(x) = f(x) - x$$

and apply the Lemma to this function for the closed rectangle U . We notice that $D_j g^i(x) = D_j f^i(x) - \delta_{ij}$, as we know the partial derivatives of the identity function x . We deduce that

$$|g(x_1) - g(x_2)| \leq n^2 \frac{1}{2n^2} |x_1 - x_2| = \frac{1}{2} |x_1 - x_2| \tag{3}$$

The choice of the neighbourhood in (2) so that the constant $1/(2n^2)$ appears on the right is motivated with the desire to get g as a contraction map (with constant $1/2$) as we see in (3). Now the triangle inequality in the form $|a| - |b| \leq |a - b|$ gives

$$|x_1 - x_2| - |f(x_1) - f(x_2)| \leq |(x_1 - x_2) - (f(x_1) - f(x_2))| = |-g(x_1) + g(x_2)| < \frac{1}{2} |x_1 - x_2|$$

$$\Rightarrow |x_1 - x_2| - \frac{1}{2}|x_1 - x_2| < |f(x_1) - f(x_2)| \Rightarrow \frac{1}{2}|x_1 - x_2| < |f(x_1) - f(x_2)| \quad (4)$$

Here x_1, x_2 are in U . We immediately see that on U the function f is injective:

$$f(x_1) = f(x_2) \Rightarrow |x_1 - x_2| = 0 \Rightarrow x_1 = x_2.$$

We still have not determined the neighbourhoods W of $f(a)$ and V of a .

Step 5: Determination of the minimum distance of $f(a)$ to the image of the boundary of U and definition of W .

We have assumed that on the closed rectangle U we have $f(x) \neq f(a)$ for $x \neq a$. This is definitely true on the boundary of U , denoted ∂U , which is a closed and bounded set, i.e. compact. The function $m(x) = |f(x) - f(a)|$ is continuous on a neighbourhood of ∂U and nonzero on it. It achieves a minimum value on ∂U (an advanced argument from Real Analysis is that the image of a compact set is compact, so that $m(\partial U)$ is compact, which means closed and bounded. Such a set has a maximum and minimum). The minimum value cannot be zero, say

$$\min_{x \in \partial U} m(x) = \min_{x \in \partial U} |f(x) - f(a)| > 0.$$

Now define

$$W = \{y \in \mathbb{R}^n, |y - f(a)| < \delta/2\}.$$

Step 6: Comparison of $|y - f(x)|$ with $|y - f(a)|$ for $x \in \partial U$, and $y \in W$. We have

$$\begin{aligned} |f(x) - f(a)| \geq \delta, \quad |y - f(a)| \leq \delta/2 &\Rightarrow -|y - f(x)| + \delta \leq -|y - f(x)| + |f(x) - f(a)| \leq |y - f(a)| < \delta/2 \\ &\Rightarrow \delta/2 = \delta - \delta/2 < |y - f(x)| \Rightarrow |y - f(a)| < \delta/2 < |y - f(x)|. \end{aligned}$$

Step 7: Show that for $y_0 \in W$ there exists a unique $x_0 \in U^0$ such that $f(x_0) = y_0$. The uniqueness is obvious from the fact that f is injective on U . The construction of such an x_0 is tricky. We define another function on U by

$$g(x) = |f(x) - y_0|^2 = \sum_{i=1}^n (f^i(x) - y_0^i)^2.$$

This function is continuously differentiable, as it is a sum of the squares of the components. On the compact set U the function g achieves its minimum, say at x_0 , i.e. $g(x_0) \leq g(x)$ for $x \in U$. We claim that x_0 is the desired point with $f(x_0) = y_0$. First we see that x_0 is in the interior of the set U . On the boundary of U the function $g(x)$ has values $> \delta/2$, by Step 6, while $g(a) < \delta/2$. So the minimum is not achieved on the boundary of U . Therefore, it is achieved in an interior point. This point has to be a critical point of g , i.e. $D_j g(x_0) = 0$, $j = 1, \dots, n$. We calculate them to be

$$2 \sum_{i=1}^n (f^i(x_0) - y_0^i) D_j f^i(x_0) = 0, \quad j = 1, \dots, n.$$

This is a homogeneous system of linear equations with unknowns $f^i(x_0) - y_0^i$ and coefficients $D_j f^i(x_0)$. The determinant of the coefficients of the system is nonzero, as $x_0 \in U$. The system has a unique solution, and this solution is the zero vector, i.e

$$0 = f^i(x_0) - y_0^i, \quad i = 1, \dots, n \Rightarrow f(x_0) = y_0.$$

Step 8: We define V and Show that $f : V \rightarrow W$ is bijective and continuous. We define $V = U^0 \cap f^{-1}(W)$. Clearly $f : V \rightarrow W$ is bijective. Moreover, V is open as the intersection of the

open set U^0 and the open set $f^{-1}(W)$, which is open as the inverse image of an open set W by the continuous function f . We now rewrite (4) as

$$|x_1 - x_2| < 2|f(x_1) - f(x_2)| \Leftrightarrow |f^{-1}(y_1) - f^{-1}(y_2)| < 2|y_1 - y_2| \quad (5)$$

with $y_1 = f(x_1)$ and $y_2 = f(x_2)$, $y_i \in W$. This shows that f^{-1} is a Lipschitz function with constant 2, so that it is continuous. Alternatively, choose $\delta = \epsilon/2$ in the definition of continuity.

Step 9: Show that f^{-1} is differentiable. Let $\mu = Df(x_1)$. Since $f^{-1} \circ f = Id$, the chain rule gives the only possible choice for $Df^{-1}(y_1) = \mu^{-1}$. Here $f(x_1) = y_1$ and later $f(x) = y$. By the definition of the derivative we have

$$f(x) - f(x_1) = \mu(x - x_1) + \phi(x - x_1), \quad \lim_{x \rightarrow x_1} \frac{|\phi(x - x_1)|}{|x - x_1|} = 0.$$

We apply to the equation the linear transformation μ^{-1} to get

$$\begin{aligned} \mu^{-1}(y - y_1) &= x - x_1 + \mu^{-1}(\phi(x - x_1)) \Rightarrow x - x_1 - \mu^{-1}(y - y_1) = \mu^{-1}(\phi(x - x_1)) \\ &\Rightarrow f^{-1}(y) - f^{-1}(y_1) - \mu^{-1}(y - y_1) = -\mu^{-1}(\phi(x - x_1)). \end{aligned}$$

By the definition of the derivative of f^{-1} at y_1 we need to show that

$$\lim_{y \rightarrow Y_1} \frac{|-\mu^{-1}(\phi(x - x_1))|}{|y - y_1|} = 0 \quad (6)$$

Since μ^{-1} is a linear transformation, we have seen that it is a bounded linear operator, i.e. there exists a constant \tilde{M} with

$$|\mu^{-1}(y)| \leq \tilde{M}|y|, \quad \forall y \in \mathbb{R}^n.$$

Since

$$\frac{|-\mu^{-1}(\phi(x - x_1))|}{|y - y_1|} \leq \frac{\tilde{M}|\phi(x - x_1)|}{|y - y_1|}$$

by the sandwich theorem it is enough to prove that

$$\lim_{y \rightarrow Y_1} \frac{|\phi(x - x_1)|}{|y - y_1|} = 0$$

We have

$$\frac{|\phi(x - x_1)|}{|y - y_1|} = \frac{|\phi(x - x_1)|}{|x - x_1|} \frac{|x - x_1|}{|y - y_1|} \leq \frac{|\phi(x - x_1)|}{|x - x_1|} \cdot 2,$$

by (5). Moreover, $y \rightarrow y_1$ iff $x \rightarrow x_1$ as f is continuous at x_1 and f^{-1} is continuous at y_1 . We know that

$$\lim_{x \rightarrow x_1} \frac{|(\phi(x - x_1))|}{|x - x_1|} = 0$$

This suffices to prove (6)

Step 10: The partial derivatives $D_j(f^{-1})^i(y)$ are continuous. We know that the Jacobian of $f^{-1}(y)$ is

$$(f^{-1})'(y) = (D_j(f^{-1})^i(y)) = [f'(f^{-1}(y))]^{-1} = (D_j f^i(x))^{-1}.$$

The inverse of the matrix $(D_j f^i(x))$ can be calculated as a quotient of two $n \times n$ determinants with entries among $D_j f^i(x)$. The denominator is the determinant of the Jacobian at x , which is nonzero for $x \in U$. The whole expression depends continuously on $x \in V$. As f^{-1} is continuous, the inverse matrix depends continuously on $y \in W$. The individual entries are the partial derivatives of f^{-1} . \square

Example 1.11. $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$(z, w) = f(x, y) = (xy, x^2 + y^2), \quad z = xy, \quad w = x^2 + y^2$$

$$f'(x, y) = \begin{pmatrix} y & x \\ 2x & 2y \end{pmatrix}$$

$$\det f'(x, y) = 2y^2 - 2x^2 = 2(y + x)(y - x)$$

$$\det f'(x, y) \neq 0 \Leftrightarrow x \neq \pm y$$

Solving:

$$y = \frac{z}{x}$$

$$\therefore w = x^2 + \frac{z^2}{x^2}$$

$$\therefore wx^2 = x^4 + z^2$$

$$\Rightarrow x^4 - wx^2 + z^2 = 0$$

Let $t = x^2 \therefore t^2 - wt + z^2 = 0$ So

$$t = \frac{w \pm \sqrt{w^2 - 4z^2}}{2}$$

$$x = \pm \sqrt{\frac{w \pm \sqrt{w^2 - 4z^2}}{2}}$$

And

$$y = \frac{z}{\pm \sqrt{\frac{w \pm \sqrt{w^2 - 4z^2}}{2}}}$$

You should be able to differentiate if $w^2 - 4z^2 \neq 0 \Leftrightarrow$ if $y \neq \pm x$

$$\begin{bmatrix} \frac{dx}{dz} & \frac{dx}{dw} \\ \frac{dy}{dz} & \frac{dy}{dw} \end{bmatrix} = (f^{-1})'(z, w) = \begin{pmatrix} y & x \\ 2x & 2y \end{pmatrix}^{-1} = \frac{1}{2(y^2 - x^2)} \begin{pmatrix} 2y & -x \\ -2x & y \end{pmatrix}$$

$$\Rightarrow \frac{dx}{dz} = \frac{2y}{2(y^2 - x^2)}$$

$$\Rightarrow \frac{dx}{dw} = \frac{-x}{2(y^2 - x^2)}$$

When we have $z = xy$, $w = x^2 + y^2$ along the lines $y = \pm x$ the circle meets the hyperbola tangentially so we cannot invert.

1.9 Implicit Function Theorem

Example 1.12.


$$x^2 + y^2 = 1, \quad y = g(x)$$

$$2x + 2y \frac{dy}{dx} = 0, \quad \frac{dy}{dx} = \frac{dg}{dx} = \frac{-x}{y}, \quad y \neq 0$$

Example 1.13.

$$y^2 + xz + z^2 - e^z - 4 = 0 \quad (\text{impossible to solve for } z)$$

$$\text{set } F(x, y, z) = y^2 + xz + z^2 - e^z - 4, \quad F(x, y, g(x, y)) = 0$$



Differentiate in x :

$$\begin{aligned} \frac{d}{dx} F(x, y, g(x, y)) &= \frac{dF}{dx} \frac{dx}{dx} + \frac{dF}{dy} \frac{dy}{dx} + \frac{dF}{dz} \frac{dz}{dx} \\ &= \frac{dF}{dx} + \frac{dF}{dz} \frac{dg}{dx} = 0 \\ \frac{dg}{dx} &= -\frac{\frac{dF}{dx}}{\frac{dF}{dz}} = -\frac{z}{x^2 + 2z - e^z} \\ \frac{dF}{dy} &= 0 \xrightarrow[\text{rule}]{\text{chain}} \frac{dF}{dy} + \frac{dF}{dz} \frac{dz}{dy} \Rightarrow \frac{dg}{dy} = -\frac{\frac{dF}{dy}}{\frac{dF}{dz}} = -\frac{2y}{x^2 + 2z - e^z} \end{aligned}$$

the point $(0, e, 2)$ satisfies $F(x, y, z) = 0$

$$e^2 + 0 \cdot 2 + 2^2 - e^2 - 4 = 0$$

$$\begin{aligned} \left. \frac{dg}{dx} \right|_{(0,e)} &= -\frac{z}{x^2 + 2z - e^z} = -\frac{2}{0 + 2 \cdot 2 - e^2} \\ \left. \frac{dg}{dy} \right|_{(0,e)} &= -\frac{2y}{x^2 + 2z - e^z} = -\frac{2e}{0 + 2 \cdot 2 - e^2} \end{aligned}$$

valid for $\frac{dF}{dz} \neq 0$

General situation: m equations with m unknowns y^1, \dots, y^m

$$\begin{aligned} f^1(x^1, \dots, x^n, y^1, \dots, y^m) &= 0 && \text{depends on } n \text{ parameters: } x^1, \dots, x^n \\ f^2(x^1, \dots, x^n, y^1, \dots, y^m) &= 0 && \text{Try to solve for: } y^1, \dots, y^m \\ \vdots &&& \vdots \\ f^m(x^1, \dots, x^n, y^1, \dots, y^m) &= 0 \end{aligned}$$

$$x = (x^1, \dots, x^n), \quad y = (y^1, \dots, y^m)$$

So we have:

$$\begin{aligned} f^1(x, y) &= 0 \\ f^2(x, y) &= 0 \\ &\vdots \\ f^m(x, y) &= 0 \end{aligned}$$

Define $f(x, y) = (f^1(x, y), \dots, f^m(x, y)) = \underbrace{0}_{\text{vector}} = \underbrace{(0, \dots, 0)}_m$

Let $a \in \mathbb{R}^n$, $b \in \mathbb{R}^m$ such that $f(a, b) = 0$ when we can find for each (x^1, \dots, x^n) near $a = (a^1, \dots, a^n)$ a unique $y = (y^1, \dots, y^m)$ near $b = (b^1, \dots, b^m)$ such that: $f(x, y) = 0$, $f(x^1, \dots, x^n, y^1, \dots, y^m) = 0$

Theorem 1.15 (Implicit Function Theorem). $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ continuously differentiable on an open set containing (a, b) , $a \in \mathbb{R}^n$, $b \in \mathbb{R}^m$. moreover $f(a, b) = 0$ consider the matrix

$$M = (D_{j+n} f^i(a, b))_{i=1, \dots, m}^{j=1, \dots, m}$$

assume $\det M \neq 0$. Then there exist two open sets $A \subset \mathbb{R}^n$, $B \subset \mathbb{R}^m$, $a \in A$, $b \in B$. such that $\forall x \in A, \exists$ unique $g(x) \in B$ such that $f(x, g(x)) = 0$ Moreover $g : A \rightarrow B$ is differentiable.

Proof. Increase the dimension of the target. Define $F : \underbrace{U}_{\in \mathbb{R}^n \times \mathbb{R}^m} \rightarrow \mathbb{R}^n \times \mathbb{R}^m$

$$F(x^1, \dots, x^n, y^1, \dots, y^m) = (x^1, \dots, x^n, f^1(x, y), \dots, f^m(x, y))$$

$$F(x, y) = (x, f(x, y))$$

F is continuously differentiable because x^1, \dots, x^n are continuously differentiable and $f^1(x, y), \dots, f^m(x, y)$ are continuously differentiable (because $f(x, y)$ is continuously differentiable)

$$F(a, b) = (a, f(a, b)) = (a, 0)$$

$$F'(a, b) = \left(\begin{array}{cccc|ccc} 1 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 1 & 0 & \cdots & 0 \\ \hline \frac{df^1}{dx^1} & \frac{df^1}{dx^2} & \cdots & \frac{df^1}{dx^n} & \frac{df^1}{dy^1} & \cdots & \frac{df^1}{dy^m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & & \vdots \\ \frac{df^m}{dx^1} & \frac{df^m}{dx^2} & \cdots & \frac{df^m}{dx^n} & \frac{df^m}{dy^1} & \cdots & \frac{df^m}{dy^m} \end{array} \right)$$

$$F'(a, b) = \left(\begin{array}{c|c} I_{n \times n} & 0_{n \times m} \\ \hline * \text{ Some } m \times n & M_{m \times m} \\ \text{matrix} & \end{array} \right)$$

$\det M \neq 0$ (reducing from top left entry).

By the inverse function theorem, \exists an open set W containing $F(a, b) = (a, 0)$ and an open set containing $(a, 0)$ which I can take to be a rectangle $A \times B$, $a \in A$, $b \in B$, A open in \mathbb{R}^n , B open in \mathbb{R}^m .

$F : A \times B \rightarrow W$ is bijective

$$\exists h = F^{-1} : W \rightarrow A \times B \text{ such that } F \cdot h = id$$

h is continuously differentiable.

$$\begin{aligned} F(x^1, \dots, x^n, y^1, \dots, y^m) &= (x^1, \dots, x^n, f^1(x, y), \dots, f^m(x, y)) \\ F(x, y) &= (x, f(x, y)) \\ F \text{ is continuously differentiable because } x^1, \dots, x^n &\text{ are continuously differentiable} \end{aligned}$$

h must have the form: $h(x, y) = (x, k(x, y))$ for some function $k : W \rightarrow B$, $B \subset \mathbb{R}^m$, k continuously differentiable.

$$F(h(x, y)) = (x, f(x, k(x, y))) = (x, y)$$

$$f(x, k(x, y)) = y$$

Set $y = 0$

$$f(x, k(x, 0)) = 0$$

The solution is $g(x) = k(x, 0)$ (solution to $f(x, y) = 0$). □

Theorem 1.16. Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^p$ be a continuously differentiable function in an open set containing a and assume that $p \leq n$. If $g(a) = 0$ and the rank of the $p \times n$ matrix

$$(D_j g^i(a))_{i=1, \dots, p, j=1, \dots, n}$$

be equal to p . Then there exists an open set $A \subset \mathbb{R}^n$ and a differentiable function $h : A \rightarrow \mathbb{R}^n$ which is bijective onto an open set V and h^{-1} is differentiable and

$$(g \circ h)(x^1, x^2, \dots, x^n) = (x^{n-p+1}, x^{n-p+2}, \dots, x^n)$$

Proof. We can consider the function g as $g : \mathbb{R}^{n-p} \times \mathbb{R}^p \rightarrow \mathbb{R}^p$. The 'easy' case is as follows:

If the $p \times n$ matrix above is such that the last p columns give a matrix M with $\det(M) \neq 0$, then we are exactly in the situation of the Implicit Function Theorem as worked out above. The notation has only slightly changed: $x^{n-p+1} = y^1, x^{n-p+2} = y^2, \dots, x^n = y^p, p = m, g = f$. We have found h with $h(x, y) = (x, k(x, y))$ and

$$(f \circ h)(x, y) = f(h(x, y)) = f(x, k(x, y)) = y,$$

and in our notation

$$(g \circ h)(x^1, x^2, \dots, x^n) = (x^{n-p+1}, x^{n-p+2}, \dots, x^n).$$

In general we cannot assume that the last columns of the matrix give nonzero determinant. We know from Linear Algebra that there will be some p columns with this property. Let these columns be j^1, j^2, \dots, j^p with

$$M = (D_{j_k} g^i(a))_{i=1, \dots, p, k=1, \dots, p}, \quad \det(M) \neq 0.$$

We rearrange the variables as follows: Let $m : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be defined by (put the variables with superscript j_k , $k = 1, 2, \dots, p$ in the last entries and order in whatever way you want the other variables)

$$m(x^1, x^2, \dots, x^n) = (\dots, x^{j^1}, x^{j^2}, \dots, x^{j^p}).$$

Then $g \circ m$ is a function of the type discussed theorem 1.15, so we can find a function $s : A \rightarrow \mathbb{R}^n$ which is bijective onto an open set V and s^{-1} is differentiable and

$$((g \circ m) \circ s)(x^1, x^2, \dots, x^n) = (x^{n-p+1}, x^{n-p+2}, \dots, x^n).$$

Then use $h = m \circ s$. □

Example 1.14.

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad f(x, y) = (xy, x^2 + y^2) = (z, w)$$

$$\begin{pmatrix} \frac{dx}{dz} & \frac{dx}{dw} \\ \frac{dy}{dz} & \frac{dy}{dw} \end{pmatrix} = \begin{pmatrix} \frac{dz}{dx} & \frac{dz}{dy} \\ \frac{dw}{dx} & \frac{dw}{dy} \end{pmatrix}^{-1}$$

$$z = xy, \quad y = \frac{z}{w}$$

$$w = x^2 + y^2 = x^2 + \frac{z^2}{x^2}$$

$$wx^2 = x^4 + z^2$$

$$x^4 - wx^2 + z^2 \quad (*)$$

$$x = g(z, w)$$

Use implicit differentiation on (*) with respect to z :

$$4x^3 \frac{dx}{dz} - w \cdot 2x \frac{dx}{dz} + 2z = 0$$

$$\frac{dx}{dz} (4x^3 - 2xw) = -2z$$

$$\frac{dx}{dz} = \frac{-2z}{4x^3 - 2xw} = \frac{-z}{x(2x^2 - w)} = \frac{-y}{2x^2 - w}$$

Valid for

$$2x^2 - w \neq 0$$

$$2x^2 - (x^2 + y^2) \neq 0$$

$$x^2 - y^2 \neq 0$$

$$\Leftrightarrow f'(x, y) \neq 0$$

$$\left. \begin{array}{ll} f(x, y) = 0 & f(a, b) = 0 \\ f(x, g(x)) = 0 & \text{solve implicitly for } y \\ x \in \mathbb{R}^n, y \in \mathbb{R}^m & g : \mathbb{R}^n \rightarrow \mathbb{R}^m \\ f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m & \end{array} \right\} \quad \text{Set up of implicit function theorem}$$

$$i = 1, \dots, m \quad f^i(x^1, \dots, x^n, g^1(x^1, \dots, x^n), \dots, g^m(x^1, \dots, x^n)) = 0$$

how to compute $D_j g^i$?

$$D_j g^i(\dots) = 0$$

$$D_1 f^i \cancel{\frac{dx^1}{dx^j}} + \cancel{\dots} + D_j f^i \underbrace{\frac{dx^j}{dx^j}}_{=1} + \cancel{\dots} + D_n f^i \cancel{\frac{dx^n}{dx^j}} + D_{n+1} f^i \frac{dg^1}{dx^j} + \dots + D_{n+m} f^i \frac{dg^m}{dx^j} = 0$$

$$\underbrace{D_{n+1} f^i \frac{dg^1}{dx^j} + \dots + D_{n+m} f^i \frac{dg^m}{dx^j}}_{m \text{ unknowns}} = -D_j f^i \frac{dx^j}{dx^j}$$

Check det of coefficients is $\neq 0$

$$\begin{bmatrix} D_{n+1} f^1 & \dots & D_{n+m} f^1 \\ \vdots & & \vdots \\ D_{n+1} f^m & \dots & D_{n+m} f^m \end{bmatrix} = M$$

2 Integration

2.1 Multiple integrals

$f : A \rightarrow \mathbb{R}$, A is a rectangle in \mathbb{R}^n $A = [a_1, b_1] \times \cdots \times [a_n, b_n]$

Recall a partition \mathcal{P} of $[a, b]$ is a collection of points: t_0, \dots, t_k with $a = t_0 < t_1 < \cdots < t_k = b$

A Partition of a rectangle $[a_1, b_1] \times \cdots \times [a_k, b_k]$ is a collection $\mathcal{P} = (P_1, \dots, P_n)$ where P_i is a partition of $[a_i, b_i]$, $i = 1, \dots, n$ Subrectangles $[s_{j-1}, s_j] \times [t_{m-1}, t_m]$ Let f be bounded on the rectangle $[a_1, b_1] \times \cdots \times [a_k, b_k]$

Definition 2.1. Let f be bounded on the rectangle $[a_1, b_1] \times \cdots \times [a_k, b_k]$ and let S be subrectangle of the partition \mathcal{P}

$$m_S(f) = \inf_{x \in S} f(x), \quad M_S(f) = \sup_{x \in S} f(x)$$

Lower Riemann sum:

$$\mathcal{L}(f, \mathcal{P}) = \sum_S m_S(f) \cdot v(S)$$

where $v(s)$ is the volume of the subrectangle

$$S = [s_{l-1}, s_l] \times [t_{j-1}, t_j] \times \cdots \times [r_{k-1}, r_k]$$

$$v(S) = (s_l - s_{l-1}) \cdot (t_j - t_{j-1}) \cdots (r_k - r_{k-1})$$

Upper Riemann sum:

$$\mathcal{U}(f, \mathcal{P}) = \sum_S M_S(f) \cdot v(S)$$

$$\mathcal{L}(f, \mathcal{P}) \leq \mathcal{U}(f, \mathcal{P})$$

Refinement: A refinement \mathcal{P}' of the partition \mathcal{P} is as follows. Given S a subrectangle of \mathcal{P}' , I can find a subrectangle T of \mathcal{P} such that $S \subset T$ and $T = \cup_{S \subset T} S$, S for \mathcal{P}'

Lemma 2.1. if \mathcal{P}' is a refinement of \mathcal{P} , then:

$$\mathcal{L}(f, \mathcal{P}) \leq \mathcal{L}(f, \mathcal{P}') \tag{1}$$

$$\mathcal{U}(f, \mathcal{P}) \geq \mathcal{U}(f, \mathcal{P}') \tag{2}$$

Proof. of (1)

Let S be a subrectangle of \mathcal{P}' and T a subrectangle of \mathcal{P} such that $S \subset T$ and

$$\begin{aligned} m_S(f) &\geq m_T(f) \\ m_S(f)v(S) &\geq m_T(f)v(S) \end{aligned}$$

now sum over all $S \subset T$, S for \mathcal{P}'

$$\begin{aligned} \sum_{S \subset T} m_S(f)v(S) &\geq \sum_{S \subset T} m_T(f)v(S) = m_T(f)v(T) \\ \mathcal{L}(f, \mathcal{P}') &= \sum_T \sum_{S \subset T} m_S(f)v(S) \geq \sum_T m_T(f)v(T) = \mathcal{L}(f, \mathcal{P}) \end{aligned}$$

□

Lemma 2.2. For any two partitions \mathcal{P} and \mathcal{P}' we have:

$$\mathcal{L}(f, \mathcal{P}) \leq \mathcal{U}(f, \mathcal{P}')$$

Proof. Take \mathcal{P}'' a refinement of \mathcal{P} and \mathcal{P}' :

$$\mathcal{L}(f, \mathcal{P}) \leq \mathcal{L}(f, \mathcal{P}'') \leq \mathcal{U}(f, \mathcal{P}'') \leq \mathcal{U}(f, \mathcal{P}')$$

□

Definition 2.2.

The lower Riemann integral

$$\int_{A-} f = \sup_{\mathcal{P}} \mathcal{L}(f, \mathcal{P}), \quad (\mathcal{P} \text{ partition of rectangle } A)$$

The upper Riemann integral

$$\int_A^{\bar{}} f = \inf_{\mathcal{P}} \mathcal{U}(f, \mathcal{P})$$

f is called integrable if

$$\int_{A-} f = \int_A^{\bar{}} f \quad \text{and} \quad \int_A f = \int_{A-} f = \int_A^{\bar{}} f$$

Theorem 2.3 (Riemann's Integrability Criterion). f is integrable over the rectangle $A \Leftrightarrow \forall \epsilon > 0, \exists$ a partition \mathcal{P} of A such that

$$\mathcal{U}(f, \mathcal{P}) - \mathcal{L}(f, \mathcal{P}) < \epsilon$$

Proof. (\Rightarrow)

$$\begin{aligned} \inf_{\mathcal{P}} (\mathcal{U}(f, \mathcal{P}) - \mathcal{L}(f, \mathcal{P})) &= 0 \\ \Leftrightarrow \inf_{\mathcal{P}} \mathcal{U}(f, \mathcal{P}) - \sup_{\mathcal{P}} \mathcal{L}(f, \mathcal{P}) &= 0 \\ \Leftrightarrow \int_{A-} f &= \int_A^{\bar{}} f \end{aligned}$$

(\Leftarrow)

Assume $\int_{A-} f = \int_A^{\bar{}} f$, fix $\epsilon > 0$

$$\text{Since } \int_{A-} f = \sup_{\mathcal{P}} \mathcal{L}(f, \mathcal{P}), \quad \text{so } \exists \mathcal{P}' \text{ s.t. } \int_{A-} f - \frac{\epsilon}{2} < \mathcal{L}(f, \mathcal{P}')$$

$$\text{Since } \int_A^{\bar{}} f = \inf_{\mathcal{P}} \mathcal{U}(f, \mathcal{P}), \quad \text{so } \exists \mathcal{P}' \text{ s.t. } \int_A^{\bar{}} f + \frac{\epsilon}{2} > \mathcal{U}(f, \mathcal{P}')$$

Take \mathcal{P}'' a common refinement of \mathcal{P} and \mathcal{P}'

$$\int_A^{\bar{}} f + \frac{\epsilon}{2} > \mathcal{U}(f, \mathcal{P}'') \geq \mathcal{L}(f, \mathcal{P}'') > \int_{A-} f - \frac{\epsilon}{2}$$

So

$$\mathcal{U}(f, \mathcal{P}'') - \mathcal{L}(f, \mathcal{P}'') < \left(\int_A^{\bar{}} f + \frac{\epsilon}{2} \right) - \left(\int_{A-} f - \frac{\epsilon}{2} \right) = \epsilon$$

□

Example 2.1. *Non-Riemann integrable function $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$*

$$f(x, y) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

$$\begin{aligned} m_S(f) &= 0 & M_S(f) &= 1 \\ \mathcal{L}(f, \mathcal{P}) &= 0 & \mathcal{U}(f, \mathcal{P}) &= 1 \end{aligned}$$

Definition 2.3. *If $C \subset \mathbb{R}^n$, define the characteristic function of C to be*

$$X_C(x) = \begin{cases} 1 & \text{if } x \in C \\ 0 & \text{if } x \notin C \end{cases}$$

If f is bounded on \bar{C} and C is contained in a rectangle A , we define

$$\int_C f = \int_A f X_C$$

$f : [a, b] \times [c, d] \rightarrow \mathbb{R}$

Fix x and consider $g_x : [c, d] \rightarrow \mathbb{R}$

$$g_x(y) = f(x, y)$$

$$I(x) = \int_c^d g_x dy = \int_c^d f(x, y) dy$$

$$\int_a^b I(x) dx = \int_a^b \left(\int_c^d f(x, y) dy \right) dx \tag{1}$$

Fix y and define $h_y : [a, b] \rightarrow \mathbb{R}$

$$h_y(x) = f(x, y)$$

$$J(y) = \int_a^b h_y dx = \int_a^b f(x, y) dx$$

$$\int_c^d J(y) dy = \int_c^d \left(\int_a^b f(x, y) dx \right) dy \tag{2}$$

$$(1) = (2)$$

2.2 Fubini's theorem

Theorem 2.4 (Fubini). *Let A be a rectangle in \mathbb{R}^n and let B be a rectangle in \mathbb{R}^m . $f : A \times B \rightarrow \mathbb{R}$ is intergrable. define:*

$$g_x : B \rightarrow \mathbb{R} \quad \text{by} \quad g_x = f(x, y), \quad \forall y \in B, \forall x \in A$$

and let:

$$\left. \begin{aligned} \mathfrak{L}(x) &= \int_{B-} g_x = \int_{B-} f(x, y) dy \\ \mathfrak{U}(x) &= \int_{\bar{B}} g_x = \int_{\bar{B}} f(x, y) dy \end{aligned} \right\} \quad \text{exists } \forall x \in A$$

Then $\mathfrak{L}(x)$ and \mathfrak{U} are intergrable over A , and:

$$\int_A \mathfrak{L}(x) dx = \int_A \left(\int_{B-} f(x, y) dy \right) dx = \int_A \left(\int_{\bar{B}} f(x, y) dy \right) dx = \int_A \mathfrak{U}(x) dx = \int_{A \times B} f$$

Proof. Let \mathcal{P}_A be a partition of A , \mathcal{P}_B be a partition of B . Let S_A a subrectangle of A , S_B a subrectangle of B . Then the rectangles $S_A \times S_B$ give a partition \mathcal{P} of $A \times B$.

We will prove:

$$\mathcal{L}(f, \mathcal{P}) \underset{(1)}{\leq} \mathcal{L}(\mathfrak{L}, \mathcal{P}_A) \underset{(2)}{\leq} \mathcal{U}(\mathfrak{L}, \mathcal{P}_A) \underset{(3)}{\leq} \mathcal{U}(\mathfrak{U}, \mathcal{P}_A) \underset{(4)}{\leq} \mathcal{U}(f, \mathcal{P})$$

Since f is integrable over $A \times B$, given $\epsilon > 0$ Riemann's integrability criterion given a partition \mathcal{P} of $A \times B$, such that: $\mathcal{U}(f, \mathcal{P}) - \mathcal{L}(f, \mathcal{P}) < \epsilon$. Then \mathcal{P} defines \mathcal{P}_A , \mathcal{P}_B partitions of A , B respectively. By the inequality above: $\mathcal{U}(\mathfrak{L}, \mathcal{P}_A) - \mathcal{L}(\mathfrak{L}, \mathcal{P}_A) < \epsilon$. By reimann's integrability criterion, \mathcal{L} is integrable over A , since:

$$\sup_{\mathcal{P}} \mathcal{L}(f, \mathcal{P}) = \inf_{\mathcal{P}} \mathcal{U}(f, \mathcal{P}) = \int_{A \times B} f \quad \Rightarrow \quad \int_A \mathfrak{L}(x) dx = \sup_{\mathcal{P}_A} \mathcal{L}(\mathfrak{L}, \mathcal{P}_A) = \inf_{\mathcal{P}_A} \mathcal{U}(\mathfrak{L}, \mathcal{P}_A) = \int_{A \times B} f$$

Works similarly with $\mathfrak{U}(x)$.

Side remark:

$$\begin{aligned} \mathcal{L}(f, \mathcal{P}) &\leq \mathcal{L}(\mathfrak{L}, \mathcal{P}_A) \\ \sup_{\mathcal{P}} \mathcal{L}(f, \mathcal{P}) &\leq \sup_{\mathcal{P}_A} \mathcal{L}(\mathfrak{L}, \mathcal{P}_A) \\ \mathcal{U}(\mathfrak{L}, \mathcal{P}_A) &\leq \mathcal{U}(f, \mathcal{P}) \\ \inf_{\mathcal{P}_A} \mathcal{U}(\mathfrak{L}, \mathcal{P}_A) &\leq \inf_{\mathcal{P}} \mathcal{L}(f, \mathcal{P}) \end{aligned}$$

$$(2) \quad \mathcal{L}(\mathfrak{L}, \mathcal{P}_A) \leq \mathcal{U}(\mathfrak{L}, \mathcal{P}_A)$$

always true for a function \mathfrak{L} , partition \mathcal{P}_A

$$(3) \quad \mathcal{U}(\mathfrak{L}, \mathcal{P}_A) \leq \mathcal{U}(\mathfrak{U}, \mathcal{P}_A)$$

$$\begin{aligned} \mathfrak{L}(x) &= \int_{B-} f(x, y) dy, \quad \mathfrak{U}(x) = \int_B f(x, y) dy \Rightarrow \mathfrak{L}(x) \leq \mathfrak{U}(x) \\ &\Rightarrow \mathcal{U}(\mathfrak{L}(x), \mathcal{P}_A) \leq \mathcal{U}(\mathfrak{U}(x), \mathcal{P}_A) \end{aligned}$$

(4) is proved similarly to (1) so we only prove (1).

$$\mathcal{L}(f, \mathcal{P}) = \sum_S m_s(f) v(S) = \sum_{S_A, S_B} m_{S_A \times S_B}(f) v(S_A \times S_B) = \sum_{S_A} \left(\sum_{S_B} m_{S_A \times S_B}(f) v(S_B) \right) v(S_A)$$

Now, if $x \in S_A$, then clearly $m_{S_A \times S_B}(f) \leq m_{S_B}(g_x)$. Consequently, for $x \in S_A$ we have

$$\sum_{S_B} m_{S_A \times S_B}(f) \cdot v(S_B) \leq \sum_{S_B} m_{S_B}(g_x) \cdot v(S_B) \leq \int_{B-} g_x = \mathfrak{L}(x).$$

Therefore

$$\sum_{S_A} \left(\sum_{S_B} m_{S_A \times S_B}(f) v(S_B) \right) v(S_A) \leq \mathcal{L}(\mathfrak{L}, \mathcal{P}_A).$$

When the function is reimenn integral □

2.3 Change of variables

Theorem 2.5. *Let $A \in \mathbb{R}^n$ be open, $g : A \rightarrow \mathbb{R}^n$ be injective and continuously differentiable with $\det g'(x) \neq 0, \forall x \in A$. Let $f : g(A) \rightarrow \mathbb{R}$ be integrable. Then we have change of variables formula:*

$$\int_{g(A)} f = \int_A (f \circ g) \cdot |\det g'(x)| dx$$

3 Calculus on Manifolds

3.1 Manifolds

Definition 3.1 (C^∞). *A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a C^∞ function if all partial derivatives of all orders of all components exists and are continuous*

Definition 3.2 (Diffeomorphism). *Let U, V be open sets in \mathbb{R}^n . A C^∞ function $h : U \rightarrow V$ bijective and $h^{-1} : V \rightarrow U$, also a C^∞ function, is called a diffeomorphism from U to V*

Definition 3.3. A set M is a K -dim manifold in \mathbb{R}^n if the following condition (M) holds. For every $x \in M$:

(M): There exists two open sets U, V of \mathbb{R}^n , $x \in U$ and a diffeomorphism $h : U \rightarrow V$ such that:

$$h(U \cap M) = \{y \in V \text{ s.t. } y^{k+1} = y^{k+2} = \dots = y^n = 0\}$$

Remark. Reminder of linear algebra

$T : \mathbb{R}^n \rightarrow \mathbb{R}^p$ Linear transformation, $\text{rank}(T) = \dim(T(\mathbb{R}^n)) \leq p$

$[T] :$ rank = max number of linearly independent rows or columns.

$\text{rank}(T) \leq \min(n, p)$

$$[T] = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{p1} & & a_{pn} \end{pmatrix}, \quad [T] \in M_{p \times n}$$

determinant of minors: If r is the max size of an $r \times r$ minor with non-zero determinant, then $\text{rank}(T) = r$.

Theorem 3.1. Let $A \subset \mathbb{R}^n$ be open and let $g : A \rightarrow \mathbb{R}^p$ be a differentiable function such that $g'(x)$ has rank p on the set $g^{-1}(0)$. Then $g^{-1}(0)$ is an $n-p$ dimensional manifold in \mathbb{R}^n .

Proof. It follows directly from theorem 1.16. Let $x \in g^{-1}(0) = M$. We take $V = A$ in theorem 1.16 so that we can find a diffeomorphism $H : V \rightarrow U$, where U is open in \mathbb{R}^n and

$$g \circ H(x^1, x^2, \dots, x^n) = (x^{n-p+1}, x^{n-p+2}, \dots, x^n)$$

Let $h = H^{-1} : U \rightarrow V$. We need to show that

$$h(U \cap M) = \{y \in V, y^{n-p+1} = y^{n-p+2} = \dots = y^n = 0\}.$$

Let $y \in U \cap M$. Then $y \in g^{-1}(0)$, i.e. $g(y) = 0$. Since

$$h(g^{-1}(0)) = H^{-1}(g^{-1}(0)) = (g \circ H)^{-1}(0) = \{y \in V, y^{n-p+1} = y^{n-p+2} = \dots = y^n = 0\},$$

clearly we have for $y \in g^{-1}(0)$ that $h(y)$ has its last p coordinates zero. The converse is also obvious: if $z \in \{y \in V, y^{n-p+1} = y^{n-p+2} = \dots = y^n = 0\}$ then set $y = H(z) \in U$ and $g(y) = g(H(z)) = (z^{n-p+1}, \dots, z^n) = (0, \dots, 0) \Rightarrow y \in g^{-1}(0) = M$ and $z = h(y) \in h(U \cap M)$. \square

Example 3.1.

$$S^2 = \{(x, y, z) \in \mathbb{R}^3; x^2 + y^2 + z^2 = 1\} \quad \text{2-dim manifold in } \mathbb{R}^3$$

$$g : \mathbb{R}^3 \rightarrow \mathbb{R}$$

$$g(x, y, z) = x^2 + y^2 + z^2 - 1 \quad S^2 = g^{-1}(0)$$

$$g'(x, y, z) = \left(\frac{dg}{dx}, \frac{dg}{dy}, \frac{dg}{dz}\right) = (2x, 2y, 2z)$$

Rank can be 0 or 1

$$\underbrace{2}_{(n-p)} = \underbrace{3}_{(n)} - \underbrace{1}_{(p)} \quad \text{Aim to show } \text{rank}(g') = 1 \text{ on } M = g^{-1}(0)$$

$$\text{rank } g' = 0 \Leftrightarrow 2x = 2y = 2z = 0$$

$$\Leftrightarrow (x, y, z) = (0, 0, 0)$$

but $(0, 0, 0) \notin g^{-1}(0)$ because $g(0, 0, 0) = 0^2 + 0^2 + 0^2 - 1 = -1$

Example 3.2. The Sphere

$S^n = \{(x^1, \dots, x^{n+1}); (x^1)^2 + \dots + (x^{n+1})^2 = 1\}$ is an n -dim manifold in \mathbb{R}^{n+1}

$g : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$

$$g(x^1, \dots, x^{n+1}) = (x^1)^2 + \dots + (x^{n+1})^2 - 1 \quad S^n = g^{-1}(0)$$

$$g'(x^1, \dots, x^{n+1}) = \left(\frac{dg}{dx^1}, \dots, \frac{dg}{dx^{n+1}} \right) = \underbrace{(2x^1, \dots, 2x^{n+1})}_{1 \times (n+1) \text{ matrix}}$$

$$\begin{aligned} \text{rank } g' = 0 &\Leftrightarrow 2x^1 = \dots = 2x^{n+1} = 0 \\ &\Leftrightarrow (x^1, \dots, x^{n+1}) = (0, \dots, 0) \end{aligned}$$

but $(0, \dots, 0) \notin S^n$

Example 3.3. Hyperbolic space

$$\mathbb{H}^n = \{(x^1, \dots, x^{n+1}) \in \mathbb{R}^{n+1}, (x^1)^2 - [(x^2)^2 + \dots + (x^{n+1})^2] = 1\}$$

n -dim hyperbolic space $g : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$

$$g(x^1, \dots, x^{n+1}) = (x^1)^2 - [(x^2)^2 + \dots + (x^{n+1})^2] - 1$$

$$\mathbb{H}^n = g^{-1}(0) \Rightarrow (g : \mathbb{R}^{n+1} \rightarrow \mathbb{R}) \rightarrow (g : A \rightarrow \mathbb{R}), \quad A = \{x \in \mathbb{R}^{n+1}, x^1 > 0\}.$$

$$g'(x^1, \dots, x^{n+1}) = (2x^1, -2x^2, \dots, -2x^{n+1})$$

$$\begin{aligned} \text{rank } g' = 0 &\Leftrightarrow x^1 = \dots = x^{n+1} = 0 \\ &\Leftrightarrow (x^1, \dots, x^{n+1}) = (0, \dots, 0) \end{aligned}$$

but $(0, \dots, 0) \notin \mathbb{H}^n$ so $\text{rank}(g') = 1$ on $g^{-1}(0)$ so by theorem 3.1 $g^{-1}(0)$ is an $(n+1) - 1$ dim manifold in \mathbb{R}^{n+1} .

Example 3.4. Ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \quad a, b, c > 0$$

$$g(x, y, z) = \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1$$

$$g'(x, y, z) = \left(\frac{2x}{a^2}, \frac{2y}{b^2}, \frac{2z}{c^2} \right) = 0 \Leftrightarrow x = y = z = 0$$

but $(0, 0, 0)$ does not belong to the ellipsoid.

Example 3.5. The graph of a differentiable function

$f : U \rightarrow \mathbb{R}, \quad U \subseteq \mathbb{R}^2$

$$M = \{(x, y, z) \in \mathbb{R}^3, z = f(x, y)\} \quad (\text{mangepatch})$$

2-dim manifold in \mathbb{R}^3

$$g(x, y, z) = f(x, y) - z$$

$$g'(x, y, z) = \left(\frac{df}{dx}, \frac{df}{dy}, -1 \right) \neq 0 \Rightarrow \text{rank}(g') = 1$$

The following theorem gives the coordinate definition of a manifold M .

Theorem 3.2. *A subset M of \mathbb{R}^n is a k -dimensional manifold iff for every point $x \in M$ the following holds:*

(C) *There exists an open set $U \subset \mathbb{R}^n$, $x \in U$ and an open set $W \subset \mathbb{R}^k$ and an injective differentiable map $f : W \rightarrow \mathbb{R}^n$ such that*

$$(i) \quad f(W) = U \cap M$$

$$(ii) \quad \text{rank } f'(y) = k \quad \forall y \in W$$

$$(iii) \quad f^{-1} : f(W) \rightarrow W \text{ is continuous.}$$

Proof. Lets assume that M is a manifold according to the definition (M). We choose the function $h : U \rightarrow V$ as in the definition. We define the set W and the function f as follows:

$$W = \{a \in \mathbb{R}^k, (a, 0) \in h(U \cap M)\}, \quad f : W \rightarrow \mathbb{R}^n, \quad f(a) = h^{-1}(a, 0).$$

Here $(a, 0)$ is the vector with the last $n - k$ coordinates equal to 0. Obviously $f(W) = U \cap M$, since

$$a \in W \Leftrightarrow (a, 0) \in h(U \cap M) \Leftrightarrow h^{-1}(a, 0) \in U \cap M \Leftrightarrow f(a) \in U \cap M.$$

We prove that W is open. For $a \in W$, we have:

$$(a, 0) \in h(U \cap M) \Leftrightarrow h^{-1}(a, 0) \in U \cap M \Rightarrow h^{-1}(a, 0) \in U.$$

Since h^{-1} is continuous, if b is sufficiently close to a , so that $(a, 0)$ and $(b, 0)$ are sufficiently close, we can deduce that $h^{-1}(b, 0)$ is close enough to $h^{-1}(a, 0)$. Because U open, if $h^{-1}(a, 0) \in U$, then also $h^{-1}(b, 0) \in U$. This gives $(b, 0) \in h(U)$. Because $h(U \cap M)$ consists exactly of the points with last $n - k$ components equal to 0, $(b, 0) \in h(U \cap M) \Leftrightarrow b \in W$. We immediately see from the definition of f and W that f^{-1} is continuous (it maps $h^{-1}(a, 0)$ to a while h is continuous).

We prove that the rank of $f'(y)$ is k on W . For this we introduce another function

$$H : U \rightarrow \mathbb{R}^k, \quad H(z) = (h^1(z), \dots, h^k(z)),$$

i.e. H has the same first k coordinates as h (and ignores the last $n - k$). We have

$$H(f(y)) = H(h^{-1}(y, 0)) = y, \quad y \in W.$$

Therefore, $H'(f(y)) \cdot f'(y) = I_{k \times k}$ or, in terms of linear transformations:

$$DH(f(y)) \circ Df(y) = Id_{\mathbb{R}^k}.$$

Because the composition is injective, $Df(y)$ is injective and the nullity plus rank theorem for $Df(y) : \mathbb{R}^k \rightarrow \mathbb{R}^n$ gives that the rank of $Df(y)$ is k .

The converse: Suppose that $f : W \rightarrow \mathbb{R}^n$ satisfies condition (C). We have $f'(y) \in M_{n \times k}$. By rearranging the coordinates in \mathbb{R}^n , we can assume that the rank of the first k rows of $f'(a)$ is k . This means

$$\det(D_j f^i(a))_{i,j=1,\dots,k} \neq 0.$$

We define

$$g : W \times \mathbb{R}^{n-k} \rightarrow \mathbb{R}^n, \quad g(a, b) = f(a) + (0, b),$$

where $(0, b)$ has the first k coordinates 0. We have

$$g^i(a, b) = f^i(a), \quad i \leq k, \quad g^i(a, b) = f^i(a) + b^i, \quad i > k.$$

We compute its Jacobian matrix. For $i \leq k$

$$D_j g^i(a, b) = D_j f^i(a) \Rightarrow D_j g^i(a, b) = D_j f^i(a), \quad j \leq k, \\ \text{and} \quad D_j g^i(a, b) = 0, \quad j > k.$$

For $i > k$, however, we have

$$D_j g^i(a, b) = D_j f^i(a) + D_j b^i \Rightarrow D_j g^i(a, b) = \delta_{ij}, \quad j > k$$

while

$$D_j g^i(a, b) = D_j f^i(a), \quad j \leq k.$$

The Jacobian matrix is therefore in block form

$$g'(a, b) = \left(\begin{array}{c|c} D_j f^i(a)_{i,j=1,\dots,k} & 0 \\ \hline D_j f^i(a)_{i=k+1,\dots,n} & I_{(n-k) \times (n-k)} \end{array} \right)$$

The calculation of the determinant in block form (which can be considered as successive expansion on the last column) gives that $\det g'(a, b) \neq 0$. By the inverse function theorem, there exists an open set V_1 with $(a, 0) \in V_1$ and an open set V_2 containing $g(a, 0) = f(a)$, such that $g : V_1 \rightarrow V_2$ has a differentiable inverse $h : V_2 \rightarrow V_1$. Then, since $f(W) = U \cap M$, we have for $(x, 0) \in V_1$, $g(x, 0) \in M \Leftrightarrow f(x) \in M$: This gives

$$V_2 \cap M = \{g(x, 0), (x, 0) \in V_1\}.$$

$$h(V_2 \cap M) = g^{-1}(V_2 \cap M) = g^{-1}(\{g(x, 0), (x, 0) \in V_1\}) = V_1 \cap (\mathbb{R}^k \times \{0\}).$$

□

Example 3.6. *2-dim torus.*

$$(x - 2)^2 = z^2 = 1$$

$$(r - 2)^2 + z^2 = 1$$

$$\begin{aligned} z &= \sin \phi & x &= r \cos \theta \\ r - 2 &= \cos \phi & y &= r \sin \theta \end{aligned}$$

$$f(\theta, \phi) = ((2 + \cos \phi) \cos \theta, (2 + \cos \phi) \sin \theta, \sin \phi) \quad \theta, \phi \in (-\pi, \pi)$$

For theorem 3.2, take $U = \mathbb{R}^3$

$$f(W) = U \cap M$$

$$f(W) = M$$

$$f : \begin{matrix} W \\ \subseteq \mathbb{R}^2 \end{matrix} \rightarrow \mathbb{R}^3$$

$$f'(\theta, \phi) = \begin{pmatrix} (2 + \cos \phi)(-\sin \theta) & -\sin \phi \cos \theta \\ (2 + \cos \phi) \cos \phi & -\sin \phi \sin \theta \\ 0 & \cos \phi \end{pmatrix}$$

$$\begin{aligned}
2 \times 2 \text{ minor: } & \begin{vmatrix} (2 + \cos \phi)(-\sin \theta) & -\sin \phi \cos \theta \\ (2 + \cos \phi) \cos \phi & -\sin \phi \sin \theta \end{vmatrix} \\
& = (2 + \cos \phi)(-\sin \phi) \begin{vmatrix} -\sin \theta & \cos \theta \\ \cos \phi & \sin \theta \end{vmatrix} \\
& = (2 + \cos \phi) \sin \phi \neq 0 \text{ iff } \sin \phi \neq 0 \Leftrightarrow \phi \neq 0
\end{aligned}$$

$\therefore \text{rank } f' = 2$ whenever $\phi \neq 0$

When $\phi = 0$

$$f'(\theta, \phi) = \begin{pmatrix} -3 \sin \theta & 0 \\ 3 \cos \theta & 0 \\ 0 & 1 \end{pmatrix}$$

if $\theta = 0$ use:

$$\begin{vmatrix} 3 \cos \theta & 0 \\ 0 & 1 \end{vmatrix} = \begin{vmatrix} 3 & 0 \\ 0 & 1 \end{vmatrix} = 3$$

if $\theta \neq 0$ use:

$$\begin{vmatrix} 3 \sin \theta & 0 \\ 0 & 1 \end{vmatrix} = 3 \sin \theta \neq 0 \text{ on } \theta \in (-\infty, \infty), \theta \neq 0$$

Example 3.7. Any nice surface of revolution is a 2-dim manifold in \mathbb{R}^3

$$\gamma(t) = (r(t), z(t)) \quad t \in (a, b)$$

γ does not have any self intersections $r(t) > 0$. If γ is differentiable and

$$\gamma'(t) = (r'(t), z'(t)) \neq 0 \quad \forall t \in (a, b)$$

then when we rotate it around the z -axis we get the surface

$$f(t, \theta) = (r(t) \cos \theta, r(t) \sin \theta, z(t)) \quad t \in (a, b), \theta \in (-\pi, \pi)$$

$$f'(t, \theta) = \begin{pmatrix} r' \cos \theta & -r \sin \theta \\ r' \sin \theta & r \cos \theta \\ z' & 0 \end{pmatrix}$$

$$\begin{vmatrix} r' \cos \theta & -r \sin \theta \\ r' \sin \theta & r \cos \theta \end{vmatrix} = r \cdot r'$$

$$r > 0, r' \neq 0 \Rightarrow \text{rank} = 2$$

We also need the definition of manifold with boundary. While a k -dimensional manifold in \mathbb{R}^n looks like a k -dim slice of \mathbb{R}^n , according to condition (M), for a manifold with boundary in \mathbb{R}^n , the part close to the boundary looks like a half-slice of dimension k . To make this precise we define the half-space

$$\mathbb{H}^k = \{x \in \mathbb{R}^k, x^k \geq 0\}.$$

Then

$$h(U \cap M) = \{y \in V : y^k \geq 0, y^{k+1} = y^{k+2} = \dots = y^n = 0\}$$

is the substitute for condition (M). More precisely:

Definition 3.4. A subset M of \mathbb{R}^n is a k -dimensional manifold with boundary if for every point x of M either condition (M) holds or (exclusive) the following condition holds:
(M') There is an open set U of \mathbb{R}^n containing x , an open set V contained in \mathbb{R}^n and a diffeomorphism $h : U \rightarrow V$ such that

$$h(U \cap M) = V \cap (\mathbb{H}^k \times \{0\}) = \{y \in V : y^k \geq 0, y^{k+1} = y^{k+2} = \dots = y^n = 0\}.$$

Moreover, $h^k(x) = 0$. The set of points where condition (M') holds is called the boundary of M and is denoted by ∂M .

3.2 Dual Space

Definition 3.5. Let $g^i : \mathbb{R}^n \rightarrow \mathbb{R}$ be a linear map, such a map is called a linear functional. The set of all linear functionals from $\mathbb{R}^n \rightarrow \mathbb{R}$ is called the dual space of \mathbb{R}^n , denoted $(\mathbb{R}^n)^*$
let g^1, \dots, g^m be linear functionals $g^i : \mathbb{R}^n \rightarrow \mathbb{R}$, then I can combine them to get a map $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $g(x) = (g^1(x), \dots, g^m(x))$
 $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear such for $x, y \in \mathbb{R}^n, \lambda \in \mathbb{R}$

$$g(\lambda x + y) = \lambda g(x) + g(y)$$

this can be seen by

$$\begin{aligned} g(\lambda x + y) &= (g^1(\lambda x + y), \dots, g^m(\lambda x + y)) \\ &= (\lambda g^1(x) + g^1(y), \dots, \lambda g^m(x) + g^m(y)) \\ &= \lambda(g^1(x), \dots, g^m(x)) + (g^1(y), \dots, g^m(y)) \end{aligned}$$

$[g^i]$ is the matrix representation of g^i

$$[g^i] = (g^i_1, \dots, g^i_n)$$

$$[g]_{m \times n} = \begin{pmatrix} g^1_1 & \dots & g^1_n \\ \vdots & & \vdots \\ g^m_1 & \dots & g^m_n \end{pmatrix}$$

Theorem 3.3. $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at a iff f^i are differentiable at $a, i = 1, \dots, m$ and $Df(a) = (Df^1, \dots, Df^m(a))$

Proof. assume f is differentiable at a we take the linear function $\Pi^i(x^1, \dots, x^m) = x^i$ and compose it with f we get

$$f^i = \Pi^i \circ f$$

this is differentiable by chain rule since f and Π^i are differentiable $\forall i = 1, \dots, m$

$$\Rightarrow Df^i = D\Pi^i(a) \cdot Df(a)$$

$$D\Pi^i = \Pi^i$$

$$\Rightarrow Df^i = \Pi^i(a) \cdot Df(a)$$

Now assume the all f^i are differentiable at $a \forall i = 1, \dots, m$

$$\begin{aligned} f(a+h) - f(a) &- (Df^1(a)(h), \dots, Df^m(a)(h)) \\ &= (f^1(a+h), \dots, f^m(a+h)) - (f^1, \dots, f^m) - (Df^1(a)(h), \dots, Df^m(a)(h)) \\ &= (f^1(a+h) - f^1(a) - Df^1(a)(h), \dots, f^m(a+h) - f^m(a) - Df^m(a)(h)) \end{aligned}$$

So

$$\begin{aligned} & \frac{|f(a+h) - f(a) - (Df^1(a)(h), \dots, Df^m(a)(h))|}{|h|} \\ & \leq \frac{|f^1(a+h) - f^1(a) - df^1(a)|}{|h|}, \dots, \frac{|f^m(a+h) - f^m(a) - df^m(a)|}{|h|} \rightarrow 0 \end{aligned}$$

□

Remark. If $T, S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear then $(T + S) : \mathbb{R}^n \rightarrow \mathbb{R}^m, (T + S)(x) = T(x) + S(x)$ is linear.

If $\lambda \in \mathbb{R}$ then $(\lambda T) : \mathbb{R}^n \rightarrow \mathbb{R}^m, (\lambda T)(x) = \lambda \cdot T(x)$ is also linear.

Definition 3.6. A linear functional f is a linear transformation $f : V \rightarrow \mathbb{R}$

$$f(\lambda x + y) = \lambda f(x) + f(y) \quad \forall x, y \in V, \forall \lambda \in \mathbb{R}$$

Definition 3.7 (Dual Space).

$$V^* = \{f : V \rightarrow \mathbb{R} : f \text{ are linear functionals}\}$$

Definition 3.8. If $f, g \in V^*$ and $\lambda \in \mathbb{R}$ then

$$f + g, \lambda f : V \rightarrow \mathbb{R}$$

with:

$$(f + g)(x) = f(x) + g(x)$$

$$(\lambda f) = \lambda f(x)$$

Proposition 3.1.

$$\dim(V^*) = \dim(V)$$

Proof. We have $\{v_1, \dots, v_n\}$ a basis of $V, \forall i = 1, \dots, n$ define $\varphi_i : V \rightarrow \mathbb{R}$ as follows:

Given $x \in V$

$$x = x^1 v_1 + \dots + x^n v_n \quad (\text{uniquely}) \quad x^i \in \mathbb{R}$$

$$\varphi_i(x) = x^i$$

φ_i is a linear functional. If $y \in V$

$$y = y^1 v_1 + \dots + y^n v_n \quad y^i \in \mathbb{R}$$

for $\lambda \in \mathbb{R}$

$$\lambda x + y = (\lambda x^1 + y^1) v_1 + \dots + (\lambda x^n + y^n) v_n$$

$$\varphi_i(\lambda x + y) = \lambda x^i + y^i = \lambda \varphi_i(x) + \varphi_i(y)$$

$$\varphi_i(v_j) = \delta_{ij}$$

Is $\{\varphi_1, \dots, \varphi_n\}$ a basis for V^* ?

Spanning:

Given $f \in V^*$, define $a^i \in \mathbb{R}$

$$f(v_i) = a^i$$

we will show

$$f = a^1\varphi_1 + \cdots + a^n\varphi_n$$

if f and $a^1\varphi_1 + \cdots + a^n\varphi_n$ agree on the basis $\{v_1, \dots, v_n\}$ then its true. For $K = 1, \dots, n$

$$f(v_k) = a^k$$

$$(a^1\varphi_1, \dots, a^n\varphi_n)(v_k) = a^1\varphi_1(v_k) + \cdots + a^k\varphi_k(v_k) + \cdots + a^n\varphi_n(v_k) = a^k \cdot 1$$

Linear Independence:

$$b^1\varphi_1 + \cdots + b^n\varphi_n = 0 \stackrel{?}{\Rightarrow} b^k = 0 \quad \forall k$$

Apply to a basis vector v_k

$$(b^1\varphi_1 + \cdots + b^n\varphi_n)(v_k) = b^1 \cdot 0 + \cdots + b^{k-1} \cdot 0 + b^k \cdot 1 + b^{k+1} \cdot 0 + \cdots + b^n \cdot 0 = b^k$$

□

Hence if $\{v_1, \dots, v_n\}$ a basis of V , then $\{\varphi_1, \dots, \varphi_n\}$ a basis of V^*

3.3 Multilinear Algebra

Definition 3.9. Let V be a vector space over \mathbb{R} , define

$$V^k = \underbrace{V \times \cdots \times V}_{k \text{ times}}$$

To be

$$V^k = \{(v_1, \dots, v_k) : v_i \in V\}.$$

This is a vector space with operations

$$(v_1, \dots, v_k) + (w_1, \dots, w_k) = (v_1 + w_1, \dots, v_k + w_k)$$

$$\lambda(v_1, \dots, v_k) = (\lambda v_1, \dots, \lambda v_k)$$

Definition 3.10. $\mathcal{T} : V^k \rightarrow \mathbb{R}$ is called multilinear if $\forall i = 1, \dots, k$

$$\begin{aligned} \mathcal{T}(v_1, \dots, v_{i-1}, v_i + v'_i, v_{i+1}, \dots, v_k) &= \mathcal{T}(v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_k) \\ &\quad + \mathcal{T}(v_1, \dots, v_{i-1}, v'_i, v_{i+1}, \dots, v_k) \end{aligned}$$

$$\mathcal{T}(v_1, \dots, v_{i-1}, \lambda v_i, v_{i+1}, \dots, v_k) = \lambda \mathcal{T}(v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_k)$$

\mathcal{T} is not linear on V^k but linear on each of its components. A \mathcal{T} like this is called a k -tensor on V .

Definition 3.11.

$$\mathcal{J}^k(V) = \{\mathcal{T} : V^k \rightarrow \mathbb{R} : k - \text{multilinear}\}$$

Note. if $k = 2$ then \mathcal{T} is called bilinear, and

$$\mathcal{T}(v_1 + v_2, w) = \mathcal{T}(v_1, w) + \mathcal{T}(v_2, w)$$

$$\mathcal{T}(\lambda v, w) = \lambda \mathcal{T}(v, w)$$

$$\mathcal{T}(v, w_1 + w_2) = \mathcal{T}(v, w_1) + \mathcal{T}(v, w_2)$$

$$\mathcal{T}(v, \lambda w) = \lambda \mathcal{T}(v, w)$$

also \mathcal{T} is called Symmetric if

$$\mathcal{T}(v, w) = \mathcal{T}(w, v)$$

\mathcal{T} is called positive definite if

$$\mathcal{T}(v, w) \geq 0 \quad \forall v, w$$

Definition 3.12 (Symmetric k-tensor). \mathcal{T} is a Symmetric k-tensor if $\forall v_i, \dots, v_k \in V$

$$\mathcal{T}(v_1, \dots, v_i, v_{i+1}, \dots, v_j, \dots, v_k) = \mathcal{T}(v_1, \dots, v_j, v_{i+1}, \dots, v_i, \dots, v_k)$$

Definition 3.13 (Alternating k-tensor). \mathcal{T} is a Alternating k-tensor if $\forall v_i, \dots, v_k \in \mathcal{V}$

$$\mathcal{T}(v_1, \dots, v_i, v_{i+1}, \dots, v_j, \dots, v_k) = -\mathcal{T}(v_1, \dots, v_j, v_{i+1}, \dots, v_i, \dots, v_k)$$

Example 3.8. $V = \mathbb{R}^2$, $V^2 = \mathbb{R}^2 \times \mathbb{R}^2$

$$\mathcal{T}(v_1, v_2) = v_1^1 v_2^2 - v_2^1 v_1^2 \quad \boxed{\text{determinant}}$$

$$v_1 = (v_1^1, v_1^2) \quad v_2 = (v_2^1, v_2^2)$$

$$\begin{vmatrix} \lambda v_1 + v_1' \\ v_2 \end{vmatrix} = \lambda \begin{vmatrix} v_1 \\ v_2 \end{vmatrix} + \begin{vmatrix} v_1' \\ v_2 \end{vmatrix}$$

$$\begin{vmatrix} v_1 \\ \lambda v_2 + v_2' \end{vmatrix} = \lambda \begin{vmatrix} v_1 \\ v_2 \end{vmatrix} + \begin{vmatrix} v_1 \\ v_2' \end{vmatrix}$$

$$\begin{vmatrix} v_1 \\ v_2 \end{vmatrix} = - \begin{vmatrix} v_2 \\ v_1 \end{vmatrix}$$

\det on $k \times k$ -matrices as a function of k vectors in \mathbb{R}^k is an alternating k-tensor.

Definition 3.14. If $\mathcal{T}, \mathcal{S} \in \mathcal{J}^k(V)$, we define:

$$(\mathcal{T} + \mathcal{S})(v_1, \dots, v_k) = \mathcal{T}(v_1, \dots, v_k) + \mathcal{S}(v_1, \dots, v_k)$$

Similarly if $\lambda \in \mathbb{R}$, $\lambda \mathcal{T} \in \mathcal{J}^k(V)$

$$(\lambda \mathcal{T})(v_1, \dots, v_k) = \lambda \mathcal{T}(v_1, \dots, v_k) \quad \forall v_i \in v$$

Definition 3.15. Let $\mathcal{T} \in \mathcal{J}^k(V)$, $\mathcal{S} \in \mathcal{J}^l(V)$, $k, l \in \mathbb{N}$, $\mathcal{T} : V^k \rightarrow \mathbb{R}$, $\mathcal{S} : V^l \rightarrow \mathbb{R}$. Define $\mathcal{T} \otimes \mathcal{S} \in \mathcal{J}^{k+l}$:

$$\mathcal{T} \otimes \mathcal{S}(v_1, \dots, v_k, v_{k+1}, \dots, v_{k+l}) = \mathcal{T}(v_1, \dots, v_k) \cdot \mathcal{S}(v_{k+1}, \dots, v_{k+l})$$

$$\mathcal{T}, \mathcal{S} \in \mathcal{J}^{k+l} \Rightarrow \mathcal{T} \otimes \mathcal{S} \neq \mathcal{S} \otimes \mathcal{T} \quad \text{in general}$$

Properties.

1. $\mathcal{T} \otimes \mathcal{S} \in \mathcal{J}^{k+l}$
2. $(\mathcal{S}_1 + \mathcal{S}_2) \otimes \mathcal{T} = \mathcal{S}_1 \otimes \mathcal{T} + \mathcal{S}_2 \otimes \mathcal{T}$
3. $\mathcal{S} \otimes (\mathcal{T}_1 + \mathcal{T}_2) = \mathcal{S} \otimes \mathcal{T}_1 + \mathcal{S} \otimes \mathcal{T}_2$

$$4. (\lambda \mathcal{S}) \otimes \mathcal{T} = \lambda(\mathcal{S} \otimes \mathcal{T}) = \mathcal{S} \otimes (\lambda \mathcal{T})$$

$$5. (\mathcal{S} \otimes \mathcal{T}) \otimes \mathcal{U} = \mathcal{S} \otimes (\mathcal{T} \otimes \mathcal{U})$$

$$6. \mathcal{J}^1(V) = V^*$$

Theorem 3.4. Let $i_1, \dots, i_k \in \{1, \dots, n\}$, V has a basis $\{v_1, \dots, v_n\}$ $\dim V = n$. Let $\{\varphi_1, \dots, \varphi_n\}$ be the basis basis of V^* , $\varphi_i(v_i) = \delta_{ij}$. Then

$$\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k} \quad \text{where} \quad \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$$

form basis for $\mathcal{J}^k(V)$

$$\dim(\mathcal{J}^k(V)) = n^k$$

Proof. Clearly $\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k} \in \mathcal{J}^k(V)$ since $\varphi_{i_j} \in V^* = \mathcal{J}_1(V)$. The set spans $\mathcal{J}^k(V)$ and is *L.I.* Let $\mathcal{T} \in \mathcal{J}^k(V)$. need to write

$$\mathcal{T} = \sum_{\substack{i_1=1, \dots, n \\ \vdots \\ i_k=1, \dots, n}} a^{i_1 i_2 \dots i_k} \varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}$$

Plug $(v_{j_1}, \dots, v_{j_k})$ into the suspected identity.

$$\begin{aligned} \mathcal{T}(v_{j_1}, \dots, v_{j_k}) &= \sum_{i_1, \dots, i_k} a^{i_1 i_2 \dots i_k} \varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}(v_{j_1}, \dots, v_{j_k}) \\ &= \sum_{i_1, \dots, i_k} a^{i_1 i_2 \dots i_k} \varphi_{i_1}(v_{j_1}) \dots \varphi_{i_k}(v_{j_k}) \\ &= \sum_{i_1, \dots, i_k} a^{i_1 i_2 \dots i_k} \delta_{i_1 j_1} \dots \delta_{i_k j_k} \\ &= a^{j_1 \dots j_k} \end{aligned}$$

Define

$$a^{i_1 i_2 \dots i_k} = \mathcal{T}(v_{j_1}, \dots, v_{j_k})$$

Let $w_1, \dots, w_k \in V$

$$w_1 = \sum_{j=1}^n a^{1j} v_j \quad \dots \quad w_k = \sum_{j=1}^n a^{kj} v_j$$

$$\begin{aligned} \mathcal{T}(w_1, \dots, w_k) &= \mathcal{T} \left(\sum_{j_1} a^{1j_1} v_{j_1}, \dots, \sum_{j_k} a^{kj_k} v_{j_k} \right) \\ &= \sum_{j_1, \dots, j_k=1}^n a^{1j_1} \dots a^{kj_k} \cdot \mathcal{T}(v_{j_1}, \dots, v_{j_k}) \\ &= \sum_{j_1, \dots, j_k} a^{1j_1} \dots a^{kj_k} \cdot a^{i_1 i_2 \dots i_k} \end{aligned}$$

$$\begin{aligned}
\sum_{i_1, \dots, i_k} a^{i_1 \dots i_k} \varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}(w_1, \dots, w_k) &= \sum_{i_1, \dots, i_k} a^{i_1 \dots i_k} \varphi_{i_1}(w_1) \otimes \dots \otimes \varphi_{i_k}(w_k) \\
&= \sum_{j_1, \dots, j_k} a^{i_1 i_2 \dots i_k} \cdot a^{1_{i_1}} \dots a^{k_{i_k}}
\end{aligned}$$

Relabel:

$$\begin{aligned}
i_1 &\mapsto j_1 \\
&\vdots \\
i_k &\mapsto j_k
\end{aligned}$$

So it Spans. Now check L.I.

$$\sum_{i_1, \dots, i_k=1}^n a^{i_1 \dots i_k} \varphi_{i_1} \otimes \dots \otimes \varphi_{i_k} = 0$$

Plus v_{j_1}, \dots, v_{j_k} in to it

$$\begin{aligned}
\sum_{i_1, \dots, i_k=1}^n a^{i_1 \dots i_k} \varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}(v_{j_1}, \dots, v_{j_k}) &= \sum_{i_1, \dots, i_k} a^{i_1 \dots i_k} \delta_{i_1 j_1} \dots \delta_{i_k j_k} \\
&= a^{j_1 \dots j_k} = 0
\end{aligned}$$

put in all possible combinations of basis vectors

$$\Rightarrow \text{all coefficients } \underline{Zero}$$

□

3.4 Alternating Tensors

Remark. for $f : \mathbb{R} \rightarrow \mathbb{R}$. If f is even:

$$f(-x) = f(x)$$

if f is odd:

$$f(-x) = -f(x)$$

Every function can be written as

$$f = \underset{\text{even}}{f_1} + \underset{\text{odd}}{f_2}$$

Where

$$f_1 = \frac{f(x) + f(-x)}{2} \quad f_2 = \frac{f(x) - f(-x)}{2}$$

or σ is a bijection on $\mathbb{R} \rightarrow \mathbb{R}$, $\sigma^2 = id$.

$$\begin{aligned}
x &\xrightarrow{\sigma} -x \\
&\frac{f(x) + f(\sigma x)}{2}
\end{aligned}$$

Let S_k be that symmetric group on k letters.

$$S_k \xrightarrow{hom} \{\pm 1\} \quad \text{multiplicative group}$$

$$\sigma \mapsto \begin{cases} +1 & \text{if } \sigma \text{ even} \\ -1 & \text{if } \sigma \text{ odd} \end{cases}$$

$$\sigma \mapsto \text{sign}(\sigma).$$

Definition 3.16. If $\mathcal{T} \in \mathcal{J}^k(V)$

$$\text{Alt}(\mathcal{T})(w_1, \dots, w_k) = \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn}(\sigma) \mathcal{T}(w_{\sigma(1)}, \dots, w_{\sigma(k)}).$$

Example 3.9. $k = 2$

$$\text{Alt}(\mathcal{T})(w_1, w_2) = \frac{1}{2!} (\mathcal{T}(w_1, w_2) - \mathcal{T}(w_2, w_1))$$

Definition 3.17. The set of alternating k -tensors is denoted by $\Lambda^k(V)$, it is a subspace of $\mathcal{J}^k(V)$

Theorem 3.5.

(a) if $\mathcal{T} \in \mathcal{J}^k(V)$, $\text{Alt}(\mathcal{T}) \in \mathcal{J}^k(V)$ and $\text{Alt}(\mathcal{T})$ is alternating

(b) if \mathcal{W} is alternating, $\text{Alt}(\mathcal{W}) = \mathcal{W}$

(c) $\text{Alt}(\text{Alt}(\mathcal{T})) = \text{Alt}(\mathcal{T})$

Proof. (c) follows from (b), use $\mathcal{W} = \text{Alt}(\mathcal{T})$ which is alternating by (a)

$$\text{Alt}(\mathcal{T}) = \mathcal{W} = \text{Alt}(\mathcal{W}) = \text{Alt}(\text{Alt}(\mathcal{T}))$$

Proof of (a):

Show $\text{Alt}(\mathcal{T}) \in \mathcal{J}^k(V)$. I will show it is alternating.

$$\text{Alt}(\mathcal{T})(w_1, \dots, w_i, \dots, w_j, \dots, w_k) = -\text{Alt}(\mathcal{T})(w_1, \dots, w_j, \dots, w_i, \dots, w_k)$$

$$\begin{aligned} i &\mapsto j \\ j &\mapsto i \\ k &\mapsto k \quad \text{if } k \neq i, j \end{aligned}$$

$$\left. \begin{aligned} S_k &\rightarrow S_k \\ \sigma &\mapsto \sigma(ij) = \sigma^1 \\ \text{even} &\mapsto \text{odd} \\ \text{odd} &\mapsto \text{even} \end{aligned} \right\} \quad \text{Bijection} \quad (3)$$

$$\begin{aligned} \text{Alt}(\mathcal{T})(w_1, \dots, w_j, \dots, w_i, \dots, w_k) &= \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn}(\sigma) \text{Alt}(\mathcal{T})(w_{\sigma(1)}, \dots, w_{\sigma(j)}, \dots, w_{\sigma(i)}, \dots, w_{\sigma(k)}) \\ &= \frac{1}{k!} \sum_{\sigma' \in S_k} \text{sgn}(\sigma') \text{Alt}(\mathcal{T})(w_{\sigma'(1)}, \dots, w_{\sigma'(i)}, \dots, w_{\sigma'(j)}, \dots, w_{\sigma'(k)}) \\ &= -\text{Alt}(\mathcal{T})(w_1, \dots, w_k) \end{aligned}$$

Proof of (b):

Let ω be alternating.

$$\begin{aligned}
\omega(w_1, \dots, w_j, \dots, w_i, \dots, w_k) &= -\omega(w_1, \dots, w_i, \dots, w_j, \dots, w_k) \\
&= \omega(w_{\sigma(1)}, \dots, w_{\sigma(k)}) \\
&= \text{sgn}(\sigma) \omega(w_1, \dots, w_k) \quad \sigma \in S_k \\
&= \text{Alt}(\omega)(w_1, \dots, w_k) \\
&= \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn}(\sigma) \omega(w_{\sigma(1)}, \dots, w_{\sigma(k)}) \\
&= \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn}(\sigma)^2 \omega(w_1, \dots, w_k) \\
&= \frac{1}{k!} |S_k| \omega(w_1, \dots, w_k) \\
&= \omega
\end{aligned}$$

$\therefore \text{Alt}(\omega) = \omega$

□

Remark. If $\omega \in \Lambda^k(V)$, $\eta \in \Lambda^l(V)$ then

$$\omega \otimes \eta \in \mathcal{J}^{k+l}(V)$$

Definition 3.18.

$$\omega \wedge \eta = \frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \eta) \in \Lambda^{k+l}(V)$$

Properties. if $\omega, \omega_1, \omega_2 \in \Lambda^k(V)$, $\eta, \eta_1, \eta_2 \in \Lambda^l(V)$

- $(\omega_1 + \omega_2) \wedge \eta = \omega_1 \wedge \eta + \omega_2 \wedge \eta$
- $\omega \wedge (\eta_1 + \eta_2) = \omega \wedge \eta_1 + \omega \wedge \eta_2$
- $(\alpha\omega) \wedge \eta = \alpha(\omega \wedge \eta) = \omega \wedge (\alpha\eta) \quad \alpha \in \mathbb{R}$
- $\omega \wedge \eta = (-1)^{k \cdot l} \eta \wedge \omega$

Definition 3.19. Let V, W be vector spaces $f : V \rightarrow W$ be a linear transformation.

If \mathcal{T} is a linear functional on W , $\mathcal{T} : W \rightarrow \mathbb{R}$ then

$$\mathcal{T} \circ f$$

is a linear functional on V .

$$\begin{array}{ccc}
V & \xrightarrow{f} & W \\
& \searrow & \downarrow \mathcal{T} \\
& \mathcal{T} \circ f & \mathbb{R}
\end{array}$$

Notation.

$$f^*(\mathcal{T}) = \mathcal{T} \circ f$$

$f^*(\mathcal{T})$ is called the pullback of \mathcal{T} by f .

$$f^* : W^* \rightarrow V^*$$

by $f^*(\mathcal{T}) = \mathcal{T} \circ f$

Definition 3.20 (Pullback of Tensors). If \mathcal{T} is a k -tensor on W ie $\mathcal{T} \in \mathcal{J}^k(W)$ we define the pullback $f^* \in \mathcal{J}^k(V)$ by:

$$f^*(\mathcal{T})(v_1, \dots, v_k) = \mathcal{T}(f(v_1), \dots, f(v_k))$$

This is a k -tensor on V .

Need to show linearity in the i^{th} entry. Let $v_i, v'_i \in V, \lambda \in \mathbb{R}$.

$$\begin{aligned} f^*(\mathcal{T})(v_1, \dots, \lambda v_i + v'_i, \dots, v_k) &= \mathcal{T}(f(v_1), \dots, f(\lambda v_i + v'_i), \dots, f(v_k)) \\ &= \mathcal{T}(f(v_1), \dots, \lambda f(v_i) + f(v'_i), \dots, f(v_k)) \\ &= \lambda \mathcal{T}(f(v_1), \dots, f(v_i), \dots, f(v_k)) + \mathcal{T}(f(v_1), \dots, f(v'_i), \dots, f(v_k)) \\ &= \lambda f^*(\mathcal{T})(v_1, \dots, v_i, \dots, v_k) + f^*(\mathcal{T})(v_1, \dots, v'_i, \dots, v_k) \end{aligned}$$

□

Properties.

(a)

$$f^*(\mathcal{T} \otimes \mathcal{S}) = f^*(\mathcal{T}) \otimes f^*(\mathcal{S}) \quad \mathcal{T} \in \mathcal{J}^k(W), \mathcal{S} \in \mathcal{J}^l(W)$$

(b)

$$f^*(\mathcal{T} \wedge \mathcal{S}) = f^*(\mathcal{T}) \wedge f^*(\mathcal{S}) \quad \mathcal{T} \in \Lambda^k(W), \mathcal{S} \in \Lambda^l(W)$$

Theorem 3.6.

(a) if $\mathcal{S} \in \mathcal{J}^k(V), \mathcal{T} \in \mathcal{J}^l(V)$ and $\text{Alt}(\mathcal{S}) = 0$ then

$$\text{Alt}(\mathcal{S} \otimes \mathcal{T}) = \text{Alt}(\mathcal{T} \otimes \mathcal{S}) = 0$$

(b) if $\omega \in \Lambda^k(V), \eta \in \Lambda^l(V), \vartheta \in \Lambda^m(V)$ then

$$\text{Alt}(\text{Alt}(\omega \otimes \eta) \otimes \vartheta) = \text{Alt}(\omega \otimes \eta \otimes \vartheta) = \text{Alt}(\omega \otimes \text{Alt}(\eta \otimes \vartheta))$$

(c) if $\omega \in \Lambda^k(V), \eta \in \Lambda^l(V), \vartheta \in \Lambda^m(V)$ then

$$(\omega \wedge \eta) \wedge \vartheta = \omega \wedge (\eta \wedge \vartheta) = \frac{(k+l+m)!}{k!l!m!} \text{Alt}(\omega \otimes \eta \otimes \vartheta)$$

Proof.

Proof of (a):

$$\text{Alt}(\mathcal{S} \otimes \mathcal{T}) = \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) (\mathcal{S} \otimes \mathcal{T})(w_{\sigma(1)}, \dots, w_{\sigma(k)}, w_{\sigma(k+1)}, \dots, w_{\sigma(k+l)})$$

Let G be the subgroup of S_{k+l} such that

$$G = \left\{ \sigma \in S_{k+l} : \begin{array}{c} \sigma(k+1) = k+1 \\ \vdots \\ \sigma(k+l) = k+l \end{array} \right\}$$

The contribution of these to the sum is:

$$\begin{aligned} & \frac{1}{(k+l)!} \sum_{\sigma \in G} \text{sgn}(\sigma) \mathcal{S}(w_{\sigma(1)}, \dots, w_{\sigma(k)}) \cdot \mathcal{T}(w_{k+1}, \dots, w_{k+l}) \\ &= \frac{1}{(k+l)!} k! \text{Alt}(\mathcal{S})(w_1, \dots, w_k) \cdot \mathcal{T}(w_{k+1}, \dots, w_{k+l}) = 0 \end{aligned}$$

Let $G\sigma_0$ be a coset of G in S_{k+l} , $\sigma_0 \neq 0$

$$G\sigma_0 = \{\sigma' \cdot \sigma_0 : \sigma' \in G\}$$

define

$$(z_1, \dots, z_{k+l}) = (w_{\sigma_0(1)}, \dots, w_{\sigma_0(k+l)})$$

The contribution of these elements is

$$\begin{aligned} & \frac{1}{(k+l)!} \sum_{\sigma' \in G} \text{sgn}(\sigma' \cdot \sigma_0) \mathcal{S}(z_{\sigma'(1)}, \dots, z_{\sigma'(k)}) \cdot \mathcal{T}(z_{\sigma'(k+1)}, \dots, z_{\sigma'(k+l)}) \\ &= \frac{1}{(k+l)!} \sum_{\sigma' \in G} \text{sgn}(\sigma') \text{sgn}(\sigma_0) \mathcal{S}(z_{\sigma'(1)}, \dots, z_{\sigma'(k)}) \cdot \mathcal{T}(z_{k+1}, \dots, z_{k+l}) \\ &= \frac{1}{(k+l)!} \text{sgn}(\sigma_0) \mathcal{T}(z_{k+1}, \dots, z_{k+l}) k! \text{Alt}(\mathcal{S})(z_1, \dots, z_k) = 0 \end{aligned}$$

Proof of (b):

$$\text{Alt}(\omega \otimes \eta) - \omega \otimes \eta = \mathcal{S}$$

$$\begin{aligned} \text{Alt}(\mathcal{S}) &= \text{Alt}(\text{Alt}(\omega \otimes \eta) - \omega \otimes \eta) = \text{Alt}(\text{Alt}(\omega \otimes \eta)) - \text{Alt}(\omega \otimes \eta) \\ &= \text{Alt}(\omega \otimes \eta) - \text{Alt}(\omega \otimes \eta) = 0 \end{aligned}$$

Apply (a) with \mathcal{S}

$$\text{alt}(\mathcal{S} \otimes \vartheta) = 0$$

$$\text{Alt}([\text{Alt}(\omega \otimes \eta) - \omega \otimes \eta] \otimes \vartheta) = 0$$

$$\text{Alt}(\text{Alt}(\omega \otimes \eta) \otimes \vartheta) - \text{Alt}(\omega \otimes \eta \otimes \vartheta) = 0$$

Proof of (c):

$$\begin{aligned} (\omega \wedge \eta) \wedge \vartheta &= \frac{((k+l)+m)!}{(k+l)! m!} \text{Alt}((\omega \wedge \eta) \otimes \vartheta) \\ &= \frac{((k+l)+m)!}{(k+l)! m!} \text{Alt}\left(\frac{(k+l)!}{k! l!} \text{Alt}(\omega \otimes \eta) \otimes \vartheta\right) \\ &= \frac{((k+l)+m)!}{(k+l)! m!} \frac{(k+l)!}{k! l!} \text{Alt}(\text{Alt}(\omega \otimes \eta) \otimes \vartheta) \\ &= \frac{(k+l+m)!}{k! l! m!} \text{Alt}(\omega \otimes \eta \otimes \vartheta) \end{aligned}$$

□

Theorem 3.7. Let $\dim V = n$, then the following is a basis for $\Lambda^k(V)$

$$\varphi_{i_1} \wedge \cdots \wedge \varphi_{i_k}, \quad 1 \leq i_1 < \cdots < i_k \leq n.$$

therefore

$$\dim \Lambda^k(V) = \binom{n}{k}$$

Since choosing a subset of k items from a set of n items and reordering.

Proof. $\mathcal{T} \in \Lambda^k(V)$ then $\text{Alt}(\mathcal{T}) = \mathcal{T}$. Since

$$\varphi_{i_1} \otimes \cdots \otimes \varphi_{i_k}$$

is a basis for $\mathcal{J}^k(V)$ we have

$$\mathcal{T} = \sum_{i_1, \dots, i_k=1, \dots, n} a^{i_1 \cdots i_k} \varphi_{i_1} \otimes \cdots \otimes \varphi_{i_k}$$

Apply Alt on both sides

$$\mathcal{T} = \sum_{i_1, \dots, i_k} a^{i_1 \cdots i_k} \text{Alt}(\varphi_{i_1} \otimes \cdots \otimes \varphi_{i_k})$$

$\text{Alt}(\varphi_{i_1} \otimes \cdots \otimes \varphi_{i_k})$ is a multiple of $\varphi_{i_1} \wedge \cdots \wedge \varphi_{i_k}$. Since

$$\varphi_{i_j} \wedge \varphi_{i_s} = -\varphi_{i_s} \wedge \varphi_{i_j}$$

you can reorder to

$$\sum_{i_1 < \cdots < i_k}$$

So $\varphi_{i_1} \wedge \cdots \wedge \varphi_{i_k}$ with $i_1 < \cdots < i_k$ span $\Lambda^k(V)$. It is easy to see that they are L.I. □

Example 3.10. $\dim V = 3, k = 1$

$$\dim \Lambda^1(V) = \binom{3}{1} = 3$$

$$\Lambda^1(V) = \mathcal{J}^1(V) = V^*$$

if $\{v_i\}_{i=1,2,3}$ is a basis of V , then the dual basis $\varphi_1, \varphi_2, \varphi_3$ is a basis of $\Lambda^1(V)$

Example 3.11. $\dim V = 3, k = 2$

$$\dim \Lambda^2(V) = \binom{3}{2} = 3$$

$$\Lambda^2(V) = \mathcal{J}^2(V) = V^*$$

basis is

$$\varphi_1 \wedge \varphi_2, \varphi_1 \wedge \varphi_3 \text{ and } \varphi_2 \wedge \varphi_3$$

$$\begin{aligned}
(\varphi_1 \wedge \varphi_2)(w_1, w_2) &= \frac{(1+1)!}{1!1!} \text{Alt}(\varphi_1 \otimes \varphi_2)(w_1, w_2) \\
&= 2! \frac{1}{2!} (\varphi_1 \otimes \varphi_2(w_1, w_2) - \varphi_1 \otimes \varphi_2(w_2, w_1)) \\
&= \varphi_1(w_1) \cdot \varphi_2(w_2) - \varphi_1(w_2) \cdot \varphi_2(w_1) \\
&= \varphi_1 \otimes \varphi_2(w_1, w_2) - \varphi_2 \otimes \varphi_1(w_1, w_2)
\end{aligned}$$

$$\begin{aligned}
\therefore \varphi_1 \wedge \varphi_2 &= \varphi_1 \otimes \varphi_2 - \varphi_2 \otimes \varphi_1 \\
\varphi_1 \wedge \varphi_3 &= \varphi_1 \otimes \varphi_3 - \varphi_3 \otimes \varphi_1 \\
\varphi_2 \wedge \varphi_3 &= \varphi_2 \otimes \varphi_3 - \varphi_3 \otimes \varphi_2
\end{aligned}$$

$$\begin{aligned}
\therefore \varphi_1 \wedge \varphi_2 &= \varphi_1 \otimes \varphi_2 - \varphi_2 \otimes \varphi_1 = -\varphi_2 \wedge \varphi_1 \\
\varphi_1 \wedge \varphi_3 &= -\varphi_3 \wedge \varphi_1 \\
\varphi_2 \wedge \varphi_3 &= -\varphi_3 \wedge \varphi_2
\end{aligned}$$

$$(\varphi_1 \wedge \varphi_1)(w_1, w_2) = \varphi_1(w_1)\varphi_1(w_2) - \varphi_1(w_1)\varphi_1(w_2) = 0$$

$$\therefore \boxed{\varphi_1 \wedge \varphi_1 = \varphi_2 \wedge \varphi_2 = \varphi_3 \wedge \varphi_3 = 0}$$

Example 3.12. $\dim V = 3, k = 3$


$$\dim \Lambda^3(V) = \binom{3}{3} = 1 \quad \text{basis : } \varphi_1 \wedge \varphi_2 \wedge \varphi_3$$

$$\begin{aligned}
(\varphi_1 \wedge \varphi_2 \wedge \varphi_3)(w_1, w_2, w_3) &= 3! \text{Alt}(\varphi_1 \otimes \varphi_2 \otimes \varphi_3) \\
&= \sum_{\sigma \in S_3} \text{sgn}(\sigma) \varphi_1 \otimes \varphi_2 \otimes \varphi_3(w_{\sigma(1)}, w_{\sigma(2)}, w_{\sigma(3)}) \\
&= \varphi_1(w_1)\varphi_2(w_2)\varphi_3(w_3) - \varphi_1(w_2)\varphi_2(w_1)\varphi_3(w_3) \\
&\quad - \varphi_1(w_3)\varphi_2(w_2)\varphi_3(w_1) - \varphi_1(w_1)\varphi_2(w_3)\varphi_3(w_2) \\
&\quad + \varphi_1(w_2)\varphi_2(w_3)\varphi_3(w_1) + \varphi_1(w_3)\varphi_2(w_1)\varphi_3(w_2)
\end{aligned}$$

$$\begin{aligned}
\varphi_1 \wedge \varphi_2 \wedge \varphi_3 &= \varphi_1 \otimes \varphi_2 \otimes \varphi_3 - \varphi_2 \otimes \varphi_1 \otimes \varphi_3 - \varphi_2 \otimes \varphi_2 \otimes \varphi_1 \\
&\quad - \varphi_1 \otimes \varphi_3 \otimes \varphi_2 + \varphi_3 \otimes \varphi_1 \otimes \varphi_2 + \varphi_2 \otimes \varphi_3 \otimes \varphi_1
\end{aligned}$$

3.5 Differential Forms

Definition 3.21 (Tangent Space).



$$\mathbb{R}_p^n = \{(p, v) : v \in \mathbb{R}^n\} \quad \text{This is the tangent space at } p$$

$$(p, v) + (p, w) = (p, v + w)$$

$$\lambda(p, v) = (p, \lambda v)$$

with these operations \mathbb{R}_p^n is a vector space. The operation $(p, v) + (q, w)$ makes no sense if $p \neq q$.

Notation. $v_p = (p, v)$

on \mathbb{R}_p^n we have:

$$\langle (p, v), (p, w) \rangle = \langle v, w \rangle$$

Definition 3.22 (Vector Field). A vector field in \mathbb{R}^n is a function

$$F : p \rightarrow F(p) \in \mathbb{R}_p^n$$

$$F(p) = (p, v)$$

$$v = (F^1(p), \dots, F^n(p))$$

Properties.

- if the components F^i , $i \in \{1, \dots, n\}$ are continuous, the vector field is continuous

$$F^i : p \rightarrow F^i(p)$$

- if the components are differentiable then the vector function is differentiable.
- if F, G are vector fields in \mathbb{R}^n , then $F + G$ is also a vector field in \mathbb{R}^n

$$(F + G)(p) = F(p) + G(p)$$

- λF is a vector field $\forall \lambda \in \mathbb{R}$ and

$$(\lambda F)(p) = \lambda \cdot F(p)$$

- if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a function (continuous and differentiable) then $f \cdot F$ is a new vector field on \mathbb{R}^n .

$$(f \cdot F)(p) = f(p) \cdot F(p)$$

Definition 3.23 (Divergence). If F is a vector field then its divergence is defined to be:

$$(\operatorname{div} F)(p) = \sum_{i=1}^n D_i F^i(p) \in \mathbb{R}$$

So $\operatorname{div} F : \mathbb{R}^n \rightarrow \mathbb{R}$.

Notation.

$$\operatorname{div} F = \nabla \cdot F$$

Definition 3.24 (k-form). Given $p \in \mathbb{R}^n$, let $\omega(p) \in \Lambda^k(\mathbb{R}_p^k)$

$$\omega(p) = \sum_{i_1 < \dots < i_k} \omega_{i_1 \dots i_k}(p) \varphi_{i_1}(p) \wedge \dots \wedge \varphi_{i_k}(p)$$

it is defined by $\binom{n}{k}$ functions $p \rightarrow \omega_{i_1 \dots i_k}(p)$, $i_1 < \dots < i_k$

Properties.

- if these functions are continuous, then the k -form is continuous.
- if these functions are differentiable, then ω is a differential k -form.
- if ω and η are differentiable k -forms on \mathbb{R}^n , $\omega + \eta$ is a differentiable k -form on \mathbb{R}^n

$$(\omega + \eta)(p) = \omega(p) + \eta(p)$$

- if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a differentiable function then $f \cdot \omega$ is a differentiable k -form.

$$(f \cdot \omega)(p) = f(p)\omega(p)$$

- if ω is a differentiable k -form and η is a differentiable l -form, then $\omega \wedge \eta$ is a differentiable $(k + l)$ -form.

$$(\omega \wedge \eta)(p) = \omega(p) \wedge \eta(p)$$

Definition 3.25. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable, then

$$Df(p) : \mathbb{R}^n \rightarrow \mathbb{R}$$

is a linear map

$$Df(p) \in (\mathbb{R}_p^n)^* = \mathcal{J}^1(\mathbb{R}_p^n) = \Lambda^1(\mathbb{R}_p^n)$$

Note. we define the following 1-form

$$df(p) \in \Lambda^1(\mathbb{R}_p^n)$$

$$df(p)(v_p) = Df(p)(v), \quad v_p = (v, p)$$

Let $f = \pi^i$, the projection into the i -component.

$$\pi^i(x^1, \dots, x^n) = x^i$$

is a linear map, sometimes denoted $x^i(x) = x^i$

$$d\pi^i(p)(v_p) = D\pi^i(p)(v) = \pi^i(p)(v) = \pi^i(v) = v^i$$

But this is the same as $\varphi_i(v)$

$$\therefore d\pi^i = \varphi_i = dx^i$$

A differentiable k -form on \mathbb{R}^n will look like

$$\omega(p) = \sum_{i_1 < \dots < i_k} \omega_{i_1 \dots i_k}(p) dx^{i_1}(p) \wedge \dots \wedge dx^{i_k}(p)$$

$$\omega = \sum_{i_1 < \dots < i_k} \omega_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

Example 3.13. $\mathbb{R}^2 : (x^1, x^2) = (x, y)$ ie $n = 2$

$$\begin{aligned} k = 0 & \quad \omega = f(x, y) \\ k = 1 & \quad \omega = f(x, y)dx + g(x, y)dy \\ k = 2 & \quad \omega = f(x, y, z)dx \wedge dy \end{aligned}$$

Example 3.14. $\mathbb{R}^3 : (x^1, x^2, x^3) = (x, y, z)$

$$\begin{aligned} k = 0 & \quad \omega = f(x, y, z) \\ k = 1 & \quad \omega = f(x, y, z)dx + g(x, y, z)dy + h(x, y, z)dz \\ k = 2 & \quad \omega = f(x, y, z)dx \wedge dy + g(x, y, z)dx \wedge dz + h(x, y, z)dy \wedge dz \\ k = 3 & \quad \omega = f(x, y, z)dx \wedge dy \wedge dz \end{aligned}$$

$$dx \wedge dx = dy \wedge dy = dz \wedge dz = 0$$

$$dx \wedge dy = -dy \wedge dx$$

$$dx \wedge dz = -dz \wedge dx$$

$$dy \wedge dz = -dz \wedge dy$$

Theorem 3.8. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable, then the 1-form df is:

$$df = D_1 f dx^1 + D_2 f dx^2 + \cdots + D_n f dx^n$$

Proof. $df(p) \in \Lambda^1(\mathbb{R}_p^n)$

$$df(p)(V_p) = df(p)(v) = (D_1 f(p), \dots, D_n f(p)) \cdot \begin{pmatrix} v^1 \\ \vdots \\ v^n \end{pmatrix} = \sum_{i=1}^n D_i f(p) v^i$$

Calculate:

$$\begin{aligned} (D_1 f dx^1 + \cdots + D_n f dx^n)(p)(V_p) &= [D_1 f(p) dx^1(p) + \cdots + D_n f(p) dx^n(p)](V_p) \\ &= D_1 f(p) dx^1(p)(V_p) + \cdots + D_n f(p) dx^n(p)(V_p) \\ &= D_1 f(p) v^1 + \cdots + D_n f(p) v^n \\ &= \sum_{i=1}^n D_i f(p) v^i \end{aligned}$$

□

Example 3.15. for \mathbb{R}^3

$$df = \frac{df}{dx} dx + \frac{df}{dy} dy + \frac{df}{dz} dz$$

Definition 3.26 (The Operator d on k -forms).

$$\omega = \sum_{i_1 < \cdots < i_k} \omega_{i_1 \dots i_k} dx^{i_1} \wedge \cdots \wedge dx^{i_k} \quad (k\text{-form})$$

$$d\omega = \sum_{i_1 < \cdots < i_k} \sum_{\alpha=1}^n D_i \omega_{i_1 \dots i_k} x^\alpha \wedge dx^{i_1} \wedge \cdots \wedge dx^{i_k} \quad (k+1)\text{-form}$$

Example 3.16. $n = 3, k = 1$

$$\omega = f dx + g dy + h dz$$

$$\begin{aligned} d\omega &= \frac{df}{dx} \cancel{dx \wedge dx} + \frac{df}{dy} dy \wedge dx + \frac{df}{dz} dz \wedge dx \\ &\quad + \frac{dg}{dx} dx \wedge dy + \frac{dg}{dy} \cancel{dy \wedge dy} + \frac{dg}{dz} dz \wedge dy \\ &\quad + \frac{dh}{dx} dx \wedge dz + \frac{dh}{dy} dy \wedge dz + \frac{dh}{dz} \cancel{dz \wedge dz} \\ &= \left(\frac{dg}{dx} - \frac{df}{dy} \right) dx \wedge dy + \left(\frac{dh}{dy} - \frac{dg}{dz} \right) dy \wedge dz + \left(\frac{dh}{dx} - \frac{df}{dz} \right) dx \wedge dz \end{aligned}$$

$$\begin{vmatrix} i & j & k \\ dx & dy & dz \\ f & g & h \end{vmatrix} = \left(\frac{dh}{dy} - \frac{dg}{dz} \right) \mathbf{i} - \left(\frac{dh}{dx} - \frac{df}{dz} \right) \mathbf{j} + \left(\frac{dg}{dx} - \frac{df}{dy} \right) \mathbf{z}$$

$$\mathbf{i} \leftrightarrow dy \wedge dz \quad \mathbf{j} \leftrightarrow dz \wedge dx \quad \mathbf{k} \leftrightarrow dx \wedge dy$$

Example 3.17. $n = 3, k = 2$

$$\omega = f dy \wedge dz + g dz \wedge dx + h dx \wedge dy$$

$$\begin{aligned} d\omega &= \frac{df^1}{dx} dx \wedge dy \wedge dz + \frac{df^1}{dy} dy \wedge dy \wedge dz + \frac{df^1}{dz} dz \wedge dy \wedge dz \\ &\quad + \frac{df^2}{dx} dx \wedge dz \wedge dx + \frac{df^2}{dy} dy \wedge dz \wedge dx + \frac{df^2}{dz} dz \wedge dz \wedge dx \\ &\quad + \frac{df^3}{dx} dx \wedge dx \wedge dy + \frac{df^3}{dy} dy \wedge dx \wedge dy + \frac{df^3}{dz} dz \wedge dx \wedge dy \\ &= \left(\frac{df^1}{dx} + \frac{df^2}{dy} + \frac{df^3}{dz} \right) dx \wedge dy \wedge dz \end{aligned}$$

Example 3.18. 2-form $\leftrightarrow F = (f^1, f^2, f^3)$

$$\operatorname{div} F = \frac{df^1}{dx} + \frac{df^2}{dy} + \frac{df^3}{dz} \leftrightarrow d\omega$$

Example 3.19. $n = 3, k = 3$

$$\omega = f dx \wedge dy \wedge dz$$

$$d\omega = 0 \quad 4\text{-form on } \mathbb{R}^3 \quad \binom{3}{4} = 0$$

Example 3.20. $n = 2, k = 1$

$$\omega = f dx + g dy$$

$$\begin{aligned} d\omega &= \frac{df}{dx} dx \wedge dx + \frac{df}{dy} dy \wedge dx + \frac{dg}{dx} dx \wedge dy + \frac{dg}{dy} dy \wedge dy \\ &= \left(\frac{dg}{dx} - \frac{df}{dy} \right) dx \wedge dy \end{aligned}$$

Example 3.21. $n = 2, k = 2$

$$\begin{aligned}\omega &= f dx \wedge dy \\ d\omega &= 0\end{aligned}$$

Example 3.22. $n = 1, k = 1$

$$\begin{aligned}\omega &= f dx \\ d\omega &= 0\end{aligned}$$

Theorem 3.9.

(a) $d(\omega + \eta) = d(\omega) + d(\eta)$

(b) if ω is a k -form, η is an l -form, $\omega \wedge \eta$ is a $(k + l)$ -form.

$$d(\omega \wedge \eta) = (d\omega) \wedge \eta + (-1)^k \omega \wedge (d\eta)$$

(c) $d(d\omega) = 0$

Proof.

(c):

$$\begin{aligned}\omega &= \sum_{i_1 < \dots < i_k} \omega_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ d(d\omega) &= d \left(\sum_{i_1 < \dots < i_k} \sum_{i=1}^n D_i \omega_{i_1 \dots i_k} x^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \right) \\ &= \sum_{i_1 < \dots < i_k} \sum_{i=1}^n \sum_{j=1}^n D_j (D_i \omega_{i_1 \dots i_k}) dx^j \wedge dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\end{aligned}$$

if $i = j$ then $dx^i \wedge dx^j = 0$, if $i \neq j : (i, j), (j, i)$

$$D_j D_i \omega_{i_1 \dots i_k} dx^j \wedge dx^i - D_i D_j \omega_{i_1 \dots i_k} \overset{\text{swapped in } i, j}{dx^j \wedge dx^i}$$

for functions which have continuous mixed partial derivatives we have proved

$$\begin{aligned}D_i D_j \omega_{i_1 \dots i_k} &= D_j D_i \omega_{i_1 \dots i_k} \\ &\Rightarrow \text{property (c)}\end{aligned}$$

(b):

Take

$$\omega \wedge \eta = \sum_{i_1 < \dots < i_k} \sum_{j_1 < \dots < j_c} \omega_{i_1 \dots i_k} \eta_{j_1 \dots j_c} dx^{i_1} \wedge \dots \wedge dx^{i_k} \wedge dx^{j_1} \wedge \dots \wedge dx^{j_c}$$

$$d(\omega \wedge \eta) = \sum_{i_1 < \dots < i_k} \sum_{j_1 < \dots < j_c} \sum_{\alpha=1}^n D_\alpha (\omega_{i_1 \dots i_k} \eta_{j_1 \dots j_c}) dx^\alpha \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \wedge dx^{j_1} \wedge \dots \wedge dx^{j_c}$$

$$\begin{aligned}
&= \sum_{i_1 < \dots < i_k} \sum_{j_1 < \dots < j_c} \sum_{\alpha=1}^n ((D_\alpha \omega_{i_1 \dots i_k}) \eta_{j_1 \dots j_c} + \omega_{i_1 \dots i_k} (D_\alpha \eta_{j_1 \dots j_c})) dx^\alpha \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \wedge dx^{j_1} \wedge dx^{j_c} \\
&= \sum_{i_1 < \dots < i_k} \sum_{j_1 < \dots < j_c} \sum_{\alpha=1}^n (D_\alpha \omega_{i_1 \dots i_k}) \eta_{j_1 \dots j_c} dx^\alpha \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \wedge dx^{j_1} \wedge dx^{j_c} \\
&\quad + \sum_{i_1 < \dots < i_k} \sum_{j_1 < \dots < j_c} \sum_{\alpha=1}^n \omega_{i_1 \dots i_k} (D_\alpha \eta_{j_1 \dots j_c}) dx^\alpha \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \wedge dx^{j_1} \wedge dx^{j_c}
\end{aligned}$$

* shift the dx^α term down to one place in front of the dx^{j_1} term.

$$\begin{aligned}
&= \left(\sum_{i_1 < \dots < i_k} \sum_{\alpha=1}^n D_\alpha \omega_{i_1 \dots i_k} dx^\alpha \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \right) \wedge \left(\sum_{j_1 < \dots < j_c} \eta_{j_1 \dots j_c} dx^{j_1} \wedge dx^{j_c} \right) \\
&\quad + \left(\sum_{i_1 < \dots < i_k} \omega_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k} \right) \wedge (-1)^k \left(\sum_{j_1 < \dots < j_c} \sum_{\alpha=1}^n D_\alpha \eta_{j_1 \dots j_c} dx^\alpha \wedge dx^{j_1} \wedge dx^{j_c} \right) \\
&= d(\omega) \wedge \eta + (-1)^k \omega \wedge d(\eta)
\end{aligned}$$

□

3.6 Closed and Exact forms

Definition 3.27. Let ω be a k -form, ω is called closed if

$$d\omega = 0$$

Definition 3.28. Let ω be a k -form, ω is called exact if \exists a $(k-1)$ -form η such that:

$$d\eta = \omega$$

Proposition 3.2. If ω is exact then it is closed.

Proof.

$$d(\omega) = d(d(\eta)) = 0$$

□

Example 3.23. $n = 2, k = 1$

$$\omega = p(x, y)dx + q(x, y)dy$$

$$d\omega = \left(-\frac{dp}{dy} + \frac{dq}{dx} \right) dx \wedge dy$$

ω is closed if $\frac{dp}{dy} = \frac{dq}{dx}$

Example 3.24. when ω is an exact 1- form we have a $\eta = f$ 0-form such that

$$\omega = d\eta = df = \frac{df}{dx}dx + \frac{df}{dy}dy$$

$$\text{grad}(f) = \frac{df}{dx}\mathbf{i} + \frac{df}{dy}\mathbf{j}$$

Definition 3.29. For a vector field

$$F = P\mathbf{i} + Q\mathbf{j}$$

if

$$\frac{dP}{dy} = \frac{dQ}{dx}$$

we call it a conservative field. Also F is a conservative field if

$$F = \text{grad}(f)$$

Example 3.25.

$$\omega = xy^2dx + ydy$$

$$\begin{aligned} d\omega &= \frac{d(xy^2)}{dy}dy \wedge dx + \frac{d(y)}{dx}dx \wedge dy \\ &= -2xydy \wedge dx \neq 0 \end{aligned}$$

\Rightarrow not closed and not exact

Example 3.26.

$$\omega = xy^2dx + x^2ydy$$

$$d\omega = 2xydy \wedge dx + 2xydx \wedge dy = 0$$

\Rightarrow closed

Is it exact?

$$\exists f = \frac{x^2y^2}{2} + h$$

$$\frac{df}{dx}dx + \frac{df}{dy}dy = xy^2dx + x^2ydy = \omega$$

so it is also exact.

Proposition 3.3. If $n > 2$, $k = 1$

$$\omega = \omega_1dx^1 + \cdots + \omega_ndx^n$$

is closed. Is it exact? i.e does there exist f st

$$\omega = df = D_1f dx^1 + \cdots + D_nf dx^n$$

$\omega_i = D_if$ and can assume $f(0) = 0$ you can recover the f by integration in one variable t . $f : \mathbb{R}^n \rightarrow \mathbb{R}$.

$$f(x) - f(\theta) = \int_0^1 \frac{d}{dt}[f(tx)]dt$$

$$\begin{aligned}
f(x) &= \int_0^1 \sum_{\alpha=1}^n D_{\alpha} f(tx) \frac{d}{dt}(tx^{\alpha}) dt \\
&= \int_0^1 \sum_{\alpha=1}^n D_{\alpha} f(tx) x^{\alpha} dt \\
&= \int_0^1 \sum_{\alpha=1}^n \omega_{\alpha}(tx) x^{\alpha} dt
\end{aligned}$$

Definition 3.30 (Star-Shaped region). *A is a star-shaped region with respect to 0 (or p) if $\forall t \in [0, 1], \forall x \in A$*

$$t \cdot x \in A$$

i.e.

$$p + t(x - p) \in A$$

Lemma 3.10 (Poincaré Lemma). *If A is star-shaped with respect to 0 and ω is a closed form on A then ω is an exact form on A.*

Proof. For any lform ω , i will define an (l-1)-form $I(\omega)$ such that

•

$$I(\lambda\omega_1 + \omega_2) = \lambda I(\omega_1) + I(\omega_2)$$

•

$$I(0) = 0$$

•

$$\underbrace{d(I(\omega))}_{l-1} + \underbrace{I(d(\omega))}_{l+1} = \omega$$

Then if ω is closed, $d\omega = 0$ so $I(d\omega) = 0$ so we get

$$d(I(\omega)) = \omega \Rightarrow \omega \text{ is exact}$$

$$\omega = \sum_{i_1 < \dots < i_l} \omega_{i_1 \dots i_l} dx^{i_1} \wedge \dots \wedge dx^{i_l}$$

$$I(\omega) = \sum_{i_1 < \dots < i_l} \sum_{\alpha=1}^l (-1)^{\alpha-1} \int_0^1 t^{l-1} \omega_{i_1 \dots i_l}(tx) x^{i_{\alpha}} dt dx^{i_1} \wedge \dots \wedge \hat{dx}^{\alpha} \wedge \dots \wedge dx^{i_l}$$

$$I(0) = 0 \quad \text{trivial to show}$$

$$I(\lambda\omega_1 + \omega_2) = \lambda I(\omega_1) + I(\omega_2) \quad \text{trivial to show}$$

□

Definition 3.31. *The set $I^k = [0, 1]^k$ is called the standard k-cube. A continuous map $C : I^k \rightarrow A$, where A is open in \mathbb{R}^n is called a singular k-cube.*


Example 3.27. $k = 1$, $C : [0, 1] \rightarrow A$ (curve)

Example 3.28. $K = 2$, $C : [0, 1]^2 \rightarrow A$ parameterisation of a curve/surface.

Example 3.29. $k = 0$, $[0, 1]^0 = \{0\}$ A singular 0-cube, $\{0\} \rightarrow A$ maps to a point on some surface.

Example 3.30. $k = 1$, $\partial(I^1) = +1, -0$ The motivation for this is $\int_0^1 f'(x)dx = f(1) - f(0)$

Example 3.31. $k = 2$ $I^2 = [0, 1]^2$



$$\partial(I^2) = \gamma_1 + \gamma_2 - \gamma_3 - \gamma_4$$

$$\gamma_1 \quad I_{(2,0)}^2 = \{(x, 0), 0 \leq x \leq 1\}$$


$$\gamma_2 \quad I_{(1,1)}^2 = \{(1, y), 0 \leq y \leq 1\}$$

$$\gamma_3 \quad I_{(2,1)}^2 = \{(x, 1), 0 \leq x \leq 1\}$$

$$\gamma_4 \quad I_{(1,0)}^2 = \{(0, y), 0 \leq y \leq 1\}$$

Note. $I_{(a,b)}^n$ means: fix a^{th} variable and set a^{th} variable to b .

Example 3.32. $k = 3$



$$top : \quad I_{(3,1)}^3 = \{(x, y, 1), 0 \leq x \leq 1, 0 \leq y \leq 1\}$$

$$base : \quad I_{(3,0)}^3 = \{(x, y, 0), 0 \leq x \leq 1, 0 \leq y \leq 1\}$$

$$front : \quad I_{(1,1)}^3 = \{(1, y, z), 0 \leq y \leq 1, 0 \leq z \leq 1\}$$

$$back : \quad I_{(1,0)}^3 = \{(0, y, z), 0 \leq y \leq 1, 0 \leq z \leq 1\}$$

$$left : \quad I_{(2,0)}^3 = \{(x, 0, z), 0 \leq x \leq 1, 0 \leq z \leq 1\}$$

$$right : \quad I_{(2,1)}^3 = \{(x, 1, z), 0 \leq x \leq 1, 0 \leq z \leq 1\}$$

$$\therefore \partial I^3 = I_{(3,1)}^3 - I_{(3,0)}^3 + I_{(1,1)}^3 - I_{(1,0)}^3 + I_{(2,0)}^3 - I_{(2,1)}^3$$

Definition 3.32. Give an n -cube $I^n = [0, 1]^n$, we define the various faces of it to be

$$I_{(i,0)}^n = \{(x^1, x^2, \dots, x^{i-1}, 0, x^{i+1}, \dots, x^n), 0 \leq x^j \leq 1\}$$

$$I_{(i,1)}^n = \{(x^1, x^2, \dots, x^{i-1}, 1, x^{i+1}, \dots, x^n), 0 \leq x^j \leq 1\}$$

We define the boundry of I^n to be

$$\partial I^n = \sum_{i=1}^n \sum_{\alpha=0}^1 (-1)^{i+\alpha} I_{(i,\alpha)}^n$$

We form formal sums of singular n -cubes with integer coefficients. (This is the construction of a certain abelian group, or \mathbb{Z} -module.)

Example 3.33.

$$C_1 : I^1 \rightarrow A$$

$$C_2 : I^1 \rightarrow A$$

$$3C_1 + (-5)C_2$$

These are singular n -chains

Definition 3.33. A singular n -chain, $C \in A$, is a finite linear combination of singular n -cubes with integer coefficients.

$$C = \sum_{j=1}^m m_j C_j \quad m_j \in \mathbb{Z}, C_j : I^n \rightarrow A$$

Definition 3.34. If C is a singular n -cube, $C : I \rightarrow A$. Then

$$\partial C = \sum_{i=1}^n \sum_{\alpha=0}^1 (-1)^{i+\alpha} C(I_{(i,\alpha)}^n)$$

for a singular n -chain

$$C = \sum_{j=1}^m m_j C_j$$

where C_j are singular n -cubes

$$\partial C = \sum_{j=1}^m m_j \partial(C_j)$$

In \mathbb{R}^k we will define integration of a k -form on a k -cube and a k -form on a k -cube face.

Let ω be a k -form on I^k

$$\forall p \in I^k : \omega(p) \in \Lambda^k(\mathbb{R}_p^k)$$

$$\omega = f(x^1, \dots, x^k) dx^1 \wedge \dots \wedge dx^k$$

$$\begin{aligned} \int_{I^k} \omega &= \int_{I^k} f(x^1, \dots, x^k) dx^1 \wedge \dots \wedge dx^k \\ &= \int_{[0,1]^k} f(x^1, \dots, x^k) dx^1 \wedge \dots \wedge dx^k \quad (\text{Riemann Integral}) \end{aligned}$$

By Fubini

$$= \int_0^1 \left(\int_0^1 \left(\cdots \left(\int_0^1 f(x^1, \dots, x^k) dx^1 \right) dx^2 \right) \cdots \right) dx^k$$

this can be evaluated

On \mathbb{R}^k , let η be a $k-1$ form on $I_{(i,\alpha)}^k$. The basis of $k-1$ forms in \mathbb{R}^k is

$$dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k \quad j = 1, \dots, k$$

Assume η is given by:

$$\eta = g(x^1, \dots, x^k) dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k$$

$$\int_{I_{(i,\alpha)}^k} \eta = \begin{cases} \int_{[0,1]^k} g(x^1, \dots, x^{j-1}, \alpha, x^{j+1}, \dots, x^k) dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

Example 3.34.

$$\begin{array}{ccc} \begin{array}{c} \cdot \\ \cdot \text{---} \cdot \\ \cdot \end{array} & \int_{I_{(2,0)}^k} dy = 0 \\ \uparrow & \int_{I_{(1,0)}^k} dx = 0 \\ \cdot & \end{array}$$

If

$$\eta = \sum_{j=1}^n g_j(x^1, \dots, x^k) dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k$$

then

$$\int_{I_{(i,\alpha)}^k} \eta = \sum_{j=1}^n \int_{I_{(i,\alpha)}^k} g_j(x^1, \dots, x^k) dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k$$

If ω is a 0-form, then ω is a function $f(x^1, \dots, x^k)$, 0-cube is the point $\{0\}$. Then

$$\int_{I^0} \omega = f(0, \dots, 0)$$

If

$$C = \sum_{j=1}^m m_j C_j$$

where C_j are all k-cubes standard, then

$$\int_C \omega = \sum_{j=1}^m m_j \int_{C_j} \omega$$

If

$$C = \sum_{j=1}^m m_j C_j$$

where C_j are all k-1 cubes, then

$$\int_C \eta = \sum_{j=1}^m m_j \int_{C_j} \eta$$

$$I^1 = 0 \dashv \vdash 1 \quad \int_{I^1} \omega, \quad \int_{5I^1} \omega = 5 \int_{I^1} \omega$$

$$I^2_{(1,0)} \left[\begin{array}{c} \overline{I^2_{(2,1)}} \\ \overline{I^2_{(2,0)}} \end{array} \right] I^2_{(1,1)}$$

$$\int_{\partial I^2} \overbrace{\eta}^{1-form} = + \int_{I^2_{(2,0)}} \eta + \int_{I^2_{(1,1)}} \eta - \int_{I^2_{(2,1)}} \eta - \int_{I^2_{(1,0)}} \eta$$

Lemma 3.11 (poincaré lemma). *If A is star-shaped with respect to 0 and ω is a closed l -form on A , then ω is exact*

$$\omega = \sum_{i_1 < \dots < i_l} \omega_{i_1 \dots i_l} dx^{i_1} \wedge \dots \wedge dx^{i_l}$$

$$I(\underbrace{\omega}_{l-form}) = \sum_{i_1 < \dots < i_l} \sum_{\alpha} (-1)^{\alpha-1} \int_0^1 t^{l-1} \omega_{i_1 \dots i_l}(tx) dt x^{i_\alpha} dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_\alpha}} \wedge \dots \wedge dx^{i_l}$$

$$I(0) = 0$$

$$I(\omega_1 + \omega_2) = I(\omega_1) + I(\omega_2)$$

$$\boxed{\text{if } \omega \text{ is closed, } d\omega = 0}$$

$$dI(\omega) + I(d\omega) = \omega \quad (unproved)$$

If ω is closed $d\omega = 0$, so

$$dI(\omega) + I(\cancel{\emptyset}) = \omega \quad \boxed{\omega \text{ is exact}}$$

Because I is a linear opperator, it suffices to prove it for

$$\omega = f(x^1, \dots, x^n) dx^{i_1} \wedge \dots \wedge dx^{i_l}$$

$$d\omega = \sum_{\beta=1}^n D_{\beta}[f(x^1, \dots, x^n)] dx^{\beta} \wedge dx^{i_1} \wedge \dots \wedge dx^{i_l}$$

$$I(\omega) = \sum_{\alpha}^l (-1)^{\alpha-1} \int_0^1 t^{l-1} f(tx) dt x^{i_{\alpha}} dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l}$$

$$\begin{aligned} d(I(\omega)) &= \sum_{\beta=1}^n \sum_{\alpha}^l (-1)^{\alpha-1} \int_0^1 t^{l-1} D_{\beta}(f(tx) x^{i_{\alpha}}) dt dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l} \\ &= \sum_{\beta=1}^n \sum_{\alpha}^l (-1)^{\alpha-1} \int_0^1 t^{l-1} (\delta_{\beta_1, i_{\alpha}} f(tx) + x^{i_{\alpha}} (D_{\beta} f)(tx) \cdot t) dt dx^{\beta} \wedge dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l} \\ &= \sum_{\alpha}^l (-1)^{\alpha-1} \int_0^1 t^{l-1} f(tx) dt dx^{i_{\alpha}} \wedge dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l} \\ &\quad + \sum_{\beta=1}^n \sum_{\alpha}^l (-1)^{\alpha-1} \int_0^1 t^l x^{i_{\alpha}} (D_{\beta} f)(tx) dt dx^{\beta} \wedge dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l} \\ &= l \int_0^1 t^{l-1} f(tx) dt dx^{i_1} \wedge \dots \wedge \underbrace{dx^{i_{\alpha}}}_{\text{put back}} \wedge \dots \wedge dx^{i_l} \\ (1) \quad &+ \sum_{\beta=1}^n \sum_{\alpha}^l (-1)^{\alpha-1} \int_0^1 t^l x^{i_{\alpha}} (D_{\beta} f)(tx) dt dx^{\beta} \wedge dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l} \end{aligned}$$

$$\begin{aligned} I(d\omega) &= \sum_{j \in \{\beta, i_1, \dots, i_l\}} \sum_{\beta}^n (-1)^{j-1} \int_0^1 t^l D_{\beta} f(tx) dt x^j (dx^{\beta} \wedge dx^{i_1} \wedge \dots \wedge \widehat{dx^j} \wedge \dots \wedge dx^{i_l}) \\ &= \sum_{\beta}^n \int_0^1 t^l (D_{\beta} f)(tx) dt x^{\beta} dx^{i_1} \wedge \dots \wedge dx^j \wedge \dots \wedge dx^{i_l} \\ (2) \quad &+ \sum_{\alpha=1}^l \sum_{\beta}^n (-1)^{\alpha} \int_0^1 t^l (D_{\beta} f)(tx) dt x^{\alpha} dx^{\beta} \wedge dx^{i_1} \wedge \dots \wedge \widehat{dx^{i_{\alpha}}} \wedge \dots \wedge dx^{i_l} \end{aligned}$$

(1) cancels (2).

$$\begin{aligned}
dI(\omega) + I(d\omega) &= l \int_0^1 t^{l-1} f(tx) dt dx^{i_1} \wedge \cdots \wedge dx^{i_\alpha} \wedge \cdots \wedge dx^{i_l} \\
&\quad + \sum_{\beta}^n (-1)^{j-1} \int_0^1 t^l (D_{\beta} f)(tx) dt x^{\beta} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^{i_l} \\
&= \left(\int_0^1 \left[lt^{l-1} f(tx) + \sum_{\beta}^n t^l x^{\beta} (D_{\beta} f)(tx) \right] dt \right) dx^{i_1} \wedge \cdots \wedge dx^{i_l} \\
&= \int_0^1 \frac{d}{dt} \left(t^l f(tx) \right) dt dx^{i_1} \cdots \wedge dx^{i_l} \\
&= t^l f(tx) \Big|_{t=0}^{t=1} dx^{i_1} \cdots \wedge dx^{i_l} \\
&= f(1 \cdot x) dx^{i_1} \wedge \cdots \wedge dx^{i_l} - 0 \\
&= \omega
\end{aligned}$$

3.7 Stoke's Theorem

Theorem 3.12 (Stoke's Theorem).

$$\int_{\partial C} \omega = \int_C d\omega$$

where ω is a $(k-1)$ -form, $d\omega$ is a k -form, ∂C is a $(k-1)$ singular chain and C is a k -singular chain.

Proof of stokes theorem on \mathbb{R}^k for ω $k-1$ form. $C = I^k$ standard k -cube

$$\int_{\partial I^k} \omega = \int_{I^k} d\omega$$

we know $\int_C \eta$ is linear in η , ie

$$\int_C \lambda \eta_1 + \eta_2 = \lambda \int_C \eta_1 + \int_C \eta_2$$

therefore it suffices to prove it for:

$$\begin{aligned}
\omega &= f(x^1, \dots, x^k) dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k \\
d\omega &= \sum_{\beta=1}^k D_{\beta} f(x^1, \dots, x^k) dx^{\beta} \wedge dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k \\
&= D_j f(x^1, \dots, x^k) dx^j \wedge dx^1 \wedge \cdots \wedge \widehat{dx^j} \wedge \cdots \wedge dx^k \\
&= (-1)^{j-1} D_j f(x^1, \dots, x^k) dx^1 \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^k
\end{aligned}$$

$$\begin{aligned}
\int_{I^k} d\omega &= (-1)^{j-1} \int_{I^k} D_j f dx^1 \wedge \cdots \wedge dx^k \\
&= (-1)^{j-1} \int_{[0,1]^k} D_j f dx^1 \cdots dx^k \\
&= (-1)^{j-1} \int_0^1 \cdots \int_0^1 \left(\int_0^1 D_j f dx^j \right) dx^1 \cdots \widehat{dx^j} \cdots dx^k \\
&= (-1)^{j-1} \int_0^1 \cdots \int_0^1 f(x^1, \dots, x^k) \Big|_{x^j=0}^{x^j=1} dx^1 \cdots \widehat{dx^j} \cdots dx^k \\
&= (-1)^{j-1} \int_0^1 \cdots \int_0^1 f(x^1, \dots, x^{j-1}, 1, x^{j+1}, \dots, x^k) dx^1 \cdots \widehat{dx^j} \cdots dx^k \\
&\quad - (-1)^{j-1} \int_0^1 \cdots \int_0^1 f(x^1, \dots, x^{j-1}, 0, x^{j+1}, \dots, x^k) dx^1 \cdots \widehat{dx^j} \cdots dx^k
\end{aligned}$$

$$\begin{aligned}
\int_{\partial I^k} \omega &= \sum_{i=1}^k \sum_{\alpha=0,1} (-1)^{i+\alpha} \int_{I_{(i,\alpha)}^k} \underbrace{\omega}_{k-1 \text{ form}} \\
\text{only } i=j \text{ remaining} &= \sum_{\alpha=0,1} (-1)^{j+\alpha} \int_{[0,1]^{k-1}} f(x^1, \dots, x^{j-1}, \alpha, x^{j+1}, \dots, x^k) dx^1 \cdots \widehat{dx^j} \cdots dx^k \\
&= (-1)^{j+1} \int_{[0,1]^{k-1}} f(x^1, \dots, x^{j-1}, 1, x^{j+1}, \dots, x^k) dx^1 \cdots \widehat{dx^j} \cdots dx^k \\
&\quad + (-1)^j \int_{[0,1]^{k-1}} f(x^1, \dots, x^{j-1}, 0, x^{j+1}, \dots, x^k) dx^1 \cdots \widehat{dx^j} \cdots dx^k
\end{aligned}$$

□

Definition 3.35 (Pullback $C^*(\omega)$). If ω is a k -form on A containing a singular k -cube C , ($C : I^k \rightarrow A$), then:

$$\int_C \omega = \int_{I^k} C^*(\omega)$$

How do we define the pullback of a k -form on C ?

Remark. If S is linear, $S^*(T) = T \cdot S$ linear

$$\begin{array}{ccc}
V & \xrightarrow{S} & W \\
& \downarrow T & \\
& \mathbb{R} &
\end{array} \quad V, W \text{ vector spaces, } T \text{ linear functional}$$

Recall the Pullback of Tensors: $T \in \mathcal{J}^k(W)$, then $S \in \mathcal{J}^k(W)$

$$S^*(T)(v_1, \dots, v_k) = T(S(v_1), \dots, S(v_k)), \quad v_i \in V$$

Let ω be a k -form on \mathbb{R}^m and let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ $f^*(\omega)(p) \in \Lambda^k(\mathbb{R}_p^n) \quad \forall p \in \mathbb{R}^n$

$$f^*(\omega)(p)(v_1, \dots, v_k) = \omega(f(p))(\underbrace{Df(v_1), \dots, Df(v_k)}_{\in \Lambda^k(\mathbb{R}_{f(p)}^m)})$$

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be differentiable, $p \in \mathbb{R}^n$, $Df(p) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ linear map. It helps us the push-forward of \mathbb{R}_p^n to $\mathbb{R}_{f(p)}^m$. If $v_p \in \mathbb{R}_p^n$, $v_p = (p, v)$, $v \in \mathbb{R}^n$, then $F_*(v_p) \in \mathbb{R}_{f(p)}^m$ defined by

$$f_*(v_p) = (f(p), Df(p)(v))$$

Proposition 3.4. $F_*(v_p) : \mathbb{R}_p^n \rightarrow \mathbb{R}_{f(p)}^m$ is linear.

Proof. If $v_p, w_p \in \mathbb{R}_p^n$, $\lambda \in \mathbb{R}$

$$\begin{aligned} f_*(\lambda v_p + w_p) &= f_*(\lambda(p, v) + (p, w)) = f_*((p, \lambda v + w)) = (f(p), Df(p)(\lambda v + w)) \\ &= (f(p), \lambda Df(p)(v) + Df(p)(w)) = \lambda(f(p), Df(p)(v)) + (f(p), Df(p)(w)) \\ &= \lambda f_*(v_p) + f_*(w_p) \end{aligned}$$

□

Definition 3.36. If $T \in \mathcal{J}^k(\mathbb{R}_{f(p)}^m)$ then $f^*(T) \in \mathcal{J}^k(\mathbb{R}_p^n)$ will be defined by

$$f^*(\omega)(p)(v_1, \dots, v_k) = \omega(f(p))(f_*(v_1), \dots, f_*(v_k)) \quad \forall p \in \mathbb{R}^n, v_i \in \mathbb{R}_p^n$$

Theorem 3.13. $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ differentiable

$$(i) \quad f^*(dx^i) = \sum_{j=1}^n D_j f^i dx^j$$

$$(ii) \quad f^*(\lambda w_1 + w_2) = \lambda f^*(w_1) + f^*(w_2)$$

$$(iii) \quad f^*(g \cdot \omega) = (g \circ f)f^*(\omega)$$

$$(iv) \quad f^*(\omega \wedge \eta) = f^*(\omega) \wedge f^*(\eta)$$

Proof.

Proof of (i):

$$f^*(dx^i) = \sum_{j=1}^n D_j f^i dx^j$$

Take $p \in \mathbb{R}^n$, $f^*(dx^i)(p) \in \Lambda^1(\mathbb{R}_p^n)$

$$f^*(dx^i)(p)(v_p) = dx^i(f(p))(f_*(v_p)) = dx^i(f(p))(f(p), Df(p)(v))$$

dx^i picks up the i^{th} component of the vector:

$$(f(p), Df(p)(v))^i = \sum_{j=1}^n D_j f^i(p) v_j$$

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$Df(p)(v) = f'(p) \cdot v = \begin{pmatrix} D_1 f^1 & \dots & D_n f^1 \\ \vdots & & \vdots \\ \hline i^{th} \text{ row} \\ \hline \vdots & & \vdots \\ D_1 f^m & \dots & D_n f^m \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ \vdots \\ \vdots \\ \vdots \\ v_n \end{pmatrix} = \sum_{j=1}^n D_j f^i v_j$$

compute:

$$\left(\sum_{j=1}^n D_j f^i dx^j \right) (p)(v_p) = \sum_{j=1}^n D_j f^i(p) \underbrace{dx^j(p)}_{\text{Picks up } j^{th} \text{ component}}(v_p) = \sum_{j=1}^n D_j f^i(p) v_j$$

Proof of (iii):

$$F^*(g \cdot \omega) = g \circ f \cdot f^*(\omega)$$

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m, g : \mathbb{R}^m \rightarrow \mathbb{R}, p \in \mathbb{R}^n, v_1, \dots, v_k \in \mathbb{R}_p^n$$

$$\begin{aligned} F^*(g \cdot \omega)(p)(v_1, \dots, v_k) &= (g\omega)(f(p))(f_*(v_1), \dots, f_*(v_k)) \\ &= g(f(p)) \cdot \omega(f(p))(f_*(v_1), \dots, f_*(v_k)) \end{aligned}$$

Compute:

$$g \circ f \cdot f^*(\omega)(p)(v_1, \dots, v_k) = (g \circ f)(p)\omega(f(p))(f_*(v_1), \dots, f_*(v_k))$$

□

Example 3.35. ω a 1-form in \mathbb{R}^3

$$\omega = P(x, y, z)dx + Q(x, y, z)dy + R(x, y, z)dz$$

$f : [0, 1] \rightarrow \mathbb{R}^3$ (parameterises curve in \mathbb{R}^3) $f^*(\omega)$ 1-form on $[0, 1]$

Let v_t be a tangent vector on \mathbb{R}_t^1 , $v_t = (t, v)$

$$\begin{aligned} f^*(\omega)(t)(v_t) &= \omega(f(t))(f_*(v_t)) \\ &= (Pdx + Qdy + Rdz)(f(t))(f_*(v_t)) \\ &= P(f(t))dx(f(t))(f_*(v_t)) + Q(f(t))dy(f(t))(f_*(v_t)) + R(f(t))dz(f(t))(f_*(v_t)) \end{aligned}$$

$$\begin{aligned} f^*(v_t) &= (f(t), Df(t)(v)) \\ &= (f(t), (Df^1(t)(v), Df^2(t)(v), Df^3(t)(v))) \\ f &= f^1 + f^2 + f^3 \end{aligned}$$

$$= P(f(t))Df^1(t)(v) + Q(f(t))Df^2(t)(v) + R(f(t))Df^3(t)(v)$$

$$\Rightarrow f^*(\omega) = (P \circ f) \frac{df^1}{dt} dt + (Q \circ f) \frac{df^2}{dt} dt + (R \circ f) \frac{df^3}{dt} dt$$

$$\begin{aligned} f^*(\omega) &= f^*(Pdx + Qdy + Rdz) \\ &= (P \circ f)f^*(dx) + (Q \circ f)f^*(dy) + (R \circ f)f^*(dz) \\ &= (P \circ f)Df^1 dt + (Q \circ f)Df^2 dt + (R \circ f)Df^3 dt \end{aligned}$$

Definition 3.37. If $C : I^k \rightarrow A$ is a singular k -cube in A and ω is a k -form on A , then:

$$\int_C \omega = \int_{I^k} C^*(\omega)$$

Example 3.36. ω 1-form on \mathbb{R}^2 ,

$$\omega = xdy$$

$$C : [0, 1] \rightarrow \mathbb{R}^2,$$

$$C(t) = (a \cos(2\pi t), b \sin(2\pi t)), \quad a, b > 0.$$

$$\begin{aligned} \int_C xdy &= \int_{[0,1]} C^*(xdy) = \int_0^1 (x \cdot C)(t) \frac{dC^2}{dt} dt = \int_0^1 a \cos(2\pi t) \cdot b 2\pi \cos(2\pi t) dt \\ &= \int_0^1 ab 2\pi \cdot \cos^2(2\pi t) dt = ab\pi \int_0^1 1 + \cos(4\pi t) dt = \pi ab \end{aligned}$$

Stokes Theorem:

$$\int_C \omega = \int_{\tilde{C}} d\omega = \int_{\tilde{C}} d(xdy) = \int_{\tilde{C}} dx \wedge dy \quad \leftarrow \text{area of reigon parameterised by } \tilde{C}$$

Call \tilde{C} the inside of the ellipse (2-cube)

$$\tilde{C}(u, t) = (au \cos(2\pi t), bu \sin(2\pi t)), \quad t \in [0, 1], \quad u \in [0, 1]$$

$$\partial \tilde{C} = C$$

Definition 3.38. If $C : I^k \rightarrow A$ is a singular k -cube in A and ω is a k -form on A then,

$$\int_C \omega = \int_{I^k} C^*(\omega)$$

If C is a singular k -chain, ie

$$C = \sum_{j=1}^m m_j C_j \quad m_j \in \mathbb{Z}, \quad C_j \text{ singular } k - \text{cubes}$$

then

$$\int_C \omega = \sum_{j=1}^m m_j \int_{C_j} \omega = \sum_{j=1}^m m_j \int_{I^k} C_j^*(\omega)$$

Theorem 3.14 (Stoke's Theorem for Singular k-chains in \mathbb{R}^k). *If ω is a $(k-1)$ -form on \mathbb{R}^k , $d\omega$ is a k -form on \mathbb{R}^k , C a k -singular chain, ∂C a $(k-1)$ -singular chain. Then:*

$$\int_{\partial C} \omega = \int_C d\omega$$

$$C = \sum_{j=1}^m m_j C_j \quad m_j \in \mathbb{Z}, \quad C_j \text{ singular } k\text{-cube}, \quad C_j : I^k \rightarrow \mathbb{R}^k$$

$$\partial C_j = C_j(\partial I^k) = \sum_{i=1}^k \sum_{\alpha=0,1} (-1)^{i+\alpha} C_j(I_{(i,\alpha)}^k)$$

Proof.

$$\sum_{j=1}^m \sum_{i=1}^k \sum_{\alpha=0,1} m_j (-1)^{i+\alpha} \int_{C_j(I_{(i,\alpha)}^k)} \stackrel{Def^n}{=} \sum_{j=1}^m \sum_{i=1}^k \sum_{\alpha=0,1} \left(m_j (-1)^{i+\alpha} \int_{I_{(i,\alpha)}^k} C_j^*(\omega) \right) \quad (1)$$

Now compute

$$\int_C d\omega = \sum_{j=1}^m m_j \int_{C_j} d\omega \stackrel{Def^n}{=} \sum_{j=1}^m m_j \int_{I^k} C_j^*(d\omega) = \sum_{j=1}^m m_j \int_{I^k} d(C_j^*(\omega))$$

since

$$\boxed{d(C_j^*(\omega)) = C_j^*(d\omega)}$$

Now apply Stoke's theorem for standard k-cube

$$= \sum_{j=1}^m m_j \int_{\partial I^k} C_j^*(\omega) = \sum_{j=1}^m m_j \sum_{i=1}^k \sum_{\alpha=0,1} (-1)^{i+\alpha} \int_{I_{(i,\alpha)}^k} C_j^*(\omega). \quad (2)$$

$$\therefore (1) = (2)$$

□

3.8 Classical Stoke's Theorem in \mathbb{R}^2

$$\int_{\gamma} P(x, y)dx + Q(x, y)dy = \iint_D \left(-\frac{dP}{dy} + \frac{dQ}{dx} \right) dx dy$$

s^1, s^2 are axes for the 2-cube

$$C(s^1, s^2) = (C^1(s^1, s^2), C^2(s^1, s^2))$$

$$\gamma : [0, 1] \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\gamma^1(t), \gamma^2(t))$$

$$\partial C \stackrel{Def^n}{=} C(\partial I^2) = \gamma$$

$$\begin{aligned}
\int_{\gamma} Pdx + Qdy &= \int_{C(\partial I^2)} Pdx + Qdy \stackrel{Defn}{=} \int_{\partial I^2} C^*(Pdx + Qdy) \\
&= \int_{\partial I^2} P(C^1(s^1, s^2), C^2(s^1, s^2)) C^*(dx) + Q(C^1(s^1, s^2), C^2(s^1, s^2)) C^*(dy) \\
&= \int_{\partial I^2} P \frac{d\gamma^1}{dt} dt + Q \frac{d\gamma^2}{dt} dt = \int_{\partial I^2} \left[P(\gamma^1(t), \gamma^2(t)) \frac{d\gamma^1}{dt} + Q(\gamma^1(t), \gamma^2(t)) \frac{d\gamma^2}{dt} \right] dt \\
&\quad \int_{\substack{\gamma \\ = \partial C}} Pdx + Qdy \stackrel{\text{Stoke's}}{\underset{\text{in } \mathbb{R}^2}{=}} \int_C \underbrace{d(Pdx + Qdy)}_{d\omega}
\end{aligned}$$

where γ is the the singular 1-cube which is the boundary of $C(I^k)$

$$\therefore \int_{\partial C} \omega = \int_C d\omega$$

$$\begin{aligned}
\int_C d(Pdx + Qdy) &= \int_C \cancel{P_x dx \wedge dx} + P_y dy \wedge dx + Q_x dx \wedge dy + \cancel{Q_y dy \wedge dy} \\
&= \int_C -\frac{dP}{dy} dx \wedge dy + \frac{dQ}{dy} dx \wedge dy = \int_C \left(-\frac{dP}{dy} + \frac{dQ}{dy} \right) dx \wedge dy \\
&\stackrel{defn}{=} \int_{I^2} C^* \left(-\frac{dP}{dy} + \frac{dQ}{dy} \right) dx \wedge dy \\
&= \int_{I^2} \left[-\frac{dP}{dy}(C^1, C^2) + \frac{dQ}{dy}(C^1, C^2) \right] C^*(dx \wedge dy) \quad (*)
\end{aligned}$$

Proposition 3.5 (What is $C^*(dx \wedge dy)$?).

$$\begin{aligned}
C^*(dx \wedge dy) &= C^*(dx) \wedge C^*(dy) \\
&= \left(\frac{dC^1}{ds^1} ds^1 + \frac{dC^1}{ds^2} ds^2 \right) \wedge \left(\frac{dC^2}{ds^1} ds^1 + \frac{dC^2}{ds^2} ds^2 \right) \\
&= \frac{dC^1}{ds^1} \frac{dC^2}{ds^2} ds^1 \wedge ds^2 + \frac{dC^1}{ds^2} \frac{dC^2}{ds^1} ds^2 \wedge ds^1 \\
&= \left(\frac{dC^1}{ds^1} \frac{dC^2}{ds^2} - \frac{dC^1}{ds^2} \frac{dC^2}{ds^1} \right) ds^1 \wedge ds^2 = \det [C'(s^1, s^2)] ds^1 \wedge ds^2
\end{aligned}$$

where

$$C'(s^1, s^2) = \begin{bmatrix} \frac{dC^1}{ds^1} & \frac{dC^1}{ds^2} \\ \frac{dC^2}{ds^1} & \frac{dC^2}{ds^2} \end{bmatrix}$$

$$(*) = \int_{I^2} \left[-\frac{dP}{dy}(C^1, C^2) + \frac{dQ}{dy}(C^1, C^2) \right] \det (C'(s^1, s^2)) ds^1 \wedge ds^2$$

\therefore ordinary double integral

Recall the change of variables formula for n-dim integrals (2 in this case):

$A \subseteq \mathbb{R}^n$, $g : A \rightarrow \mathbb{R}^n$ injective and differentiable $\det[g'(x)] \neq 0 \forall x \in A$. If $f : g(A) \rightarrow \mathbb{R}$ is integrable

$$\int_{g(A)} f = \int_A (f \circ g) |\det g'|$$

Assuming that anticlockwise orientation it can be shown that $\det(C'(s^1, s^2)) > 0$

$$\int_{I^2} \left(-\frac{dP}{dy}(C^1, C^2) + \frac{dQ}{dy}(C^1, C^2) \right) \det(C'(s^1, s^2)) ds^1 ds^2 = \int_D \left(-\frac{dP}{dy}(x, y) + \frac{dQ}{dy}(x, y) \right) dx dy$$

Theorem 3.15 (Gauss or Divergence Theorem). *Solid T in \mathbb{R}^3 with boundary surface S and a vector function $F = (F^1, F^2, F^3)$*

$S_x \equiv$ Tangent plane to the solid at point $x \in S$

$n(x) \equiv$ Outward unit normal vector

$$\int_S \langle \vec{F}, \vec{n} \rangle dA = \iiint_T (\text{div } \vec{F}) dx dy dz$$

S_x has dim 2 (tangent plane at x)

$$\dim \Lambda^2(S_x) = 1$$

$v, w \in S_x$

$\omega(v, w) = (v \times w) \cdot n = \langle v \times w, n \rangle$ Scalar triple product

$$\omega(v, w) = \begin{vmatrix} v^1 & v^2 & v^3 \\ w^1 & w^2 & w^3 \\ n^1 & n^2 & n^3 \end{vmatrix} = \begin{vmatrix} n^1 & n^2 & n^3 \\ v^1 & v^2 & v^3 \\ w^1 & w^2 & w^3 \end{vmatrix}$$

Choose $\bar{a}, \bar{b} \in S_x$ such that $\bar{a}, \bar{b}, \bar{n}$ are an orthonormal system, right-handed.

Notation (Orthonormal System). Call $\omega(v, w) = dA(v, w)$ or $\omega = dA$ where $\omega(\bar{a}, \bar{b}) = 1$

Theorem 3.16.

$$dA = n^1 dy \wedge dz + n^2 dz \wedge dx + n^3 dx \wedge dy$$

Proof.

$$dA(v, w) = \begin{vmatrix} n^1 & n^2 & n^3 \\ v^1 & v^2 & v^3 \\ w^1 & w^2 & w^3 \end{vmatrix} = n^1(v^2 w^3 - w^2 v^3) + n^2(-v^1 w^3 + v^3 w^1) + n^3(v^1 w^2 - v^2 w^1)$$

$$\begin{aligned}
(dy \wedge dz)(v, w) &= (dy \otimes dz - dz \otimes dy)(v, w) \\
&= dy(v)dz(w) - dz(v)dy(w) \\
&= v^2w^3 - w^2v^3
\end{aligned}$$

$$(dx \wedge dz)(v, w) = -v^1w^3 + v^3w^1$$

$$(dx \wedge dy)(v, w) = v^1w^2 - v^2w^1$$

□

Theorem 3.17.

$$n^1 dA = dy \wedge dz$$

$$n^2 dA = dz \wedge dx$$

$$n^3 dA = dx \wedge dy$$

Proof.

$$(*) \quad (dy \wedge dz)(v, w) = v^2w^3 - w^2v^3 \quad \text{where } v, w \in S_x$$

Since v and w are perpendicular to n , then $v \times w = \lambda n$, $\lambda \in \mathbb{R}$.

$$n^1 dA(v, w) = n^1(\lambda n, n) = n^1 \lambda \quad \text{since } |n| = 1$$

$$\langle v \times w, i \rangle = \langle \lambda n, i \rangle = \lambda n^1$$

$$\langle v \times w, i \rangle = \begin{vmatrix} i & j & k \\ v^1 & v^2 & v^3 \\ w^1 & w^2 & w^3 \end{vmatrix} \cdot i = v^2w^3 - w^2v^3 = (dy \wedge dz)(v, w) \quad \text{by } (*)$$

$$\therefore n^1 dA = dy \wedge dz$$

Similarly for other equations.

□

Proof of Divergence Theorem. Given $\bar{F} = (F^1, F^2, F^3) = F^1 \underline{i} + F^2 \underline{j} + F^3 \underline{k}$

$$\text{div}(\bar{F}) = \frac{dF^1}{dx} + \frac{dF^2}{dy} + \frac{dF^3}{dz}$$

To \bar{F} assign the 2-form:

$$\eta = F^1 dy \wedge dz + F^2 dz \wedge dx + F^3 dx \wedge dy \quad (2\text{-form})$$

Calculate:

$$d\eta = \frac{dF^1}{dx} dx \wedge dy \wedge dz + \frac{dF^2}{dy} dy \wedge dz \wedge dx + \frac{dF^3}{dz} dz \wedge dx \wedge dy$$

$$_{3\text{-form}} d\eta = \left(\frac{dF^1}{dx} + \frac{dF^2}{dy} + \frac{dF^3}{dz} \right) dx \wedge dy \wedge dz$$

$$\int_T d\eta = \int_T (\text{div } F) dx \wedge dy \wedge dz \stackrel{*}{=} \iiint_T f \text{div } F \cdot dx dy dz$$

* Change of variables for I^3 to T . By Stoke's theorem

$$\begin{aligned}
\int_T d\eta &= \int_{\partial T} \eta = \int_S F^1 dy \wedge dz + F^2 dz \wedge dx + F^3 dx \wedge dy \\
&= \int_S F^1 n^1 dA + F^2 n^2 dA + F^3 n^3 dA \\
&= \int_S (F^1 n^1 + F^2 n^2 + F^3 n^3) dA \\
&= \int_S \bar{F} \cdot \bar{n} dA
\end{aligned}$$

□

Recall.

Definition 3.39. A set M is a K -dim manifold in \mathbb{R}^n if the following condition (M) holds. For every $x \in M$:

(M): There exists two open sets U, V of \mathbb{R}^n , $x \in U$ and a diffeomorphism $h : U \rightarrow V$ such that:

$$h(U \cap M) = \{y \in V \text{ s.t. } y^{k+1} = y^{k+2} = \dots = y^n = 0\}$$

Definition 3.40. A subset M of \mathbb{R}^n is a k -dimensional manifold iff for every point $x \in M$ the following holds:

(C) There exists an open set $U \in \mathbb{R}^n$, $x \in U$ and an open set $W \subset \mathbb{R}^k$ and an injective differentiable map $f : W \rightarrow \mathbb{R}^n$ such that

$$(i) \quad f(W) = U \cap M$$

$$(ii) \quad \text{rank } f'(y) = k \quad \forall y \in W$$

$$(iii) \quad f^{-1} : f(W) \rightarrow W \text{ is continuous.}$$

Definition 3.41. A subset M of \mathbb{R}^n is a k -dimensional manifold with boundary if $\forall x \in M$ either (M) holds or (exclusive) (M') holds

(M') \exists open set U of \mathbb{R}^n containing x , an open set V contained in \mathbb{R}^n and a diffeomorphism $h : U \rightarrow V$ such that

$$h(U \cap M) = V \cap (\mathbb{H}^k \times \{0\}) = \{y \in V : y^k \geq 0, y^{k+1} = y^{k+2} = \dots = y^n = 0\}.$$

Moreover, $h^k(x) = 0$. The set of points where condition (M') holds is called the boundary of M and is denoted by ∂M .

Definition 3.42. $\forall v \in \mathbb{R}^k$

$$(a, v) \rightarrow (f(a), Df(a)(v)) \in \mathbb{R}_{f(a)}^n = \mathbb{R}_x^n$$

(a, v) is pushed forward to give a vector in \mathbb{R}_x^n . $f : W \rightarrow U \cap M$, $v_a = (a, v)$. Let $a \in W$ such that $f(a) = x$ then

$$f_*(v_a) = (x, Df(a)(v)) \in \mathbb{R}_x^n$$

$\mathbb{R}_a^k = \{(a, v) : v \in \mathbb{R}^k\}$ a vector space.

Definition 3.43. The tangent space of M at x is defined to be $M_x = f_*(\mathbb{R}_a^k)$ $\dim M_x = k$, given $x = f(a)$, f is a chart.

Definition 3.44. A Vector field on M is a function F on M such that $\forall x \in M$:

$$F(x) \in M_x.$$

Let $x = f(a)$, $f : W \rightarrow U$ $f(W) = U \cap M$. Let $G(a) \in \mathbb{R}_a^k$ such that:

$$f_*(G(a)) = F(f(a)) = F(x)$$

$G(a)$ is unique, since $f_* : \mathbb{R}_a^k \rightarrow M_x$ is injective

Definition 3.45. A vector field on M is called continuous (or differentiable) if $\forall x \in M$ the vector field G on W is continuous (or differentiable).

Definition 3.46. ω is a (differential) p -form on M if $\forall x \in M$

$$\omega(x) \in \Lambda^p(M_x)$$

Then $f^*(\omega)$ is (differentiable) p -form on W .

If $f^*(\omega)$ is differential then ω is differential on $W \subseteq \mathbb{R}^k$.

Definition 3.47. If ω is a p -form on M which k -dim in \mathbb{R}^n , $x \in M$.

$$\omega(x) = \sum_{i_1 < \dots < i_p} \omega_{i_1, \dots, i_p}(x) dx^{i_1} \wedge \dots \wedge dx^{i_p}$$

ω is continuous if $f^*(\omega)$ is continuous on W

ω is differentiable if $f^*(\omega)$ is differentiable on W

We have difficulty with $D_j(\omega_{i_1 \dots i_p})$ since $\omega_{i_1 \dots i_p}$ is not defined on an open set $U \ni x$

Theorem 3.18. Given a differential p -form on M which is k -dim in \mathbb{R}^n , there exists a unique differential $(p+1)$ -form $d\omega$ on M such that $\forall x \in M$

$$d(f^*(\omega)) = f^*(d\omega)$$

and $f : W \rightarrow U \cap M$ is a chart. $d(\omega) \in \Lambda^{p+1}(M_x)$, $v_i \in M_x$, $d\omega(x)(v_1, \dots, v_{p+1})$. Since f_* is a bijection, $f_* : \mathbb{R}_a^k \rightarrow M_x$, \exists unique vectors $w_1, \dots, w_{p+1} \in \mathbb{R}_a^k$ such that

$$f_*(w_i) = v_i$$

$$d\omega(x)(v_1, \dots, v_{p+1}) = d \underbrace{f^*(\omega)(a)}_{\in \Lambda^{p+1}(\mathbb{R}_a^k)}(w_1, \dots, w_{p+1})$$

Aim. To understand Stoke's theorem for M k -dim manifold in \mathbb{R}^n with boundry ∂M

$$\int_{\partial M} \omega + \int_M d\omega$$

where ω is a differential $(k-1)$ -form on M .

Definition 3.48 (Orientation on Vector spaces).

$$\begin{aligned} \text{Bases} : \mathcal{F} &= \{v_1, \dots, v_n\} \\ \mathcal{B} &= \{w_1, \dots, w_n\} \end{aligned}$$

We say \mathcal{F} and \mathcal{B} define the same orientation if $\det[Id]_{\mathcal{F}}^{\mathcal{B}} > 0$ and opposite orientation if $\det[Id]_{\mathcal{F}}^{\mathcal{B}} < 0$. Also we have

$$\det[Id]_{\mathcal{F}}^{\mathcal{B}} = [\det[Id]_{\mathcal{B}}^{\mathcal{F}}]^{-1}$$

$\mathcal{F} \sim \mathcal{B}$ iff they have the same orientation. This is an equivalence relation.

Standard orientation on \mathbb{R}^n : $\mathcal{F} = [e_1, \dots, e_n]$

Example 3.37. $\{e_1, e_2\}$ has opposite orientation to $\{e_2, e_1\}$

This standard orientation is denoted

$$\mu = [e_1, \dots, e_n]$$

When $f(a) = x$ on \mathbb{R}_a^k we have the standard basis $[(e_1)_a, \dots, (e_k)_a]$.

Basis for M_x : $[f_*((e_1)_a), \dots, f_*((e_k)_a)]$

$$\mu_x = [f_*((e_1)_a), \dots, f_*((e_k)_a)]$$

If $b \in W$, then

$$\mu_{f(b)} = [f_*((e_1)_b), \dots, f_*((e_k)_b)]$$

If we have $z = f(c)$ and $z = f(d)$ we assign two orientations at z

$$[f_*((e_1)_c), \dots, f_*((e_k)_c)] = [g_*((e_1)_d), \dots, g_*((e_k)_d)]$$

If the two orientations are equal, ie $\det(Id) > 0$ on these two bases, then we say f and g define consistent orientations at point z . Hopefully this is true on $f(\omega) \cap g(\omega')$ then we call the two orientations consistent.

If there exists consistent orientation on all of M , we say that M is orientable and the manifold is oriented once we fix orientation. If S is a surface and the manifold is oriented once we fix orientation. If S is a surface in \mathbb{R}^3 which is orientable, let

$$\mu_x = [v_1, v_2] \quad x \in S \text{ (2-manifold)}$$

Draw the line perpendicular to S_x at x . Pick a unit vector in $n(x)$ such that $[n(x), v_1, v_2]$ is that standard orientation in \mathbb{R}^3 , then $n(x)$ is the outer unit normal.

M is a k -dim manifold with boundry in \mathbb{R}^3 . $(\partial M)_x$ has a basis

$$[f_*((e_1)_a), \dots, f_*((e_{k-1})_a)]$$

then let $v_0 \in \mathbb{R}_a^k$ such that $f_*(v_0)$ is perpendicular at B , then $|f_*(v_0)| = 1$ and $n(x) = f_*(v_0)$

Definition 3.49 (Integrals). Let C be a singular p -cube on m k -dim manifold. $C : I^k \rightarrow M$. Let ω be a p -form on M . We define:

$$\int_C \omega \stackrel{\text{pullback}}{=} \int_{I^k} C^*(\omega)$$

If C is a k -cube in M a k -manifold and $I^k \subseteq W$, $f : W \rightarrow U \cap M$ is the chart and $C(x) = f(x) \forall x \in I^k$. if f is perserving orientation, then we say C is orientation preserving singular k -cube on M . If ω is a k -form on M with $\omega(y) = 0, \forall y \in C(I^k)$ then we define

$$\int_M \omega \stackrel{\text{Defn}}{=} \int_C \omega$$

$$\boxed{d(f^*(\omega)) = f^*(d\omega) \quad f_1^*(\omega) \text{ a } k-1 \text{ - form on } \mathbb{R}^k}$$

can partition W in to sections W_i then

$$\int_M \omega = \int_{f(W_1)} \omega + \dots + \int_{f(W_i)} \omega + \dots$$

use partitions of unity to define $\int_M \omega$ k -form, $\int_{\partial M} \eta$ $(k-1)$ -form

Theorem 3.19. Let M be a compact, oriented k -manifold with boundry ∂M and ω be a differential $(k-1)$ -form on M , then:

$$\int_{\partial M} \omega = \int_M d\omega$$

Proof.

Recall (Classical Stoke's theorem). • M is a oriented 2-dim manifold with boundary

• F differentiable vector field on M

$$\int_{\partial M} \vec{F} \cdot \vec{T} ds = \int_M \text{curl} \vec{F} \cdot \vec{n} dA$$

Let M be a compact orientated, 2-dim manifold with bounry ∂M in \mathbb{R}^3 . Let T be a vector field on ∂M such that $ds(M) = 1$ where ds is the length element of ∂M , \vec{F} be a differentiable vector fied on an open set containing M , \vec{n} be the outer unit normal on M . Then

$$\int_{\partial M} \vec{F} \cdot \vec{T} ds = \int_M \text{curl} \vec{F} \cdot \vec{n} dA$$

if

$$F = (F^1, F^2, F^3) = F^1 \underline{i} + F^2 \underline{j} + F^3 \underline{k}$$

we define 1-form

$$\omega = F^1 dx + F^2 dy + F^3 dz$$

then we calculate

$$\begin{aligned} d\omega &= \frac{dF^1}{dy} dy \wedge dx + \frac{dF^1}{dz} dz \wedge dx + \frac{dF^2}{dx} dx \wedge dy + \frac{dF^2}{dz} dz \wedge dy + \frac{dF^3}{dx} dx \wedge dz + \frac{dF^3}{dy} dy \wedge dz \\ &= G^1 dy \wedge dz + G^2 dz \wedge dx + G^3 dx \wedge dy \end{aligned}$$

$$G^1 \underline{i} + G^2 \underline{j} + G^3 \underline{k} = \text{curl}(F)$$

We Know

$$n^1 dA = dy \wedge dz$$

$$n^2 dA = dz \wedge dx$$

$$n^3 dA = dx \wedge dy$$

$$\begin{aligned} \int_M G^1 dy \wedge dz + G^2 dz \wedge dx + G^3 dx \wedge dy &= \int_M (G^1 n^1 + G^2 n^2 + G^3 n^3) dA \\ &= \int_M \vec{G} \cdot \vec{n} dA \stackrel{\text{Defn of } G}{=} \int_M \text{curl}(F) \cdot \vec{n} \end{aligned}$$

According to Stoke's general theorem

$$\int_{\partial M} \omega = \int_M d\omega = \int_M \text{curl}(\vec{F}) \cdot \vec{n} dA$$

Since $ds(T) = 1$, we can prove as in that

$$dx = T^1 ds$$

$$dy = T^2 ds$$

$$dz = T^3 ds$$

$$\begin{aligned} \int_{\partial M} \omega &= \int_{\partial M} F^1 dx + F^2 dy + F^3 dz \\ &= \int_{\partial M} F^1 T^1 ds + F^2 T^2 ds + F^3 T^3 ds \\ &= \int_{\partial M} \vec{F} \cdot \vec{T} ds \end{aligned}$$

□