YG OBSTETRIC THEATRE					
Patient Status	**Procedure**	**Staff**	**PPE Level**	**Duration**	**Notes**
COVID - 19 positive and COVID – 19 suspected patients: ➢ Febrile, temp >37.8˚C *or* ➢ Symptomatic *or* ➢ *or* ➢ awaiting test result. Minimum staff in theatre Asymptomatic birthing partner can be present as per normal cases	General Anaesthetic for any procedure	All	**RED**	Whole Case	All obstetric GA are intubated. Aerosol takes 20min to go. During intubation by anaesthetist and ODP the remaining staff will stay in the changing room corridor
	Regional anaesthesia for: e.g. C/S, Trial, Manual removal of placenta, suturing, Cervical suture	All	**GREEN** **Scrubbed in Sterile Green** **Others in Non-sterile Green**	Whole case	Ensure epidural or spinal is working well. If in doubt and time critical give GA. PPE is then RED for all. See note 2 below.
Low Risk of COVID – 19: Temp <37˚8 and No cough, respiratory symptoms *or* COVID – 19 test negative	General Anaesthetic or Spinal or Epidural cases	All staff	**GREEN**	Whole case	As in pre-COVID era

Notes

1. Bleep for Category 1 Section should state "COVID" if this is the case to instruct all staff.
2. Avoid GA if possible. Likelihood of needing to convert from regional to general at same time as surgeon needing to stay at the table is very small. The mother bleeding is not an indication for a GA. Hence if need to convert it is very likely there will be time for all to don PPE FFP3.
3. At all Team Briefs discuss PPE requirements for each case.
4. Use suction on diathermy for all (?)COVID
5. Always observe donning and de-gowning procedures.
6. Avoid staff movement from (red) theatre to green areas unless absolutely necessary
7. Pens to remain with (?)COVID-case, not staff

PPE COLOUR CODING	
RED	FFP3 Mask, visor, surgical gown, long gloves over gown, theatre hat
GREEN	**Sterile**: Surgical facemask, visor/goggles, sterile surgical gown, gloves, theatre hat. **Non-sterile:** Type IIR fluid resistant surgical facemask, plastic apron and gloves, theatre hat. Eye protection (visor or goggles) to be worn for those who will be within 2m of Aerosol Generating Procedure or potential for being splashed by pulse lavage, blood or body fluids.

YG LABOUR WARD					
Patient Status	**Procedure**	**Staff**	**PPE Level**	**Duration**	**Notes**
COVID - 19 positive and COVID – 19 suspected patients: ➢ Febrile, temp >37.8˚C *or* ➢ Symptomatic *or* ➢ awaiting test result. Asymptomatic birthing partner can be present with Surgical face mask	First and second stage labour	All	**GREEN**	Whole case	Non-sterile to protect staff
	Epidural	Anaesthetist	**GREEN**	Siting of Epidural	Sterile to protect patient
Low Risk of COVID – 19: Temp <37˚8 and No cough, no respiratory symptoms *or* Negative COVID – 19 test	First and Second Stage and Epidurals	All	**GREEN**	Whole case	As in pre-COVID era

References:

1. PHW Recommended PPE for healthcare workers by secondary care inpatient clinical setting, NHS and independent sector (April 2nd 2020) https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/877728/T1_Recommended_PPE_for_healthcare_workers_by_secondary_care_clinical_context_poster.pdf
2. Advice regarding PPE to be worn when managing pregnant women with known or suspected COVID – 19 (ICM, ICS, AAGBI, RCOA 20TH March 2020) https://static1.squarespace.com/static/5e6613a1dc75b87df82b78e1/t/5e74a80a1ff9b93077d05e4e/1584703502911/PPE-infographic-20.03.20.pdf
3. Guidance for antenatal and postnatal services in the evolving coronavirus (COVID-19) pandemic (RCOG 30th March 2020) https://www.rcog.org.uk/globalassets/documents/guidelines/2020-03-30-guidance-for-antenatal-and-postnatal-services-in-the-evolving-coronavirus-covid-19-pandemic-20200331.pdf