Sugammadex & Contraception

YG CG 442 MAY 2021

REVIEW BY: DEC 2027

Author(s): Dr N Boyer
Reviewer: Departmental Guideline Group

Key Points

- Sugammadex interacts with progesterone reducing the effectiveness of hormone-based contraception
- Women of child-bearing age using progesterone-containing contraception must be counselled
- Patient advice leaflet available in recovery and theatre 4

Background

Sugammadex (brand name Bridion®) is a modified cyclodextrin which is used to rapidly reverse the effects of the amino-steroidal non-depolarising neuromuscular blockers rocuronium and pancuronium.

Sugammadex and progesterone

As well as its effects on amino-steroidal NMBS, sugammadex binds to progesterone.

Progesterone is used widely in contraceptives. It is found in oral contraceptives (combined oral contraceptives and progesterone-only pill) as well as injectables (e.g. depo-provera), the implant (Nexplanon), the intra-uterine system and the vaginal ring.

A bolus dose of 4mg/kg sugammadex may reduce progesterone exposure by up to 34%, which is equivalent to missing one oral contraceptive dose. The effect of sugammadex on oestrogen is much lower and is therefore thought to be negligible.

All women of child-bearing age who receive a dose of sugammadex must be counseled about this interaction and how to avoid an unplanned pregnancy.

For women taking a progesterone-containing oral contraceptive, they should follow the “missed pill” advice in the package insert.

For women utilising a non-oral, hormonal form of contraception they should use barrier contraception for 7 days following administration of sugammadex.

To improve patient retention of this advice there is a patient information leaflet on sugammadex and contraception in theatre 4 anaesthetic room, recovery (in the sugammadex book) and to download here.

References

https://www.merck.com/product/usa/pi_circulars/b/bridion/bridion_pi.pdf

https://www.ema.europa.eu/en/documents/product-information/bridion-epar-product-information_en.pdf

BNF

YG CG 442 Patient Leaflet: Contraceptive Advice Following Sugammadex

During many surgeries it is necessary to administer medication to relax your muscles. Prior to waking you up it is important to reverse the effects of these medications with “reversal” drugs. You received one of these “reversal” drugs called sugammadex (Bridion®).
Sugammadex can interfere with hormone-based contraception, reducing the effectiveness and increasing the chance of an unplanned pregnancy.

Oral Contraception

If you use oral contraception you should follow the “missed pill” advice in the package insert. Keep taking your pill.

Hormone-based, non-oral contraception

If you use any hormone-based contraception other than a pill; such as an implant, injection, intra-uterine system (doesn’t include the copper coil) or vaginal ring; you should use barrier contraception (condom/diaphragm) for 7 days.

If you have any questions or concerns your GP or pharmacist can help. Show them this letter. They can find advice on sugammadex in the British National Formulary (BNF).

Key Points

- You received a drug called sugammadex which can interfere with hormone-based contraception
- For oral contraceptives follow the missed pill advice in the package insert
- For non-oral contraception use a barrier method (such as a condom/diaphragm) for 7 days

YG CG442 Taflen Cleifion: Cyngor Atal cenhedlu yn dilyn Sugammadex.

Yn ystod llawer o feddygfeydd mae angen rhoi meddyginiaeth i ymlacio'ch cyhyrau. Cyn eich deffro mae'n bwysig gwrthdroi effeithiau'r meddyginiaethau hyn â chyffuriau "gwrthdroi". Fe wnaethoch chi dderbyn un o'r cyffuriau “gwrthdroi” hyn o'r enw sugammadex (Bridion®). Gall Sugammadex ymyrryd ag atal cenhedlu ar sail hormonau, gan leihau effeithiolrwydd a chynyddu'r siawns o feichiogrwydd heb ei gynllunio.

Atal Cenhedlu Llafar

Os ydych chi'n defnyddio dulliau atal cenhedlu geneuol dylech ddilyn y cyngor “colli bilsen” yn y pecyn. Daliwch ati i gymryd eich bilsen.

Atal cenhedlu heb fod yn geg ar sail hormonau

Os ydych chi'n defnyddio unrhyw ddulliau atal cenhedlu sy'n seiliedig ar hormonau heblaw am bilsen; megis mewnblaniad, pigiad, system fewn-groth (nid yw'n cynnwys y coil copr) neu gylch y fagina; dylech ddefnyddio dulliau atal cenhedlu rhwystr (condom / diaffram) am 7 diwrnod.

Os oes gennych unrhyw gwestiynau neu bryderon gall eich meddyg teulu neu fferyllydd helpu. Dangoswch y llythyr hwn iddyn nhw. Gallant ddod o hyd i gyngor ar sugammadex yn y Fformiwlari Cenedlaethol Prydeinig (BNF).

Pwyntiau Allweddol

- Fe wnaethoch chi dderbyn cyffur o'r enw sugammadex a all ymyrryd ag atal cenhedlu sy'n seiliedig ar hormonau
- Ar gyfer dulliau atal cenhedlu geneuol dilynwch y cyngor bilsen a gollwyd yn y pecyn
- Ar gyfer atal cenhedlu nad yw'n geg, defnyddiwch ddull rhwystr (fel condom / diaffram) am 7 diwrnod