Ysbyty Gwynedd ICU/HDU Standard Drug Prescriptions and Infusion Strengths

*Ideally via a large/central vein
**Must be via a central vein
NS = normal saline
D5W = 5% dextrose
Version 1 (November 2022). Please provide feedback/suggestions via QR code:

DRUG NAME	MIXTURE	FINAL DILUTION	INFUSION RATE
Adrenaline**	5mg in 50mL of D5W	80microgram/mL	0-1.0microgram/kg/min - *Titrate to MAP target*
Alfentanil	25mg in 50mL of NS (ready diluted)	500microgram/mL	0-10mL/hour
Aminophylline	Loading: 5mg/kg in 250mL NS/G Maintenance: 0.5mg/kg/hr in 500mL NS/G	Loading: 5mg/kg in 250mL NS/G Maintenance: 0.5mg/kg/hr in 500mL NS/G	Loading: 30 minutes Maintenance: Over 24 hours Max rate = 25mg/min
Amiodarone*	Loading: 300mg in 100mL of D5W Maintenance: 600-900mg in 500mL of D5W	Loading: 300mg/100mL = 3mg/mL Maintenance 600-900mg/500mL = 1.2-1.8mg/mL	Loading over 20-120mins Maintenance over 24 hours = 21mL/h
Vasopressin**	20 units in 50mL of NS	0.4 units/mL	1.5-6mL/h - *Titrate to MAP target*
Atracurium	500mg in 50mL ready diluted	10mg/mL	270-1770microgram/kg/hour = 4.5-29.5microgram/kg/min *Titrate to TOF/peripheral nerve stimulator*
Calcium Chloride*	10mL of 10% pre-made solution	10mL of 10% pre-made solution	Slow IV injection (over 10mins)
Clonidine*	750microgram in 50mL of NS or D5W	15microgram/mL	Bolus can be given up to 75microgram 0.5-2microgram/kg/hour (can increase to 2-4microgram/kg/hour after Cons discussion)
Dobutamine**/*	**250mg in 50mL of NS or D5W *250mg in 250mL of NS or D5W	5mg/mL 1mg/mL	0-10microgram/kg/min
Fentanyl	2.5mg in 50mL of NS	50microgram/mL	0-5microgram/kg/hour
Furosemide*	50mg in 50mL NS	1mg/mL	0-6mL/hour (increase after Cons discussion)
GTN (nitroglycerin)*	50mg in 50mL ready diluted	1mg/mL	Start 1mL/hour - *Titrate to MAP target*
Haloperidol	0.5-5mg/dose	Max 18mg/24hours	IV bolus over 3-5 minutes (higher doses under Cons guidance)

Insulin/Dextrose for Hyperkalaemia*	10 units ActRapid in 50mL of 50% Dextrose	10 units ActRapid in 50mL of 50% Dextrose	Over 30 minutes
Ketamine	500mg in 50mL NS	10mg/mL	0.6-3mg/kg/hour
Labetalol**/*	**5mg/mL ready diluted ampoules *200mg in 200mL of D5W	**5mg/mL *1mg/mL	0-160mL/hour (=Max rate of 2mg/min) *Start 20ml/h + double every 30 minutes to MAP target*
Levosimendan**	12.5mg/5mL diluted to 250mL D5W	50microgram/mL	0.1microgram/kg/minute for ≤24 hours
Magnesium (replacement) **/*	20mmol in 100mL NS	0.2mmol/mL (1g = 4mmol)	Peripherally: over >4 hours Centrally: over >1-2 hours
Metaraminol*	20mg in 40mL NS or D5W	0.5mg/mL	0-10mL/hour (Bolus 1-2mL) - *Titrate to MAP target*
Midazolam	50mg in 50mL ready diluted or D5W	1mg/mL	0-10mL/hour
Milrinone**	10mg in 50mL of NS	200microgram/mL	Loading: 50microgram/kg over 10 mins Infusion: 0.375microgram/kg/min - 0.75microgram/kg/min (not to exceed 1.13mg/kg/day)
Morphine	50mg in 50mL of NS	1mg/mL	Titrate to effect (usually PCA)
Neostigmine	10mg in 50mL of NS	200microgram/mL	2mL/hour (400microgram/hour)
Nimodipine*	10mg in 50mL ready diluted	200microgram/mL	5mL/hour for first two hours, increased to 10mL/h
Noradrenaline**	Single Strength: 8mg in 50mL D5W Double Strength 16mg in 50mL D5W	SS: 160microgram/mL DS: 320microgram/mL	0-1.0microgram/kg/min - *Titrate to MAP target*
Potassium**	40mmol in 100mL of Water for Inj	0.4mmol/mL	Maximum rate 20mmol/hour**
Propofol	Either 1% or 2% available	1% =10mg/mL or 2% = 20mg/mL	0-4mg/kg/hour - *Titrate to RASS target*
Rocuronium	50mg/5mL ready diluted	10mg/mL	Bolus: 0.6-1.2mg/kg (IBW) Maintenance: 300-600microgram/kg/hour *Titrate to TOF/peripheral nerve stimulator*
Salbutamol*	Injection: 2.5mg in 50mL NS Infusion: 5mg in 500mL NS or D5W	Injection: 50microgram/mL Infusion: 10microgram/mL	Bolus: 250microgram over 3 minutes Infusion: 5-20microgram/min = 3-12mL/hour