# RECTUS SHEATH QUICK GUIDE

Ysbyty Gwynedd
Anaesthetics Department
Adran Anaestheteg

## CATHETER INSERTION

- **WHEN:** Before closure
- **WHERE:** Superior end of wound within peritoneal-muscle interface
- **INITIAL BOLUS:** 0.25% Chirocaine 15-20ml per catheter

OR

## ORDER PUMP

(08:30-15:00)

(Mon-Fri)

Prescribe on 'Once Only' section of drug chart:

**0.125% Levobupivicaine 5ml/hour via Elastomeric Pump per Catheter. Remove catheter(s) on day 3 post-op**

- Call **4137** (Aseptic Department) to request 2 Pajunk wound infusion elastomeric pumps & give patient details
- Send porter to pharmacy (Alaw ward corridor entrance) with drug chart
- Connect each pump to the filter via a Vygon Octopus 2 connector
- Set pump rate to **5ml/hr** with spanner

## SELF PREPARATION

IF NO PUMP AVAILABLE FROM PHARMACY

**You need:**

Pajunk Fuser-Pump set, 90x90 sterile drape, sterile gloves, filler needle, iv infusion set with 3-way tap, 250ml 0.9% saline, 80ml 0.5% Chirocaine

- Strict ANTT
- Connect pump to both catheters via Y-connector
- Set pump rate to **8ml/hr** with spanner

**Minimise Disconnection:**
wrap catheter/pump connection with Tegaderm

**Maximise Identification:**
Ensure bottle and line are clearly labelled

**Document:**
Complete acute pain referral form



