RECTUS SHEATH QUICK GUIDE

CATHETER INSERTION

- **WHEN**: before closure
- **WHERE:** superior end of wound within the peritoneum-muscle interface
- **INITIAL BOLUS:** 0.25% Chirocaine- 15-20ml per catheter, then 10ml via Y-connector

ORDER PUMP

OR

(08:30-15:00)
(Mon-Fri)

- Prescribe on 'Once Only' section of drug chart:
  - **0.125% Levobupivicaine 8ml/hour via Elastomeric Pump. Remove catheter on day 3 post op.**
- Call **4137** (Aseptic Department) to request a Baxter XLV wound infusion elastomeric pump & give patient details
- Send porter to pharmacy (Alaw ward corridor entrance) with drug chart

SELF PREPARATION

•IF NO PUMP AVAILABLE FROM PHARMACY

•**You need:**

•Pajunk Fuser-Pump set, 250ml 0.9% saline, 80ml 0.5% chirocaine, 90x90 drape, sterile gloves, filler needle, iv infusion set with 3-way tap.

•Strict aseptic technique

•Set pump rate to 8ml/hr with spanner

Minimise Disconnection:

- wrap catheter/pump connection with Tegaderm

Maximise Identification:

- Ensure bottle and line are clearly labelled

Document

- Complete acute pain referral form



