



Review article

Applications of biomaterials for immunosuppression in tissue repair and regeneration



Peng Shen^a, Yanxin Chen^a, Shuai Luo^a, Zhiyuan Fan^b, Jilong Wang^a, Jiang Chang^{a,c,**}, Junjie Deng^{a,*}

^a Engineering Research Center of Clinical Functional Materials and Diagnosis & Treatment Devices of Zhejiang Province, Wenzhou Institute, University of Chinese Academy of Sciences, Wenzhou, China

^b Department of Materials Science and Engineering, Drexel University, Philadelphia, PA 19104, USA

^c State Key Laboratory of High Performance Ceramics and Superfine Microstructure, Shanghai Institute of Ceramics, Chinese Academy of Sciences, Shanghai, China

ARTICLE INFO

Article history:

Received 28 October 2020

Revised 24 February 2021

Accepted 8 March 2021

Available online 17 March 2021

Keywords:

Biomaterial

Immunosuppression

Macrophage polarization

Regulatory T cell

Tissue repair and regeneration

ABSTRACT

The immune system plays an essential role in tissue repair and regeneration. Regardless of innate or adaptive immune responses, immunosuppressive strategies such as macrophage polarization and regulatory T (Treg) cell induction can be used to modulate the immune system to promote tissue repair and regeneration. Biomaterials can improve the production of anti-inflammatory macrophages and Treg cells by providing physiochemical cues or delivering therapeutics such as cytokines, small molecules, microRNA, growth factors, or stem cells in the damaged tissues. Herein, we present an overview of immunosuppressive modulation by biomaterials in tissue regeneration and highlight the mechanisms of macrophage polarization and Treg cell induction. Overall, we foresee that future biomaterials for regenerative strategies will entail more interactions between biomaterials and the immune cells, and more mechanisms of immunosuppression related to T cell subsets remain to be discovered and applied to develop novel biomaterials for tissue repair and regeneration.

Statement of Significance

Immunosuppression plays a key role in tissue repair and regeneration, and biomaterials can interact with the immune system through their biological properties and by providing physiochemical cues. Here, we summarize the studies on biomaterials that have been used for immunosuppression to facilitate tissue regeneration. In the first part of this review, we demonstrate the crucial role of macrophage polarization and induction of T regulatory (Treg) cells in immunosuppression. In the second part, distinct approaches used by biomaterials to induce immunosuppression are introduced, which show excellent performance in terms of promoting tissue regeneration.

© 2021 Acta Materialia Inc. Published by Elsevier Ltd. All rights reserved.

1. Introduction

The main function of the immune system is to defend the body against pathogens; it also plays key roles in tissue devel-

opment, homeostasis, and tissue regeneration [1,2]. The immune system can determine the outcome of a healing process, which normally ranges from complete regeneration that is identical to the undamaged tissue to incomplete regeneration such as fibrosis and scarring. The relationships between the immune system and tissue healing are complex because the immune responses can exert either beneficial or inhibitory effects on tissue repair and regeneration [3,4]. Normally, an inflammatory response is the initial response of the immune system to tissue injury [5,6]. However, long-time inflammation can inhibit the wound-healing process and delay the restoration of normal tissue architecture, thus making controlled inflammation necessary for improving the

* Corresponding author at: Wenzhou Institute, University of Chinese Academy of Sciences, Engineering Research Center of Clinical Functional Materials and Diagnosis & Treatment Devices of Zhejiang Province, No.1 Jinlian Road, Wenzhou, 325001, China.

** Corresponding author at: Shanghai Institute of Ceramics, Chinese Academy of Sciences, 1295 Dingxi Road, Shanghai, 200050, China.

E-mail addresses: jchang@mail.sic.ac.cn (J. Chang), j.deng@ucas.ac.cn (J. Deng).

healing process [2,7]. Therefore, immunosuppression strategies can be used to modulate components of the immune system to control undesired inflammatory responses and facilitate tissue repair and regeneration [8].

Immunosuppressive factors such as cytokines, microRNAs, small molecules, growth factors, and stem cells have been used as regenerative strategies [9–13]. These factors act through binding to receptors such as Toll-like receptors (TLR) and T cell receptors (TCR) [14] and then polarizing macrophages or inducing regulatory T (Treg) cell in the healing process [15,16]. Given that systemic approaches to applying immunosuppressive factors may result in severe off-target toxicity and low efficacy, biomaterials can serve as carriers for targeted delivery of immunosuppressive agents and control their release [1,17]. Moreover, biomaterials can also provide physical support to recruit immune cells and modulate them by tailoring their physiochemical properties for tissue remodeling [18].

This brief review focuses on the applications of biomaterials for immunosuppression in tissue repair and regeneration. We highlight immunosuppression-mediated mechanisms including macrophage polarization and Treg cell induction, which correlate with the innate and adaptive immune responses, respectively. We then propose future avenues in this field, including engineering biomaterials to target certain immune cell populations in a particular tissue or organ and the combination of typical biomaterials and delivery system platforms for seeking an immune-centric approach to enhance tissue regeneration.

2. The role of immune system in tissue repair and regeneration

2.1. Innate immune response and macrophage polarization

Innate and adaptive immune systems are two main parts of the human immune system. The innate immune system can be triggered by infections or injury. The activation process involves components of the immune system which includes the complement system and patrolling cells such as macrophages, neutrophils, and dendritic cells (DCs) [19–22]. Pattern recognition receptors (PRRs) expressed on patrolling cells can bind to danger signals such as damage-associated molecular patterns (DAMPs) and pathogen-associated molecular patterns (PAMPs). PAMPs are related to simple molecules or regular patterns of molecular structures shared by pathogens, while DAMPs are related to the intracellular molecules released by damaged extracellular matrix (ECM) and necrotic or stressed cells [23,24]. Both DAMPs and PAMPs contribute to inflammation and inflammatory diseases [25,26]. After danger signals are recognized by PRRs, inflammation is triggered by activating the transcription factors NF- κ B or interferon (IFN) regulatory factors [27]. Activated patrolling cells secrete more chemokines such as C-C motif ligand (CCL)-2 to recruit more neutrophils and circulating monocytes to the site, which release cytokines such as tumor necrosis factor- α (TNF- α), IL-1 β , and IL-6 to induce inflammatory responses [28]. Recruited circulating monocytes rapidly differentiate into inflammatory macrophages at the injury within three days [29]. Both neutrophils and inflammatory macrophages provide the first defense of innate immunity by phagocytizing invading pathogens, apoptotic cells, and cellular debris and efficiently destroying them with degradative enzymes [30,31]. In addition, innate immune cells such as neutrophils, macrophages, and DCs play key roles in tissue repair and regeneration. Neutrophils produce vascular endothelial growth factor (VEGF)- α and IL-17 to recruit more inflammatory cells, enhance angiogenesis, and activate the proliferation of epithelial cells, fibroblasts, and keratinocytes [32,33]. Activated DCs modulate the proliferation and differentiation of myoblasts in the skeletal muscle [34]. Furthermore, DCs can release extracellular vesicles to promote mesenchymal stem cell

(MSC) recruitment for bone repair and regeneration [35]. The effects of macrophages in tissue repair and regeneration will be introduced in detail in the following text.

Intensive investigation for several decades has revealed that apart from being scavenger cells, macrophages contribute to tissue repair and regeneration in much more complex roles [15,31]. At the damage site, resident and recruited macrophages proliferate and switch into distinct subtypes with different functions. They can be roughly classified into pro-inflammatory “M1” macrophages and anti-inflammatory “M2” macrophages. Macrophages are highly plastic, and their phenotype is determined by tissue microenvironments. M1 macrophages are induced by danger signals such as DAMPs, PAMPs, and cytokines such as TNF- α and IFN- γ [36], while M2 macrophages are generated by exposure to signals such as hormones, immune complexes, or cytokines such as IL-4, IL-10, IL-13, or IL-33. Accordingly, the functions of these two phenotypes are significantly different. M1 macrophages participate in phagocytosis and angiogenesis, and they secrete cytokines such as IL-1 β , IL-6, and TNF- α to recruit additional inflammatory cells [37]. The long-term presence of M1 macrophages triggers chronic inflammation that can exacerbate tissue injury and inhibit the healing process [38–40]. Mosser et al. [41] suggested a new grouping of macrophage populations based on three different homeostatic activities: host defense, wound healing, and immune regulation. Macrophages are classified into classically activated macrophages, wound-healing macrophages, and regulatory macrophages. Similar to M2 macrophages, wound-healing macrophages arise in response to IL-4, while regulatory macrophages are generated in response to various stimuli, including immune complexes, prostaglandins, G-protein-coupled receptor ligands, glucocorticoids, apoptotic cells, or IL-10 [42,43]. M2 macrophages can also be classified into four subtypes—M2a, M2b, M2c, and M2d [44]. These subtypes can lead to the suppression of inflammatory responses through different pathways [45]. For instance, M2a macrophages upregulate the expression of arginase, CD206, and MHC II molecules and secretion of IL-10 and TGF- β [46]. M2b macrophages produce both anti- and pro-inflammatory cytokines and promote Th2-type immune response [47]. M2c macrophages secrete high amounts of anti-inflammatory cytokines IL-10 and TGF- β and reduce M1 macrophage-initiated immune responses [48]. M2d macrophages are a novel M2 subset and referred to as tumor-associated macrophages (TAMs) that play a detrimental protumor role in promoting tumor metastasis and progression [49,50]. The profibrotic effects of M2 macrophages can stimulate ECM assembly and remodeling which is necessary for the wound healing process, but fibrosis can destroy normal tissue architecture and lead to fibrotic diseases [51–53]. Given that the inflammatory microenvironment of a damaged site hinders the wound healing process, immunosuppression is indispensable to switch macrophages from pro-inflammatory M1 to anti-inflammatory M2 phenotype, which may inhibit inflammation, enhance tissue repair, and drive tissue regeneration. However, because of the pro-tumoral and profibrotic properties of M2 macrophages, the processes of macrophage polarization need to be carefully controlled.

2.2. Adaptive immune response and Treg cell induction

The adaptive immune response is initiated by antigen presenting cells (APCs) such as macrophages, DCs, and B cells after the activation of the innate immune system [54]. After recognizing tissue damage, immature DCs residing in peripheral tissues take up antigens and migrate to draining lymph nodes as mature nonphagocytic DCs. Mature DCs process antigens into short peptides (signal 1) and present them on the major histocompatibility complex (MHC) molecules to form peptide-MHC complexes, which can be recognized by specific TCRs on T cells. Mature DCs also

display co-stimulatory ligands CD80/CD86 (signal 2) that can bind to receptor CD28 and synergize with signal 1 to initiate priming of naïve T cells [55–58]. Then, naïve T cells are activated and become effector T cells, which can be classified into CD4⁺ T cells including helper T (Th)1, Th2, Th17 cells, Treg cells, and cytotoxic CD8⁺ T cells. T cell subtypes play different roles in the tissue repair and regeneration [59–62]. Cytotoxic T cells are responsible for delaying fracture healing in bone regeneration. Reinke et al. [60] demonstrated that endogenous fracture regeneration was enhanced after the depletion of cytotoxic T cells, while the regenerative process was impaired by the transfer of cytotoxic T cells in a mouse model. It was suggested that TNF- α and INF- γ mainly produced by cytotoxic T cells inhibited osteogenic differentiation and the survival of MSCs, which was important for bone formation [63]. The effects of Th cells are distinct according to their subtypes. Th1 and Th17 cells produce cytokines and immune complexes [64], which promote the generation of M1 macrophages and result in more pro-inflammatory cytokines and further tissue damage. On the other hand, Th2 cells secrete cytokines such as IL-4, IL-5, and IL-13 that regulate the activities of M2 macrophages for suppressing Th1 and Th17-driven inflammation [65,66]. The contribution of Th2 cells to tissue repair and regeneration depends on M2 macrophages, and excessive activation of Th2 cells may result in allergy and pathological fibrosis [31].

Treg cells are a crucial part of regulatory mechanisms in the immune system, and they maintain immune homeostasis by suppressing autoreactive and hyperactive immune responses [67]. Because of expressing the transcription factor forkhead box protein P3 (Foxp3), Treg cells are often identified as CD4⁺Foxp3⁺ T cells. Research studies have proved that Treg cells are crucial for orchestrating tissue repair and regeneration in various tissues such as skin [68], bone [69], lung [70], and cardiac muscle [71]. Treg cells generally reside in lymphoid organs, but they can be recruited to injured sites, where they facilitate inflammation resolution by multiple mechanisms of suppression to directly and indirectly regulate tissue healing [72,73]. In the indirect way, Treg cells can produce arginase and anti-inflammatory cytokines such as IL-10 and TGF- β to suppress conventional T cells [74–76]. The anti-inflammatory microenvironment provided by the secreted factors can drive the local immune response from inflammation to tissue healing process such as the modulation of neutrophils [77], regulation of Th2 cells [78,79], and polarization of M2 macrophage [80,81]. Moreover, Treg cells can directly facilitate tissue regeneration by activating tissue-specific stem cells and progenitor cells [82–84]. Treg cells in skin preferentially localize to hair follicles, and the high expression of Jagged 1 by Treg cells is closely related to the proliferation and differentiation of hair follicle stem cells [68]. Similarly, *in vitro* induced Treg cells can effectively enhance the expansion of muscle satellite cells and inhibit their myogenic differentiation [82]. Given the important role of Treg cells in tissue regeneration, the induction of Treg cells and macrophage polarization can be used for immunosuppression to facilitate tissue repair and regeneration.

3. Biomaterials for immunosuppression in tissue repair and regeneration

The immune system participates in tissue repair and regeneration through both innate and adaptive immune responses. Therefore, it is possible to regulate immune components for tissue repair and regeneration through immunosuppression such as M2 macrophage polarization and Treg cell induction. Biomaterials have provided an ideal platform for immunosuppression in tissue healing processes. They can improve biocompatibility and alleviate foreign body reaction through optimizing physiochemical properties and delivering immunosuppressive factors [85–88]. An extensive

array of biomaterials has been investigated for immunosuppression, including ceramics, decellularized ECM, collagen, elastin, alginate and hyaluronic acid (HA). Regardless of the types of biomaterial, delicate design and engineering are required for their applications in immunosuppression and tissue regeneration [89–91]. As shown in Fig. 1, the strategies of immunosuppression modulation through biomaterials in regenerative medicine can be roughly categorized into biomaterial-based, stem cell-based, and molecular signaling-based strategies [92]. Scaffolds can provide physical support to transplanted cells or host cells in the biomaterial-based strategy [93]; stem cells can rapidly proliferate and differentiate to compensate the lost tissue cells in the cell-based strategy [94]; growth factors such as insulin-like growth factor (IGF)-1 and VEGF can promote angiogenesis and tissue remodeling in the molecular signaling-based strategy [95,96]. Additionally, these three strategies can also be combined through scaffolds to co-deliver transplanted stem cells and growth factors [97,98]. Furthermore, the therapeutic effects can be further improved if biomaterials can modulate immune components and suppress undesired immune responses. The immunosuppression provided by biomaterials has significant effects on tissue repair and regeneration, and such applications of biomaterials are discussed in the following part.

3.1. Biomaterials for macrophage polarization

The pro-inflammatory function of M1 macrophages is required for triggering the healing process in a damaged site, but the resolution of inflammation is important for the healing process because prolonged inflammation can exacerbate tissue damage and inhibit tissue repair [99–102]. Furthermore, anti-inflammatory M2 phenotype has been shown to indicate constructive tissue remodeling [103,104]. Therefore, it is a promising strategy to polarize macrophage phenotype to improve tissue repair and regeneration.

3.1.1. Polarization to M2 macrophages via physiochemical biomaterials properties

Macrophages are present in a complex microenvironment with many physical and chemical signals that may influence their phenotype polarization. In the past few decades, it has been proved that ECM stiffness, architecture, and composition can regulate macrophage phenotypes [105–107]. For instance, Wu et al. [105] reported a family of thermoresponsive nanohybrid elastomer scaffolds with different stiffness softening effects. They found that stiffer substrates led to severe foreign body reaction, while softer substrates promoted M2 macrophage polarization. Yang et al. [106] fabricated hydroxyapatite (HA) bioceramics with different microscale circular patterns and nanotopographies. They found that the nanoneedle-structured HA surface could induce the M2 polarization of macrophages. Furthermore, the micro/nano hierarchical structures with the micropattern sizes of 36 μm could more significantly enhance M2 polarization of macrophages than HA bioceramics with sizes of 4 μm and 12 μm . Pinto et al. [107] compared the effect of normal and tumor ECM on macrophage polarization and found that macrophages in tumor matrices differentiated toward an anti-inflammatory M2-like phenotype. When biomaterials are implanted in the damaged tissues, macrophages are among the first immune cells to contact them. Thus, physiochemical cues of implanted biomaterials can directly change the tissue microenvironment for modulating phenotypes and functions of macrophages. After implantation of biomaterials, the initial presence of M1 macrophages triggers a necessary inflammation process, but prolonged inflammation may result in chronic inflammatory events and inhibit tissue regeneration. Therefore, appropriate physiochemical cues of biomaterials can be employed to drive macrophage polarization from M1 to M2 phenotype in regenerative strategies [108–110]. Therefore, appropriate physiochemical cues of




Fig. 1. A schematic illustration of biomaterials for immunosuppression through macrophage polarization and induction of Treg cells in tissue regeneration. Physiochemical cues of biomaterials, such as topography, pore size, particle size, surface charge, and hydrophobicity, can polarize macrophages from M1 to M2 phenotype. Biological cues such as anti-inflammatory molecules, MSCs, and microRNA can participate in macrophage polarization and induction of Treg cells. The process of macrophage polarization also affects the induction of Treg cells.

biomaterials can be used to induce macrophage polarization in regenerative strategies.

Physical properties such as topography, pore size, substrate stiffness, and particle size have been implicated in modulating the macrophage responses [111]. Different sizes of the microstructure on a surface can exhibit various performance on macrophages. For instance, Mcwhorter et al. [108] showed that mouse bone marrow-derived macrophages (BMDMs) could be elongated and differentiated toward M2 phenotype by 20 μm wide lines on a polydimethylsiloxane (PDMS) surface *in vitro*. However, if the width of lines was changed to 50 μm , BMDMs showed no difference from the control group without patterns. Moreover, Chen et al. [109] designed nanoporous anodic alumina with different pore sizes to examine modulatory effects of the nanopore structure and pore size on macrophage responses. They found that 50 nm nanoporous structure could effectively polarize macrophages to M2 phenotype, leading to anti-inflammatory reactions. Micro- and nano-scale fibers that are widely applied in tissue engineering can modulate macrophages through fiber arrangement or diameters. Jia et al. [110] compared the effects of aligned and random poly(L-lactic acid-co- ϵ -caprolactone) nanofibers on the development of macrophage phenotypes (Fig. 2A and B). Aligned nanofibers significantly induced macrophage elongation along the nanofibers and their polarization to M2 macrophages, while random nanofibers induced M1 macrophages (Fig. 2C and D). Interestingly, aligned nanofibers drastically promoted Schwann cell infiltration and axonal regrowth than random nanofibers *in vivo* (Fig. 2E and F). Furthermore, Garg et al. [112] found that mouse BMDMs were polarized from M1 to M2 phenotype when polydioxanone (PDO) fiber diameter increased from 0.35 μm to 2.8 μm and the intra pore size increased from 1 μm to 15 μm . Moreover, substrate stiffness can also affect macrophage polarization. Sridharan et al.

[113] demonstrated that stiff polyacrylamide gels (323 kPa) primed macrophages toward M1 phenotype, while softer gels (11 and 88 kPa) induced M2 macrophages. In addition, surface roughness is also an important parameter of biomaterials. Rough surfaces in bone facing implants can improve the performance of osteoblasts and MSCs [114]. Compared to the flat surface, a rough surface is preferred by macrophages to adhere, which can induce secretion of inflammatory cytokines [115].

Surface charge, hydrophilicity, and hydrophobicity also play key roles in regulating macrophage phenotypes and functions [116,117]. For example, Fush et al. [116] studied the effects of surface charge on human macrophages by using carboxyl-(COOH) and amino-(NH₂) functionalized polystyrene nanoparticles (PSNPs). M2 polarization was strongly inhibited by both cationic and anionic PSNPs, but the expression of M1 macrophage markers was not significantly affected. Moreover, hydrophilicity and hydrophobicity also play an important role in macrophage polarization. Rostam et al. [117] cultured human monocytes for 6 days on untreated hydrophobic polystyrene and O₂ plasma-etched hydrophilic polystyrene surfaces. Their data showed that monocytes on the hydrophilic surface were switched to M1 phenotype with enhanced expression of pro-inflammatory transcription factors STAT1 and IRF5. In contrast, monocytes on the hydrophobic surface were polarized to M2 macrophages with high secretion of anti-inflammatory cytokines IL-10 and CCL18. Similarly, McBane et al. [118] demonstrated that a biomaterial surface coated with degradable polar hydrophobic ionic polyurethane (D-PHI) could drive human monocytes to become M2 macrophages. This was proved by the decreasing amount of pro-inflammatory cytokines TNF- α , IL-1 β , and HMGB1 and the increasing amount of anti-inflammatory cytokine IL-10, which was not observed in control groups coated with tissue culture polystyrene.




Fig. 2. Elongated macrophages are shown on aligned nanofibers (A) compared with random nanofibers (B) under SEM imaging. (C–D) Quantification of TNF- α (mainly from M1 macrophages) and IL-10 (mainly from M2 macrophages) secreted by macrophages cultured on different nanofiber scaffolds or culture plates by ELISA assay. (E–F) Statistical analysis of S-100 (Schwann cells) positive areas and axon numbers between the random and aligned nerve guidance conduit groups for assessing nerve regeneration *in vivo*. Error bars represent standard deviation, * indicates $p < 0.05$ and *** indicates $p < 0.001$ [110].

Biomaterials that use ECM components or mimic ECM structures can create a microenvironment that benefits immunomodulation and tissue repair. Biomaterials composed of decellularized ECM have shown notable success in eliciting an anti-inflammatory (M2-like) host response and promoting constructive and functional tissue remodeling [119,120]. Decellularized ECM scaffolds derived from xenogeneic tissues are well tolerated by the host without showing hyperacute or delayed rejection [121]. Furthermore, the degradation of ECM scaffolds can release bioactive factors such as chemoattractant [122], growth factors [123], and extracellular vesicles [124] to promote M2 macrophage polarization and recruit endogenous stem cells for tissue regeneration. Sicari et al. [125] found that the degradation products of ECM such as solubilized small intestinal submucosa promoted M2 macrophage phenotype and enhanced the migration and myogenesis of skeletal muscle progenitor cells. In addition, the functions of ECM can be affected by multiple factors, including the use of chemical cross-linking [104], the source tissue of ECM [126] and the supplementary use of anti-inflammatory drugs [127]. Recently, Fan et al.

[128] reported a red blood cell membrane-derived macroporous hydrogel that mimics ECM structure, which exhibited high performance on M2-like phenotype polarization (Fig. 3A). The microporous scaffolds were formed through chemical crosslinking of vesicles derived from red blood cell membranes (RBCMs) (Fig. 3B). In the *in vivo* experiment, RBCM scaffolds showed significantly higher infiltration of macrophages than DCs and neutrophils (Fig. 3C–E). High percentages of these macrophages had M2 phenotype as confirmed by gene expression and surface markers (Fig. 3F–G). As a major constituent of human ECM, HA is also a compelling biomaterial. The effects of HA on macrophage polarization depend on the molecular weight of the polymer chains [129]. High-molecular-weight HA is indispensable for maintaining homeostasis by facilitating anti-inflammatory responses [130]. In contrast, low-molecular-weight HA induces M1 macrophages as proved by increased expression of pro-inflammatory genes NO and TNF- α [129]. In addition, molecular weights also influence the effects of keratin biomaterials on augmenting M2 phenotype *in vitro* [131]. Both high-molecular-weight keratin (KOS^H) and keratin peptide




Fig. 3. (A) A schematic illustration of macroporous red blood cell membrane (RBCM)-derived scaffold. (B) RBCM scaffolds under SEM imaging. Scale bar = 100 μ m. (C–E) Percentages of dendritic cells, neutrophils, and macrophages in RBCM and MA-alginate scaffolds after scaffold injection *in vivo*. (F–G) qPCR analysis of Arg-1 (M2 macrophage marker) and iNOS (M1 macrophage marker) expression in cells from RBCM and MA-alginate scaffolds and M0, M1, and M2 BMDMs. Error bars represent standard deviation, and * indicates $p < 0.05$ [128].

(KOS^P) coatings show potentials to drive macrophage polarization to anti-inflammation, but KOS^P exhibits more effectiveness than KOS^H.

Bioceramics can release bioactive ions to modulate macrophage phenotypes and regulate stem cells to enhance tissue regeneration [132–134]. For example, Huang et al. [3] demonstrated that silicate bioceramics—akermanite (AKT) and nagelschmidtite (NAGEL) could release silicon (Si), magnesium (Mg), and calcium (Ca) ions to inhibit inflammatory cytokine secretion by macrophages as compared to tricalcium phosphate (TCP) bioceramics *in vitro* and *in vivo*. Furthermore, Zhou et al. [135] showed that mesoporous bioactive glass incorporated with ionic Cu²⁺ could induce VEGF secretion of BMSCs and triggered the production of regenerative M2 macrophages, leading to enhanced angiogenesis and immune responses. Dong et al. [136] found that bioactive glass ion extracts could polarize M2 macrophages and enhance paracrine effects between macrophages and repairing cells. In addition, Shi et al. [137] reported that europium-doped mesoporous silica nanospheres could stimulate pro-inflammatory responses of macrophages to activate osteogenesis. Similarly, copper-doped mesoporous silica nanospheres could also initiate proper pro-inflammatory responses for inducing osteogenesis [138].

The combination of physiochemical properties can synergize to direct the differentiation of macrophages to the pro-healing M2 phenotype. For example, Chen et al. [139] adjusted the surface chemistry and scale of nanotopography simultaneously to modulate the osteoimmune environment. The biomaterial with carboxyl acid-tailored and 68 nm nanoparticle-coated surface showed promising performance in terms of increasing M2 macrophages and triggering the osteogenic differentiation of MSCs. Bachhuka et al. [140] fabricated nanoporous surfaces with controlled pore size and different surface charges. The data showed that the biomaterial with nanopores of 200 nm and carboxyl functionality significantly downregulated pro-inflammatory cytokines such as IL-6, IL-1 β , and TNF- α and upregulated anti-inflammatory cytokines such as arginase, IL-1 receptor antagonist, and IL-10.

3.1.2. Polarization to M2 macrophages through the delivery of biological cues

Anti-inflammatory cytokines such as IL-4 and IL-10 have been frequently studied for regenerative strategies in recent years because of their capability to polarize M2 macrophages [46,103,141]. IL-10 can cause a complete regeneration in fetal mice models, and IL-10-deficient fetal mice show scar formation [10,142,143]. As a




Fig. 4. (A) A schematic illustration of NHG-MS modified by heparin and loaded with IL-4. (B–C) NHG-MS with the highly porous nanofibrous architecture under SEM measurement. (D–G) Immunohistochemical assessment of macrophage phenotypes through quantitative analysis of iNOS (M1 macrophage marker), CD68 (pan-macrophage marker), and CD206 (M2 macrophage marker). (D) CD68 showed significantly higher expression in the three DM groups than in the normal control group. (E–G) Fewer M1 macrophages and more M2 macrophages observed in the control and IL4-loaded NHG-MS DM groups. Error bars represent standard deviation, and ** indicates $p < 0.01$ [145].

regenerative strategy, IL-4 is usually delivered to the injured site as proteins [144]. For instance, Hu et al. [145] developed nanofibrous heparin-modified gelatin microspheres (NHG-MS) for delivering IL-4 to provide a healing microenvironment in diabetes mellitus (DM). IL-4 was incorporated in the microsphere and could bind to heparin for stabilization (Fig. 4A). The microspheres effectively increased the number of M2 macrophages, which enhanced the bone repair and regeneration process under the DM condition (Fig. 4D–G). Bonito et al. [146] designed heparin/IL-4-functionalized supramolecular elastomers to modulate macrophage phenotypes and protein secretion. The supramolecular approach was successfully translated to a three-dimensional (3D) electrospun scaffold to prolong the bioactivity of IL-4 under conditions mimicking hemodynamic microenvironments.

Compared to IL-4 delivered as proteins, IL-10 is mostly delivered to the target through plasmid DNA or virus vectors. For example, Bridges et al. [147] designed a PLG scaffold with IL-10 viral vectors for localized and sustained IL-10 expression to resolve inflammation. Leukocytes such as macrophages, DCs, and neutrophils were modulated by the scaffold in their numbers and relative proportions. The scaffold showed high performance in terms of increasing IL-10 secretion by macrophages and decreasing IFN- γ expression in DCs, macrophages, and CD4⁺ T cells. Holladay et al. [148] designed an IL-10 plasmid-loaded collagen scaffold for enhancing stem cell transplantation in myocardial infarction. Increased stem cell retention and the recovery of cardiac functions in animal models positively correlated with the induction of IL-10. Compared to the scaffold control, the scaffold with IL-10 plasmids modulated the immune response by downregulating CD80⁺ macrophages (M1) while upregulating CD163⁺ macrophages (M2). Furthermore, Jain et al. [149] fabricated IL-10 plasmid-loaded alginate nanoparticles for treating rheumatoid arthritis (RA). The IL-10 plasmid was encapsulated into alginate nanoparticles, and tuftsin peptide was modified on the surface of nanocarrier for targeting macrophages. This method significantly decreased the expression

of systemic and joint tissue pro-inflammatory cytokines such as IL-1 β , IL-6, and TNF- α and attenuated the inflammation and joint destruction.

Biomaterials can also sequentially release different immunomodulators to stimulate M1 macrophages at first and rapidly polarize them to M2 macrophages, which can maximize their regenerative activity and minimize their side effects [150,151]. In a natural tissue healing process, the injured site witnesses the sequential activities of pro-inflammation and anti-inflammation, which are respectively dominated by M1 and M2 macrophages [152–154]. Neovascularization is closely related to macrophage polarization, and it has been proved that M1 phenotype can initiate angiogenesis and M2 phenotype can enhance vessel maturation [155]. Thus, it is possible to induce an effective regenerative microenvironment by modulating numerous immune signals involved in macrophage polarization simultaneously or sequentially. Chen et al. [156] designed a titania nanotube system with double hydrogel layers to modulate the release of IFN- γ and IL-4. In this system, pro-inflammatory cytokine IFN- γ could be totally released in 3 days for triggering M1 macrophages. The release of anti-inflammatory cytokine IL-4 was sustained even after 4 days; thus, it could switch macrophages to the M2 phenotype. The results proved that macrophage phenotype switch correlated with the sequential release of these two cytokines and that vascularization in the site was enhanced more than that achieved using IFN- γ or IL-4 alone. Similarly, Spiller et al. [144] developed decellularized bone scaffolds for sequentially releasing IFN- γ and IL-4. In this scaffold, IFN- γ was physically adsorbed on the scaffold so that it could be quickly released. IL-4 was attached through biotin-streptavidin binding, and its release lasted a few days longer.

MSCs are widely used in cell-based tissue engineering due to their powerful regenerative potential [157,158]. Interactions between MSCs and macrophages can determine the initiation of tissue regenerative processes and the formation of scar tissue [159].




Fig. 5. (A) A schematic illustration of the process for developing nanoparticle/miR-223*-laden GelMA hydrogels and the use of these adhesive hydrogels for wound healing. (B) A quantitative evaluation of wound areas after treating wounds with DPBS, GelMA, Neg miRNA, or miR-223* for 12 days. (C–F) miRNA transfection and macrophage polarization studies in J774A.1 macrophages *in vitro*. M1 represents cells that received LPS+IFN- γ treatment for 16 h before incubation with either groups. Neg miRNA represents negative control miRNA and miR-223* represents miR-223 5p mimic. (C–E) Decreased expression of proinflammatory cytokines (TNF- α , IL-6, and IL-1 β mRNA) was observed upon 24 and 48 h post-incubation with GelMA/NP/Neg miRNA and GelMA/NP/miR-223* hydrogels. (F) A significantly higher ratio of Arg-1 (M2 macrophage marker)/iNOS-2 (M1 macrophage marker) gene expression was observed in the M1/GelMA/NP/miR-223* group. Error bars represent standard deviation; * indicates $p < 0.05$ and ** indicates $p < 0.01$ [169].

Macrophages are necessary for recruiting and regulating the differentiation of MSCs during tissue regeneration [160,161]. In return, MSCs also regulate the functions of macrophages. The beneficial effect of MSCs can potentially be enhanced by infiltrating more macrophages in the defect site. Seebach et al. [162] reported that fibrin implants embedded with MSCs could rapidly attract M1 macrophages and endothelial progenitor cells to improve vascularization and bone regeneration. MSCs can also be used for macrophage polarization. In a co-culture model, MSCs dramatically enhanced the expression of anti-inflammatory cytokine IL-10 and suppressed the production of pro-inflammatory cytokines such as TNF- α , IL-1 β , and IL-6 by macrophages [163,164]. Furthermore, Qiu et al. [165] showed that MSCs and decellularized ECM could synergistically drive macrophages toward the M2 phenotype to promote skeletal muscle tissue regeneration.

Apart from anti-inflammatory cytokines and MSCs, many other factors could be used to polarize M2 macrophages, such as pro-resolution mediators, microRNAs (miRNAs), and immunosuppressive drugs. Pro-resolution mediators such as resolvins and lipoxins can shift the polarization balance toward M2 macrophages

[166,167]. For instance, Shi et al. [168] developed a polycaprolactone vascular graft loaded with Aspirin-Triggered Resolvin D1 (AT-RvD1), which significantly promoted macrophage polarization into M2 macrophages *in vitro* and *in vivo* and enhanced smooth muscle regeneration in rats. miRNAs can control the translation and degradation of messenger RNAs; thus, the modulation of miRNAs is essential to regulate macrophage polarization. Although miRNA-based regenerative strategies are rare, studies have shown that miRNAs such as miR-124, miR-125a-3p, miR-223, let-7c, miR-132, and miR-146a could switch M1 to M2 macrophages and contribute to tissue healing [169–172]. Saleh et al. [169] developed adhesive gelatin methacryloyl (GelMA)-based hydrogels by incorporating miR-223 5p mimic (miR-223*)-laden HA nanoparticles (Fig. 5A). The adhesive hydrogels could efficiently reduce the expression of proinflammatory cytokines and induce macrophage polarization to M2 phenotype *in vitro* (Fig. 5C–F). In the *in vivo* experiment, the percentage of wound closure for the miR-223* treatment group was significantly higher than that for Neg miRNA-, GelMA-, or phosphate-buffered saline (DPBS)-treated wounds (Fig. 5B). The adhesive hydrogels triggered the resolution of the inflammation




Fig. 6. (A) A schematic illustration of multifunctionalized PLLA NF-SMS with IL-2/TGF- β -loaded MSN and miR-10a-loaded PLGA MS. (B) The highest gene expression levels of anti-inflammatory cytokine IL-10 and Foxp3 in total T cells were observed in the miR-10a/IL-2/TGF- β group. (C) The bone volume changes in a mouse periodontal disease model showed that PLLA NF-SMS rescued bone resorption. Error bars represent standard deviation, and *** indicates $p < 0.001$ [11].

and enhanced the formation of new and vascularized skin tissue to accelerate the tissue regeneration rate. The immunosuppressive drug glucocorticoid is highly potent in anti-inflammation. Dexamethasone (Dex) is a type of glucocorticoids and has been shown to shape macrophage phenotypes for tissue repair and regeneration [173]. For example, Jiang et al. [12] developed a PDMS-based 3D microporous scaffold platform for locally delivering the immunosuppressive drug Dex in a diabetic mouse model. The studies showed a dose-dependent impact of Dex on M2 macrophage polarization. The elevated M2 macrophage levels suppressed the inflammatory pathways and maintained the bioactivity of islets. Given the similar effects of immunosuppressive drugs on anti-inflammatory cytokines, it is reasonable to co-deliver Dex and IL-4 to promote macrophage switch from M1 to M2 phenotype. Kumar et al. [174] designed an injectable silk hydrogel loaded with IL-4 and Dex for treating type 1 diabetes (T1D). The mild gelation of the hydrogel provided a protective 3D support for transplanted islets without compromising their functions. The hydrogel showed high performance in terms of islet viability, insulin secretion, and endothelial cell maintenance.

3.2. Biomaterials for Treg cell induction

Treg cells play a pivotal role in the healing process, and Treg cell induction can be an effective approach for tissue repair and regeneration [59,175]. Treg cells are in the downstream of immune responses; thus, their development and maintenance are regulated by macrophages, tolerogenic dendritic cells, and other immune cells [176–178]. For instance, Li et al. [179] demonstrated that RelB gene-silenced DCs could selectively promote Treg generation. Treg cells can also be induced by the delivery of cytokines, growth factors, and MSCs through biomaterials [69,154,180]. The induced Treg cells can also contribute to the resolution of inflammation and switch macrophages from M1 to M2 phenotype [16,181].

Both IL-2 and TGF- β play an important role in the induction and regulation of Treg cells. TGF- β needs IL-2 to enhance Foxp3 transcription factor expression in CD4⁺CD25⁺ Treg cells [182,183]. In contrast, IL-2- and IL-2R-deficient mice develop extensive lymphadenopathy and systemic autoimmunity [184]. Thus, the co-delivery of IL-2 and TGF- β can induce Treg cells. For example, McHugh et al. [180] developed biodegradable PLGA nanoparticles loaded with IL-2 and TGF- β to induce Treg cells *in vitro* and expanded their population *in vivo*. Induced Treg cells were stable; thus, they could exert longer immunosuppressive effects in the inflammatory microenvironment. To manipulate Treg cells *in situ*, Liu et al. [11] designed a combined delivery vehicle including poly(L-lactic acid) (PLLA) nanofibrous spongy microspheres (NF-SMS), PLLA/polyethylene glycol (PEG) co-functionalized mesoporous silica nanoparticles (MSN), and poly(lactic acid-co-glycolic acid) microspheres (PLGA MS) (Fig. 6A). In this system, IL-2/TGF- β -loaded MSN and miR-10a-loaded PLGA MS locally recruited T cells and induced their differentiation into Treg cells. PLLA NF-SMS as a scaffold helped Treg cells to adhere and proliferate (Fig. 6B). In addition, the multifunctionalized NF-SMS system enriched Treg cells *in vivo*, which suppressed the exacerbation of host immune response in mouse periodontal disease models and rescued periodontal bone loss (Fig. 6C).

Cytokine IL-33 acts as an endogenous danger signal, which attracts Treg cells to accumulate in an injured tissue [185,186]. Although the role of IL-33 in tissue regeneration has not yet been elucidated, IL-33 can induce the proliferation of Treg cells in different tissues within a few days [187–189]. For example, Liu et al. [154] developed a PLGA scaffold for the local delivery of IL-33 to facilitate allogeneic islet transplantation. IL-33 exhibited beneficial immunosuppressive effects in the animal model by upregulating Treg cells and reducing cytotoxic CD8⁺ T cells, resulting in a prolonged graft survival and delayed islet engraftment. The rationale was that IL-33 induced the proliferation of IL-33R (ST2)-expressing Treg cells and these cells produced the

epidermal growth factor-like molecule amphiregulin to mediate tissue repair [190–192]. Furthermore, IL-33 is a promising pathway for skeletal muscle regeneration as a recent study reported that supplementation with IL-33 enhanced Treg cells for aging skeletal muscle regeneration in a mouse model [193].

The MSC-based strategy is one of the essential regenerative strategies. MSCs can produce soluble factors to inhibit T cell responses and induce Treg cells [194]. For instance, Cosenza et al. [195] reported that microparticles and exosomes derived from MSCs could inhibit T cell proliferation and reduce the percentages of CD4⁺ and CD8⁺ T cell subsets. In contrast, the number and percentage of Treg cell populations were elevated. Liu et al. [69] developed a hydroxyapatite/tricalcium phosphate (TCP-HA) bone scaffold with bone marrow MSCs to enhance bone regeneration. Bone marrow MSCs could induce T cell apoptosis, which resulted in the upregulation of Treg cells. The scaffold significantly improved the number of Treg cells and reduced IFN- γ and TNF- α levels in the implanted site.

In addition, *ex vivo* cultured Treg cells have been delivered directly into the injured site. For example, Roballo et al. [8] developed a degradable poly(ethylene glycol) norbornene (PEGNB) hydrogel for delivering Treg cells to prolong the survival of peripheral nerve allografts. The hydrogel sustained the release of Treg cells over two weeks. The released Treg cells effectively infiltrated the graft and blocked the host immune response. The performance of tissue regeneration in the recipient rats was comparable to that of allograft control.

4. Conclusion and future perspectives

Many current advances in biomaterials have been used for immunosuppression through macrophage polarization or Treg cell induction. The outcomes demonstrated their high performance in facilitating tissue repair and regeneration. The immunomodulators and immune cells mentioned above are only a fraction of the complex immune system. In light of the deep interactions between the immune system and the tissue healing process, it is promising to use immunosuppression by biomaterials to treat tissue damages; this area, however, requires further investigations on the mechanisms of immune responses and tissue regeneration.

To enhance the regenerative effects of biomaterials for immunosuppression, several challenges remain to be solved. First, the detailed understanding of biomaterial-immune cell interactions is lacking; thus, it is challenging to develop effective immunomodulatory biomaterials. Second, macrophages have been frequently studied during the past decades, and they play a crucial role in the tissue healing process. However, it is indicated that the induction routes and regulated processes of macrophage polarization are complex network systems rather than simple pathways. In addition, M2 macrophages have four subtype, namely M2a, M2b, M2c, and M2d, which are activated by different cytokines and can express different immunomodulatory signals. These different mechanisms can directly affect the regenerative effects of biomaterials. Third, the regenerative strategies provided by biomaterials need to follow the natural healing process. The aim of macrophage polarization is to control immune responses to enhance tissue regeneration rather than to simply polarize M1 to M2 phenotype. Although large numbers of M1 macrophages can delay tissue repair, they are necessary to induce angiogenesis. In contrast, M2 macrophages are necessary for inflammation resolution and tissue remodeling, but excessive levels of M2 macrophages may lead to pathological fibrosis. Therefore, sequential delivery of cytokines is a feasible approach for the temporospatial control of macrophage polarization to enhance tissue regeneration, which can be easily realized by delicate designs of biomaterials. Finally, compared to the large amount of studies on macrophages, research studies on T cell sub-

sets for tissue regeneration are limited. Evidence has shown that T cell subsets such as CD8⁺ T and Th1 cells have anti-regenerative properties and other cell subsets such as Th2, $\gamma\delta$ T, and Treg cells have pro-regenerative properties. Thus, biomaterials can be used to modulate T cell activities to enhance regeneration. However, current applications of T cell switch and inhibition of cytotoxic T cells for the healing process are scarce, which may be due to unclear mechanisms of T cells. Even for the induction of Treg cells, it is mainly used to induce immune tolerance for treating autoimmune disorders, where the effects of Treg cells are often unexpectedly altered by conventional T cells.

Presently, because of our limited knowledge of the immune system, the regenerative effects of biomaterials for immunosuppression are restricted. The evolution of biomaterials can provide an opportunity to sprint, and this opportunity can be facilitated by new discoveries in immunology.

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (31971260), Zhejiang Provincial Natural Science Funds for Distinguished Young Scholar (LR20C100001), Zhejiang Provincial Natural Science Foundation of China (LY19C100002), Key Laboratory of Orthopaedics of Zhejiang Province (grant no: ZJGK1804Y), Wenzhou Scientific Research Project (Y20190136), and a seed grant from the Wenzhou Institute, University of Chinese Academy of Sciences (WIUCASQD2019004).

References

- [1] M.O. Dellacherie, B.R. Seo, D.J. Mooney, Macroscale biomaterials strategies for local immunomodulation, *Nat. Rev. Mater.* 4 (6) (2019) 379–397.
- [2] Z. Julier, A.J. Park, P.S. Briquez, M.M. Martino, Promoting tissue regeneration by modulating the immune system, *Acta Biomater.* 53 (2017) 13–28.
- [3] Y. Huang, C. Wu, X. Zhang, J. Chang, K. Dai, Regulation of immune response by bioactive ions released from silicate bioceramics for bone regeneration, *Acta Biomater.* 66 (2018) 81–92.
- [4] L. Chung, D.R. Maestas, Jr., F. Housseau, J.H. Elisseeff, Key players in the immune response to biomaterial scaffolds for regenerative medicine, *Adv. Drug Deliv. Rev.* 114 (2017) 184–192.
- [5] C.A. Loynes, J.A. Lee, A.L. Robertson, M.J. Steel, F. Ellett, Y. Feng, B.D. Levy, M.K. Whyte, S.A. Renshaw, PGE2 production at sites of tissue injury promotes an anti-inflammatory neutrophil phenotype and determines the outcome of inflammation resolution in vivo, *Sci. Adv.* 4 (9) (2018) eaar8320.
- [6] I.A. Leaf, S. Nakagawa, B.G. Johnson, J.J. Cha, K. Mittelsteadt, K.M. Guckian, I.G. Gomez, W.A. Altemeier, J.S. Duffield, Pericyte MyD88 and IRAK4 control inflammatory and fibrotic responses to tissue injury, *J. Clin. Invest.* 127 (1) (2017) 321–334.
- [7] S. Uderhardt, A.J. Martins, J.S. Tsang, T. Lämmermann, R.N. Germain, Resident macrophages cloak tissue microlesions to prevent neutrophil-driven inflammatory damage, *Cell* 177 (3) (2019) 541–555.
- [8] K.C. Santos Roballo, S. Dhungana, Z. Jiang, J. Oakey, J.S. Bushman, Localized delivery of immunosuppressive regulatory T cells to peripheral nerve allografts promotes regeneration of branched segmental defects, *Biomaterials* 209 (2019) 1–9.
- [9] J. Park, J.T. Decker, D.R. Smith, B.J. Cummings, A.J. Anderson, L.D. Shea, Reducing inflammation through delivery of lentivirus encoding for anti-inflammatory cytokines attenuates neuropathic pain after spinal cord injury, *J. Control. Release* 290 (2018) 88–101.
- [10] U. Park, M.S. Lee, J. Jeon, S. Lee, M.P. Hwang, Y. Wang, H.S. Yang, K. Kim, Coacervate-mediated exogenous growth factor delivery for scarless skin regeneration, *Acta Biomater.* 90 (2019) 179–191.
- [11] Z. Liu, X. Chen, Z. Zhang, X. Zhang, L. Saunders, Y. Zhou, P.X. Ma, Nanofibrous spongy microspheres to distinctly release miRNA and growth factors to enrich regulatory T cells and rescue periodontal bone loss, *ACS Nano* 12 (10) (2018) 9785–9799.
- [12] K. Jiang, J.D. Weaver, Y. Li, X. Chen, J. Liang, C.L. Stabler, Local release of dexamethasone from macroporous scaffolds accelerates islet transplant engraftment by promotion of anti-inflammatory M2 macrophages, *Biomaterials* 114 (2017) 71–81.
- [13] S. Zhang, K.Y.W. Teo, S.J. Chuah, R.C. Lai, S.K. Lim, W.S. Toh, MSC exosomes alleviate temporomandibular joint osteoarthritis by attenuating inflammation and restoring matrix homeostasis, *Biomaterials* 200 (2019) 35–47.

- [14] Q. Liang, Z. Liu, C. Zhu, B. Wang, X. Liu, Y. Yang, X. Lv, H. Mu, K. Wang, Intra-hepatic T-cell receptor β immune repertoire is essential for liver regeneration, *Hepatology* 68 (5) (2018) 1977–1990.
- [15] T.D. Smith, R.R. Nagalla, E.Y. Chen, W.F. Liu, Harnessing macrophage plasticity for tissue regeneration, *Adv. Drug Deliv. Rev.* 114 (2017) 193–205.
- [16] J. Li, J. Tan, M.M. Martino, K.O. Lui, Regulatory T-cells: potential regulator of tissue repair and regeneration, *Front. Immunol.* 9 (2018) 585.
- [17] B.J. Kwee, D.J. Mooney, Biomaterials for skeletal muscle tissue engineering, *Curr. Opin. Biotechnol.* 47 (2017) 16–22.
- [18] J.-E. Won, Y.S. Lee, J.-H. Park, J.-H. Lee, Y.S. Shin, C.-H. Kim, J.C. Knowles, H.-W. Kim, Hierarchical microchanneled scaffolds modulate multiple tissue-regenerative processes of immune-responses, angiogenesis, and stem cell homing, *Biomaterials* 227 (2020) 119548.
- [19] N. Robertson, M. Rappas, A.S. Doré, J. Brown, G. Bottegoni, M. Koglin, J. Cansfield, A. Jazayeri, R.M. Cooke, F.H. Marshall, Structure of the complement C5a receptor bound to the extra-helical antagonist NDT9513727, *Nature* 553 (7686) (2018) 111–114.
- [20] J. Di Domizio, C. Belkhdja, P. Chenuet, A. Fries, T. Murray, P.M. Mondéjar, O. Demaria, C. Conrad, B. Homey, S. Werner, The commensal skin microbiota triggers type I IFN-dependent innate repair responses in injured skin, *Nat. Immunol.* 21 (9) (2020) 1034–1045.
- [21] R.A. Houseright, E.E. Rosowski, P.-Y. Lam, S.J. Tauzin, O. Mulvaney, C.N. Dewey, A. Huttenlocher, Cell type specific gene expression profiling reveals a role for complement component C3 in neutrophil responses to tissue damage, *Sci. Rep.* 10 (1) (2020) 1–15.
- [22] A. Iwasaki, R. Medzhitov, Regulation of adaptive immunity by the innate immune system, *Science* 327 (5963) (2010) 291–295.
- [23] S. Pandey, T. Kawai, S. Akira, Microbial sensing by Toll-like receptors and intracellular nucleic acid sensors, *Cold Spring Harb. Perspect. Biol.* 7 (1) (2015) a016246.
- [24] K. Hayashi, F. Nikolos, Y. Lee, A. Jain, E. Tsouko, H. Gao, A. Kasabian, H. Leung, A. Osipov, S. Jung, Tipping the immunostimulatory and inhibitory DAMP balance to harness immunogenic cell death, *Nat. Commun.* 11 (1) (2020) 1–13.
- [25] K.J. Kieser, J.C. Kagan, Multi-receptor detection of individual bacterial products by the innate immune system, *Nat. Rev. Immunol.* 17 (6) (2017) 376.
- [26] A.J. Russo, S.O. Vasudevan, S.P. Méndez-Huergo, P. Kumari, A. Menoret, S. Duduskar, C. Wang, J.M. Pérez Sáez, M.M. Fettes, C. Li, R. Liu, A. Wanchoo, K. Chandiran, J. Ruan, S.K. Vanaja, M. Bauer, C. Sponholz, G.A. Hudalla, A.T. Vella, B. Zhou, S.D. Deshmukh, G.A. Rabinovich, V.A. Rathinam, Intracellular immune sensing promotes inflammation via gasdermin D-driven release of a lectin alarmin, *Nat. Immunol.* (2021), doi:10.1038/s41590-020-00844-7.
- [27] Y. Xin, Q. Yuan, C. Liu, C. Zhang, D. Yuan, MiR-155/GSK-3 β mediates anti-inflammatory effect of Chikusetsusaponin IVa by inhibiting NF- κ B signaling pathway in LPS-induced RAW264.7 cell, *Sci. Rep.* 10 (1) (2020) 18303.
- [28] O. Takeuchi, S. Akira, Pattern recognition receptors and inflammation, *Cell* 140 (6) (2010) 805–820.
- [29] G.C. Gurtner, S. Werner, Y. Barrandon, M.T. Longaker, Wound repair and regeneration, *Nature* 453 (7193) (2008) 314–321.
- [30] W. Yang, Y. Tao, Y. Wu, X. Zhao, W. Ye, D. Zhao, L. Fu, C. Tian, J. Yang, F. He, L. Tang, Neutrophils promote the development of reparative macrophages mediated by ROS to orchestrate liver repair, *Nat. Commun.* 10 (1) (2019) 1076.
- [31] T.A. Wynn, K.M. Vannella, Macrophages in tissue repair, regeneration, and fibrosis, *Immunity* 44 (3) (2016) 450–462.
- [32] S. Jhunjhunwala, Neutrophils at the biological–material interface, *ACS Biomater. Sci. Eng.* 4 (4) (2017) 1128–1136.
- [33] K. Theilgaard-Mönch, S. Knudsen, P. Follin, N. Borregaard, The transcriptional activation program of human neutrophils in skin lesions supports their important role in wound healing, *J. Immunol.* 172 (12) (2004) 7684–7693.
- [34] L. Ladislau, D.M. Portilho, T. Courau, A. Solares-Pérez, E. Negroni, J. Lainé, D. Klatzmann, A. Bonomo, Y. Allenbach, O. Benveniste, I. Riederer, W. Savino, V. Mouly, G. Butler-Brown, C.F. Benjamim, Activated dendritic cells modulate proliferation and differentiation of human myoblasts, *Cell Death Dis.* 9 (5) (2018).
- [35] A.M. Silva, M.I. Almeida, J.H. Teixeira, A.F. Maia, G.A. Calin, M.A. Barbosa, S.G. Santos, Dendritic cell-derived extracellular vesicles mediate mesenchymal stem/stromal cell recruitment, *Sci. Rep.* 7 (1) (2017) 1–15.
- [36] X. Zhang, D. Mosser, Macrophage activation by endogenous danger signals, *J. Pathol.* 214 (2) (2008) 161–178.
- [37] B. Chazaud, Macrophages: supportive cells for tissue repair and regeneration, *Immunobiology* 219 (3) (2014) 172–178.
- [38] A.D. Lynn, T.R. Kyriakides, S.J. Bryant, Characterization of the in vitro macrophage response and in vivo host response to poly (ethylene glycol)-based hydrogels, *J. Biomed. Mater. Res. A* 93 (3) (2010) 941–953.
- [39] U. Cho, B. Kim, S. Kim, Y. Han, Y.S. Song, Pro-inflammatory M1 Macrophage enhances metastatic potential of ovarian cancer cells through NF- κ B activation, *Mol. Carcinog.* 57 (2) (2018) 235–242.
- [40] A. Kimball, M. Schaller, A. Joshi, F.M. Davis, A. denDekker, A. Boniakowski, J. Bernick, A. Obi, B. Moore, P.K. Henke, Ly6Chi blood monocyte/macrophage drive chronic inflammation and impair wound healing in diabetes mellitus, *Arterioscler. Thromb. Vasc. Biol.* 38 (5) (2018) 1102–1114.
- [41] D.M. Mosser, J.P. Edwards, Exploring the full spectrum of macrophage activation, *Nat. Rev. Immunol.* 8 (12) (2008) 958–969.
- [42] C.E. Olingy, C.L. San Emeterio, M.E. Ogle, J.R. Krieger, A.C. Bruce, D.D. Pfau, B.T. Jordan, S.M. Pearce, E.A. Botchwey, Non-classical monocytes are biased progenitors of wound healing macrophages during soft tissue injury, *Sci. Rep.* 7 (1) (2017) 1–16.
- [43] W.-K. Wu, O.P. Llewellyn, D.O. Bates, L.B. Nicholson, A.D. Dick, IL-10 regulation of macrophage VEGF production is dependent on macrophage polarisation and hypoxia, *Immunobiology* 215 (9–10) (2010) 796–803.
- [44] A. Shapouri-Moghaddam, S. Mohammadian, H. Vazini, M. Taghadosi, S.A. Esmaeili, F. Mardani, B. Seifi, A. Mohammadi, J.T. Afshari, A. Sahebkar, Macrophage plasticity, polarization, and function in health and disease, *J. Cell Physiol.* 233 (9) (2018) 6425–6440.
- [45] T. Roszer, Understanding the mysterious M2 macrophage through activation markers and effector mechanisms, *Mediators Inflamm.* 2015 (2015) 816460.
- [46] M.-Z. Zhang, X. Wang, Y. Wang, A. Niu, S. Wang, C. Zou, R.C. Harris, IL-4/IL-13-mediated polarization of renal macrophages/dendritic cells to an M2a phenotype is essential for recovery from acute kidney injury, *Kidney Int.* 91 (2) (2017) 375–386.
- [47] L.X. Wang, S.X. Zhang, H.J. Wu, X.L. Rong, J. Guo, M2b macrophage polarization and its roles in diseases, *J. Leukoc. Biol.* 106 (2) (2019) 345–358.
- [48] G. Zizzo, B.A. Hilliard, M. Monestier, P.L. Cohen, Efficient clearance of early apoptotic cells by human macrophages requires M2c polarization and MerTK induction, *J. Immunol.* 189 (7) (2012) 3508–3520.
- [49] Q. Wang, H. Ni, L. Lan, X. Wei, R. Xiang, Y. Wang, Fra-1 protooncogene regulates IL-6 expression in macrophages and promotes the generation of M2 macrophages, *Cell Res.* 20 (6) (2010) 701–712.
- [50] D. Duluc, Y. Delneste, F. Tan, M.-P. Moles, L. Grimaud, J. Lenoir, L. Preisser, I. Anegón, L. Catala, N. Ifrah, Tumor-associated leukemia inhibitory factor and IL-6 skew monocyte differentiation into tumor-associated macrophage-like cells, *Blood* 110 (13) (2007) 4319–4330.
- [51] J.P. Pradere, J. Kluewe, S. De Minicis, J.J. Jiao, G.Y. Gwak, D.H. Dapito, M.K. Jang, N.D. Guenther, I. Mederacke, R. Friedman, Hepatic macrophages but not dendritic cells contribute to liver fibrosis by promoting the survival of activated hepatic stellate cells in mice, *Hepatology* 58 (4) (2013) 1461–1473.
- [52] C.E. Witherell, D. Abebayehu, T.H. Barker, K.L. Spiller, Macrophage and fibroblast interactions in biomaterial-mediated fibrosis, *Adv. Healthc. Mater.* 8 (4) (2019) 1801451.
- [53] P.M.-K. Tang, D.J. Nikolic-Paterson, H.-Y. Lan, Macrophages: versatile players in renal inflammation and fibrosis, *Nat. Rev. Nephrol.* 15 (3) (2019) 144–158.
- [54] S. Hong, Z. Zhang, H. Liu, M. Tian, X. Zhu, Z. Zhang, W. Wang, X. Zhou, F. Zhang, Q. Ge, B cells are the dominant antigen-presenting cells that activate naive CD4⁺ T cells upon immunization with a virus-derived nanoparticle antigen, *Immunity* 49 (4) (2018) 695–708.
- [55] L. Chen, D.B. Flies, Molecular mechanisms of T cell co-stimulation and co-inhibition, *Nat. Rev. Immunol.* 13 (4) (2013) 227–242.
- [56] C. June, J. Ledbetter, M. Gillespie, T. Lindsten, C. Thompson, T-cell proliferation involving the CD28 pathway is associated with cyclosporine-resistant interleukin 2 gene expression, *Mol. Cell. Biol.* 7 (12) (1987) 4472–4481.
- [57] P.S. Linsley, E.A. Clark, J.A. Ledbetter, T-cell antigen CD28 mediates adhesion with B cells by interacting with activation antigen B7/BB-1, *Proc. Natl. Acad. Sci. USA* 87 (13) (1990) 5031–5035.
- [58] M. Azuma, D. Ito, H. Yagita, K. Okumura, J. Phillips, L. Lanier, C. Somoza, B70 antigen is a second ligand for CTLA-4 and CD28, *Nature* 366 (6450) (1993) 76–79.
- [59] Y. Dombrowski, T. O'hagan, M. Dittmer, R. Penalva, S.R. Mayoral, P. Bankhead, S. Fleville, G. Eleftheriadis, C. Zhao, M. Naughton, Regulatory T cells promote myelin regeneration in the central nervous system, *Nat. Neurosci.* 20 (5) (2017) 674–680.
- [60] S. Reinke, S. Geissler, W.R. Taylor, K. Schmidt-Bleek, K. Juelke, V. Schwachmeyer, M. Dahne, T. Hartwig, L. Akyüz, C. Meisel, Terminally differentiated CD8⁺ T cells negatively affect bone regeneration in humans, *Sci. Transl. Med.* 5 (177) (2013) 177ra36–177ra36.
- [61] J. Zhang, Z. Xiao, C. Qu, W. Cui, X. Wang, J. Du, CD8 T cells are involved in skeletal muscle regeneration through facilitating MCP-1 secretion and Gr1high macrophage infiltration, *J. Immunol.* 193 (10) (2014) 5149–5160.
- [62] B.J. Kwee, E. Budina, A.J. Najibi, D.J. Mooney, CD4 T-cells regulate angiogenesis and myogenesis, *Biomaterials* 178 (2018) 109–121.
- [63] Y. Liu, L. Wang, T. Kikui, K. Akiyama, C. Chen, X. Xu, R. Yang, W. Chen, S. Wang, S. Shi, Mesenchymal stem cell-based tissue regeneration is governed by recipient T lymphocytes via IFN- γ and TNF- α , *Nat. Med.* 17 (12) (2011) 1594.
- [64] W.H. Mahallawi, O.F. Khabour, Q. Zhang, H.M. Makhadmeh, B.A. Suliman, MER-S-CoV infection in humans is associated with a pro-inflammatory Th1 and Th17 cytokine profile, *Cytokine* 104 (2018) 8–13.
- [65] S.J. Jenkins, D. Ruckerl, P.C. Cook, L.H. Jones, E.D. Finkelman, N. van Rooijen, A.S. MacDonald, J.E. Allen, Local macrophage proliferation, rather than recruitment from the blood, is a signature of Th2 inflammation, *Science* 332 (6035) (2011) 1284–1288.
- [66] R.L. Gieseck III, M.S. Wilson, T.A. Wynn, Type 2 immunity in tissue repair and fibrosis, *Nat. Rev. Immunol.* 18 (1) (2018) 62.
- [67] D. Zemmour, R. Zilionis, E. Kiner, A.M. Klein, D. Mathis, C. Benoist, Single-cell gene expression reveals a landscape of regulatory T cell phenotypes shaped by the TCR, *Nat. Immunol.* 19 (3) (2018) 291–301.
- [68] N. Ali, B. Zirik, R.S. Rodriguez, M.L. Pauli, H.-A. Truong, K. Lai, R. Ahn, K. Corbin, M.M. Lowe, T.C. Scharschmidt, Regulatory T cells in skin facilitate epithelial stem cell differentiation, *Cell* 169 (6) (2017) 1119–1129.
- [69] Y. Liu, R. Yang, S. Shi, Systemic infusion of mesenchymal stem cells improves cell-based bone regeneration via upregulation of regulatory T cells, *Tissue Eng. Part A* 21 (3–4) (2015) 498–509.
- [70] A. Zhang, A. Lacy-Hulbert, S. Anderton, C. Haslett, J. Savill, Apoptotic cell-directed resolution of lung inflammation requires myeloid α v integrin-

- mediated induction of regulatory T lymphocytes, *Am. J. Pathol.* (2020), doi:10.1016/j.ajpath.2020.02.010.
- [71] E.H. Choo, J.-H. Lee, E.-H. Park, H.E. Park, N.-C. Jung, T.-H. Kim, Y.-S. Koh, E. Kim, K.-B. Seung, C. Park, Infarcted myocardium-primed dendritic cells improve remodeling and cardiac function after myocardial infarction by modulating the regulatory T cell and macrophage polarization, *Circulation* 135 (15) (2017) 1444–1457.
 - [72] T.J. Murphy, N.N. Choleain, Y. Zang, J.A. Mannick, J.A. Lederer, CD4+ CD25+ regulatory T cells control innate immune reactivity after injury, *J. Immunol.* 174 (5) (2005) 2957–2963.
 - [73] H. Lei, K. Schmidt-Bleek, A. Dienelt, P. Reinke, H.-D. Volk, Regulatory T cell-mediated anti-inflammatory effects promote successful tissue repair in both indirect and direct manners, *Front. Pharmacol.* 6 (2015) 184.
 - [74] L. Schmidleithner, Y. Thabet, E. Schönfeld, M. Köhne, D. Sommer, Z. Abdullah, T. Sadlon, C. Osei-Sarpong, K. Subbaramaiah, F. Copperi, Enzymatic activity of HPGD in Treg cells suppresses Tconv cells to maintain adipose tissue homeostasis and prevent metabolic dysfunction, *Immunity* 50 (5) (2019) 1232–1248.
 - [75] G. Liu, H. Ma, L. Qiu, L. Li, Y. Cao, J. Ma, Y. Zhao, Phenotypic and functional switch of macrophages induced by regulatory CD4+ CD25+ T cells in mice, *Immunol. Cell Biol.* 89 (1) (2011) 130–142.
 - [76] E.M. Shevach, Mechanisms of foxp3+ T regulatory cell-mediated suppression, *Immunity* 30 (5) (2009) 636–645.
 - [77] A.N. Mathur, B. Zirak, I.C. Boothby, M. Tan, J.N. Cohen, T.M. Mauro, P. Mehta, M.M. Lowe, A.K. Abbas, N. Ali, Treg-cell control of a CXCL5-IL-17 inflammatory axis promotes hair-follicle-stem-cell differentiation during skin-barrier repair, *Immunity* 50 (3) (2019) 655–667.
 - [78] F.R. D'Alessio, K. Tushima, N.R. Aggarwal, E.E. West, M.H. Willett, M.F. Britos, M.R. Pipeling, R.G. Brower, R.M. Tuder, J.F. McDyer, CD4+ CD25+ Foxp3+ Tregs resolve experimental lung injury in mice and are present in humans with acute lung injury, *J. Clin. Invest.* 119 (10) (2009) 2898–2913.
 - [79] J. Weirather, U.D. Hofmann, N. Beyersdorf, G.C. Ramos, B. Vogel, A. Frey, G. Ertl, T. Kerkau, S. Frantz, Foxp3+ CD4+ T cells improve healing after myocardial infarction by modulating monocyte/macrophage differentiation, *Circ. Res.* 115 (1) (2014) 55–67.
 - [80] S. Qin, J.-h. Zheng, Z.-h. Xia, J. Qian, C.-I. Deng, S.-I. Yang, CTHRC1 promotes wound repair by increasing M2 macrophages via regulating the TGF- β and notch pathways, *Biomed. Pharmacother.* 113 (2019) 108594.
 - [81] S. Zhang, Y. Liu, X. Zhang, D. Zhu, X. Qi, X. Cao, Y. Fang, Y. Che, Z.-C. Han, Z.-X. He, Prostaglandin E2 hydrogel improves cutaneous wound healing via M2 macrophages polarization, *Theranostics* 8 (19) (2018) 5348.
 - [82] A. Castiglioni, G. Corna, E. Rigamonti, V. Basso, M. Vezzoli, A. Monno, A.E. Almada, A. Mondino, A.J. Wagers, A.A. Manfredi, FOXP3+ T cells recruited to sites of sterile skeletal muscle injury regulate the fate of satellite cells and guide effective tissue regeneration, *PLoS One* 10 (6) (2015) e0128094.
 - [83] A. Nosbaum, N. Prevel, H.A. Truong, P. Mehta, M. Ettinger, T.C. Scharschmidt, N.H. Ali, M.L. Pauli, A.K. Abbas, M.D. Rosenblum, Cutting edge: regulatory T cells facilitate cutaneous wound healing, *J. Immunol.* 196 (5) (2016) 2010–2014.
 - [84] L.L. McIntyre, S.A. Greilach, S. Othy, I. Sears-Kraxberger, B. Wi, J. Ayala-Angulo, E. Vu, Q. Pham, J. Silva, K. Dang, Regulatory T cells promote remyelination in the murine experimental autoimmune encephalomyelitis model of multiple sclerosis following human neural stem cell transplant, *Neurobiol. Dis.* (2020) 104868.
 - [85] O. Veisheh, J.C. Doloff, M. Ma, A.J. Vegas, H.H. Tam, A.R. Bader, J. Li, E. Langan, J. Wyckoff, W.S. Loo, Size- and shape-dependent foreign body immune response to materials implanted in rodents and non-human primates, *Nat. Mater.* 14 (6) (2015) 643–651.
 - [86] J.C. Doloff, O. Veisheh, A.J. Vegas, H.H. Tam, S. Farah, M. Ma, J. Li, A. Bader, A. Chiu, A. Sadraei, S. Aresta-Dasilva, M. Griffin, S. Jhunjunwala, M. Weber, S. Siebert, K. Tang, M. Chen, E. Langan, N. Dholokia, R. Thakrar, M. Qi, J. Oberholzer, D.L. Greiner, R. Langer, D.G. Anderson, Colony stimulating factor-1 receptor is a central component of the foreign body response to biomaterial implants in rodents and non-human primates, *Nat. Mater.* 16 (6) (2017) 671–680.
 - [87] D. Zhang, Q. Chen, C. Shi, M. Chen, K. Ma, J. Wan, R. Liu, Dealing with the foreign-body response to implanted biomaterials: strategies and applications of new materials, *Adv. Funct. Mater.* (2020) 2007226.
 - [88] A.J. Vegas, O. Veisheh, J.C. Doloff, M. Ma, H.H. Tam, K. Bratlie, J. Li, A.R. Bader, E. Langan, K. Olejnik, Combinatorial hydrogel library enables identification of materials that mitigate the foreign body response in primates, *Nat. Biotechnol.* 34 (3) (2016) 345–352.
 - [89] J. Park, M.H. Gerber, J.E. Babensee, Phenotype and polarization of autologous T cells by biomaterial-treated dendritic cells, *J. Biomed. Mater. Res. A* 103 (1) (2015) 170–184.
 - [90] N.A. Hotaling, L. Tang, D.J. Irvine, J.E. Babensee, Biomaterial strategies for immunomodulation, *Annu. Rev. Biomed. Eng.* 17 (2015) 317–349.
 - [91] Y. Deng, J. Ren, G. Chen, G. Li, X. Wu, G. Wang, G. Gu, J. Li, Injectable in situ cross-linking chitosan-hyaluronic acid based hydrogels for abdominal tissue regeneration, *Sci. Rep.* 7 (1) (2017) 1–13.
 - [92] J. Liu, D. Saul, K.O. Boker, J. Ernst, W. Lehman, A.F. Schilling, Current methods for skeletal muscle tissue repair and regeneration, *Biomed. Res. Int.* 2018 (2018) 1984879.
 - [93] N.J. Shah, A.S. Mao, T.Y. Shih, M.D. Kerr, A. Sharda, T.M. Raimondo, J.C. Weaver, V.D. Vrbancic, M. Deruaz, A.M. Tager, D.J. Mooney, D.T. Scadden, An injectable bone marrow-like scaffold enhances T cell immunity after hematopoietic stem cell transplantation, *Nat. Biotechnol.* 37 (3) (2019) 293–302.
 - [94] A.T. Ho, A.R. Palla, M.R. Blake, N.D. Yucel, Y.X. Wang, K.E. Magnusson, C.A. Holbrook, P.E. Kraft, S.L. Delp, H.M. Blau, Prostaglandin E2 is essential for efficacious skeletal muscle stem-cell function, augmenting regeneration and strength, *Proc. Natl. Acad. Sci. USA* 114 (26) (2017) 6675–6684.
 - [95] G.S. Baht, A. Bareja, D.E. Lee, R.R. Rao, R. Huang, J.L. Huebner, D.B. Bartlett, C.R. Hart, J.R. Gibson, I.R. Lanza, Meteorin-like facilitates skeletal muscle repair through a Stat3/IGF-1 mechanism, *Nat. Metab.* 2 (3) (2020) 278–289.
 - [96] B. Xia, Y. Lv, Dual-delivery of VEGF and NGF by emulsion electrospray nanofibrous scaffold for peripheral nerve regeneration, *Mater. Sci. Eng. C* 82 (2018) 253–264.
 - [97] R. Zhang, L. Xie, H. Wu, T. Yang, Q. Zhang, Y. Tian, Y. Liu, X. Han, W. Guo, M. He, Alginate/aponite hydrogel microspheres co-encapsulating dental pulp stem cells and VEGF for endodontic regeneration, *Acta Biomater.* 113 (2020) 305–316.
 - [98] S. Jimi, A. Jaguparov, A. Nurkesh, B. Sultankulov, A. Saparov, Sequential delivery of cryogel released growth factors and cytokines accelerates wound healing and improves tissue regeneration, *Front. Bioeng. Biotechnol.* 8 (2020) 345.
 - [99] M.C.P. Sok, M.C. Tria, C.E. Olingy, C.L. San Emeterio, E.A. Botchwey, Aspirin-Triggered Resolvin D1-modified materials promote the accumulation of pro-regenerative immune cell subsets and enhance vascular remodeling, *Acta Biomater.* 53 (2017) 109–122.
 - [100] T.B. Wissing, E.E. van Haaften, S.E. Koch, B.D. Ippel, N.A. Kurniawan, C.V. Bouten, A.I. Smits, Hemodynamic loads distinctively impact the secretory profile of biomaterial-activated macrophages—implications for in situ vascular tissue engineering, *Biomater. Sci.* 8 (1) (2020) 132–147.
 - [101] W. Sun, H. Zhang, H. Wang, Y.G. Chiu, M. Wang, C.T. Ritchlin, A. Kiernan, B.F. Boyce, L. Xing, Targeting notch-activated M1 macrophages attenuates joint tissue damage in a mouse model of inflammatory arthritis, *J. Bone Miner. Res.* 32 (7) (2017) 1469–1480.
 - [102] W. Wang, L. Sun, P. Zhang, J. Song, W. Liu, An anti-inflammatory cell-free collagen/resveratrol scaffold for repairing osteochondral defects in rabbits, *Acta Biomater.* 10 (12) (2014) 4983–4995.
 - [103] M. Jung, Y. Ma, R.P. Iyer, K.Y. DeLeon-Pennell, A. Yabluchanskiy, M.R. Garrett, M.L. Lindsey, IL-10 improves cardiac remodeling after myocardial infarction by stimulating M2 macrophage polarization and fibroblast activation, *Basic Res. Cardiol.* 112 (3) (2017) 33.
 - [104] B.N. Brown, R. Londono, S. Tottey, L. Zhang, K.A. Kukla, M.T. Wolf, K.A. Daly, J.E. Reing, S.F. Badylak, Macrophage phenotype as a predictor of constructive remodeling following the implantation of biologically derived surgical mesh materials, *Acta Biomater.* 8 (3) (2012) 978–987.
 - [105] L. Wu, A. Magaz, E. Maughan, N. Oliver, A. Darbyshire, M. Loizidou, M. Emberton, M. Birchall, W. Song, Cellular responses to thermoresponsive stiffness memory elastomer nanohybrid scaffolds by 3D-TIPS, *Acta Biomater.* 85 (2019) 157–171.
 - [106] C. Yang, C. Zhao, X. Wang, M. Shi, Y. Zhu, L. Jing, C. Wu, J. Chang, Stimulation of osteogenesis and angiogenesis by micro/nano hierarchical hydroxyapatite via macrophage immunomodulation, *Nanoscale* 11 (38) (2019) 17699–17708.
 - [107] M.L. Pinto, E. Rios, A.C. Silva, S.C. Neves, H.R. Caires, A.T. Pinto, C. Duraes, F.A. Carvalho, A.P. Cardoso, N.C. Santos, C.C. Barrias, D.S. Nascimento, O.P. Pinto-do, M.A. Barbosa, F. Carneiro, M.J. Oliveira, Decellularized human colorectal cancer matrices polarize macrophages towards an anti-inflammatory phenotype promoting cancer cell invasion via CCL18, *Biomaterials* 124 (2017) 211–224.
 - [108] F.Y. McWhorter, T. Wang, P. Nguyen, T. Chung, W.F. Liu, Modulation of macrophage phenotype by cell shape, *Proc. Natl. Acad. Sci. USA* 110 (43) (2013) 17253–17258.
 - [109] Z. Chen, S. Ni, S. Han, R. Crawford, S. Lu, F. Wei, J. Chang, C. Wu, Y. Xiao, Nanoporous microstructures mediate osteogenesis by modulating the osteo-immune response of macrophages, *Nanoscale* 9 (2) (2017) 706–718.
 - [110] Y. Jia, W. Yang, K. Zhang, S. Qiu, J. Xu, C. Wang, Y. Chai, Nanofiber arrangement regulates peripheral nerve regeneration through differential modulation of macrophage phenotypes, *Acta Biomater.* 83 (2019) 291–301.
 - [111] A. Mantovani, S.K. Biswas, M.R. Galdiero, A. Sica, M. Locati, Macrophage plasticity and polarization in tissue repair and remodelling, *J. Pathol.* 229 (2) (2013) 176–185.
 - [112] K. Garg, N.A. Pullen, C.A. Oskeritzian, J.J. Ryan, G.L. Bowlin, Macrophage functional polarization (M1/M2) in response to varying fiber and pore dimensions of electrospray scaffolds, *Biomaterials* 34 (18) (2013) 4439–4451.
 - [113] R. Sridharan, B. Cavanagh, A.R. Cameron, D.J. Kelly, F.J. O'Brien, Material stiffness influences the polarization state, function and migration mode of macrophages, *Acta Biomater.* 89 (2019) 47–59.
 - [114] L. Ponsonnet, K. Reybier, N. Jaffrezic, V. Comte, C. Lagneau, M. Lissac, C. Martelet, Relationship between surface properties (roughness, wettability) of titanium and titanium alloys and cell behaviour, *Mater. Sci. Eng. C* 23 (4) (2003) 551–560.
 - [115] A.K. Refai, M. Textor, D.M. Brunette, J.D. Waterfield, Effect of titanium surface topography on macrophage activation and secretion of proinflammatory cytokines and chemokines, *J. Biomed. Mater. Res. A* 70 (2) (2004) 194–205.
 - [116] A.-K. Fuchs, T. Syrovets, K.A. Haas, C. Loos, A. Musyanovych, V. Mailänder, K. Landfester, T. Simmet, Carboxyl- and amino-functionalized polystyrene nanoparticles differentially affect the polarization profile of M1 and M2 macrophage subsets, *Biomaterials* 85 (2016) 78–87.
 - [117] H.M. Rostam, S. Singh, F. Salazar, P. Magennis, A. Hook, T. Singh, N.E. Vrana, M.R. Alexander, A.M. Ghaemmaghami, The impact of surface chemistry

- modification on macrophage polarisation, *Immunobiology* 221 (11) (2016) 1237–1246.
- [118] J.E. McBane, L.A. Matheson, S. Sharifpoor, J.P. Santerre, R.S. Labow, Effect of polyurethane chemistry and protein coating on monocyte differentiation towards a wound healing phenotype macrophage, *Biomaterials* 30 (29) (2009) 5497–5504.
- [119] J.L. Dziki, L. Huleihel, M.E. Scarritt, S.F. Badylak, Extracellular matrix bioscaffolds as immunomodulatory biomaterials, *Tissue Eng. Part A* 23 (19–20) (2017) 1152–1159.
- [120] T.J. Keane, R. Londono, N.J. Turner, S.F. Badylak, Consequences of ineffective decellularization of biologic scaffolds on the host response, *Biomaterials* 33 (6) (2012) 1771–1781.
- [121] S.F. Badylak, D.O. Freytes, T.W. Gilbert, Extracellular matrix as a biological scaffold material: structure and function, *Acta Biomater.* 5 (1) (2009) 1–13.
- [122] F. Qu, J.L. Holloway, J.L. Esterhai, J.A. Burdick, R.L. Mauck, Programmed biomolecule delivery to enable and direct cell migration for connective tissue repair, *Nat. Commun.* 8 (1) (2017) 1–11.
- [123] J. Schnieder, A. Mamazhakypov, A. Birnhuber, J. Wilhelm, G. Kwapiszewska, C. Ruppert, P. Markart, L. Wujak, K. Rubio, G. Barreto, Loss of LRP1 promotes acquisition of contractile-myofibroblast phenotype and release of active TGF- β 1 from ECM stores, *Matrix Biol.* 88 (2020) 69–88.
- [124] L. Huleihel, G.S. Hussey, J.D. Naranjo, L. Zhang, J.L. Dziki, N.J. Turner, D.B. Stolz, S.F. Badylak, Matrix-bound nanovesicles within ECM bioscaffolds, *Sci. Adv.* 2 (6) (2016) e1600502.
- [125] B.M. Sicari, J.L. Dziki, B.F. Siu, C.J. Medberry, C.L. Dearth, S.F. Badylak, The promotion of a constructive macrophage phenotype by solubilized extracellular matrix, *Biomaterials* 35 (30) (2014) 8605–8612.
- [126] J.L. Dziki, D.S. Wang, C. Pineda, B.M. Sicari, T. Rausch, S.F. Badylak, Solubilized extracellular matrix bioscaffolds derived from diverse source tissues differentially influence macrophage phenotype, *J. Biomed. Mater. Res. A* 105 (1) (2017) 138–147.
- [127] C.L. Dearth, P.F. Slivka, S.A. Stewart, T.J. Keane, J.K. Tay, R. Londono, Q. Goh, F.X. Pizsa, S.F. Badylak, Inhibition of COX1/2 alters the host response and reduces ECM scaffold mediated constructive tissue remodeling in a rodent model of skeletal muscle injury, *Acta Biomater.* 31 (2016) 50–60.
- [128] Z. Fan, J. Deng, P.Y. Li, D.R. Chery, Y. Su, P. Zhu, T. Kambayashi, E.P. Blankenhorn, L. Han, H. Cheng, A new class of biological materials: cell membrane-derived hydrogel scaffolds, *Biomaterials* 197 (2019) 244–254.
- [129] J.E. Rayahin, J.S. Buhrman, Y. Zhang, T.J. Koh, R.A. Gemeinhart, High and low molecular weight hyaluronic acid differentially influence macrophage activation, *ACS Biomater. Sci. Eng.* 1 (7) (2015) 481–493.
- [130] H. Siiskonen, S. Oikari, S. Pasonen-Seppanen, K. Rilla, Hyaluronan synthase 1: a mysterious enzyme with unexpected functions, *Front. Immunol.* 6 (2015) 43.
- [131] M. Waters, P. VandeVord, M. Van Dyke, Keratin biomaterials augment anti-inflammatory macrophage phenotype in vitro, *Acta Biomater.* 66 (2018) 213–223.
- [132] Y. Zhou, C. Wu, J. Chang, Bioceramics to regulate stem cells and their microenvironment for tissue regeneration, *Mater. Today* 24 (2019) 41–56.
- [133] Z. Chen, T. Klein, R.Z. Murray, R. Crawford, J. Chang, C. Wu, Y. Xiao, Osteoimmunomodulation for the development of advanced bone biomaterials, *Mater. Today* 19 (6) (2016) 304–321.
- [134] D. Zhai, L. Chen, Y. Chen, Y. Zhu, Y. Xiao, C. Wu, Lithium silicate-based bioceramics promoting chondrocyte maturation by immunomodulating M2 macrophage polarization, *Biomater. Sci.* 8 (16) (2020) 4521–4534.
- [135] Y. Zhou, S. Han, L. Xiao, P. Han, S. Wang, J. He, J. Chang, C. Wu, Y. Xiao, Accelerated host angiogenesis and immune responses by ion release from mesoporous bioactive glass, *J. Mater. Chem. B* 6 (20) (2018) 3274–3284.
- [136] X. Dong, J. Chang, H. Li, Bioglass promotes wound healing through modulating the paracrine effects between macrophages and repairing cells, *J. Mater. Chem. B* 5 (26) (2017) 5240–5250.
- [137] M. Shi, L. Xia, Z. Chen, F. Lv, H. Zhu, F. Wei, S. Han, J. Chang, Y. Xiao, C. Wu, Europium-doped mesoporous silica nanosphere as an immune-modulating osteogenesis/angiogenesis agent, *Biomaterials* 144 (2017) 176–187.
- [138] M. Shi, Z. Chen, S. Farnaghi, T. Friis, X. Mao, Y. Xiao, C. Wu, Copper-doped mesoporous silica nanospheres, a promising immunomodulatory agent for inducing osteogenesis, *Acta Biomater.* 30 (2016) 334–344.
- [139] Z. Chen, A. Bachhuka, S. Han, F. Wei, S. Lu, R.M. Visalakshan, K. Vasilev, Y. Xiao, Tuning chemistry and topography of nanoengineered surfaces to manipulate immune response for bone regeneration applications, *ACS Nano* 11 (5) (2017) 4494–4506.
- [140] A. Bachhuka, R. Madathiparambil Visalakshan, C. Law, A. Santos, H. Ebendorf-Heidepriem, S. Karnati, K. Vasilev, Modulation of macrophages differentiation by nanoscale-engineered geometric and chemical features, *ACS Appl. Bio Mater.* 3 (3) (2020) 1496–1505.
- [141] D. Kwon, B.G. Cha, Y. Cho, J. Min, E.-B. Park, S.-J. Kang, J. Kim, Extra-large pore mesoporous silica nanoparticles for directing in vivo M2 macrophage polarization by delivering IL-4, *Nano Lett.* 17 (5) (2017) 2747–2756.
- [142] K.L. Singampalli, S. Balaji, X. Wang, U.M. Parikh, A. Kaul, J. Gilley, R.K. Birla, P.L. Bollyky, S.G. Keswani, The role of an IL-10/hyaluronan axis in dermal wound healing, *Front. Cell Dev. Biol.* 8 (636) (2020) 1–15.
- [143] M. Monavarian, S. Kader, S. Moenizadeh, E. Jabbari, Regenerative scar-free skin wound healing, *Tissue Eng. Part B Rev.* 25 (4) (2019) 294–311.
- [144] K.L. Spiller, S. Nassiri, C.E. Witherell, R.R. Anfang, J. Ng, K.R. Nakazawa, T. Yu, G. Vunjak-Novakovic, Sequential delivery of immunomodulatory cytokines to facilitate the M1-to-M2 transition of macrophages and enhance vascularization of bone scaffolds, *Biomaterials* 37 (2015) 194–207.
- [145] Z. Hu, C. Ma, X. Rong, S. Zou, X. Liu, Immunomodulatory ECM-like microspheres for accelerated bone regeneration in diabetes mellitus, *ACS Appl. Mater. Interfaces* 10 (3) (2018) 2377–2390.
- [146] V. Bonito, A. Smits, O. Goor, B.D. Ippel, A. Driessen-Mol, T. Munker, A.W. Bosman, T. Mes, P.Y.W. Dankers, C.V.C. Bouten, Modulation of macrophage phenotype and protein secretion via heparin-IL-4 functionalized supramolecular elastomers, *Acta Biomater.* 71 (2018) 247–260.
- [147] R.M. Gower, R.M. Boehler, S.M. Azarin, C.F. Ricci, J.N. Leonard, L.D. Shea, Modulation of leukocyte infiltration and phenotype in microporous tissue engineering scaffolds via vector induced IL-10 expression, *Biomaterials* 35 (6) (2014) 2024–2031.
- [148] C.A. Holladay, A.M. Duffy, X. Chen, M.V. Sefton, T.D. O'Brien, A.S. Pandit, Recovery of cardiac function mediated by MSC and interleukin-10 plasmid functionalized scaffold, *Biomaterials* 33 (5) (2012) 1303–1314.
- [149] S. Jain, T.H. Tran, M. Amiji, Macrophage repolarization with targeted alginate nanoparticles containing IL-10 plasmid DNA for the treatment of experimental arthritis, *Biomaterials* 61 (2015) 162–177.
- [150] T. Li, M. Peng, Z. Yang, X. Zhou, Y. Deng, C. Jiang, M. Xiao, J. Wang, 3D-printed IFN- γ -loading calcium silicate- β -tricalcium phosphate scaffold sequentially activates M1 and M2 polarization of macrophages to promote vascularization of tissue engineering bone, *Acta Biomater.* 71 (2018) 96–107.
- [151] M.M. Alvarez, J.C. Liu, G. Trujillo-de Santiago, B.-H. Cha, A. Vishwakarma, A.M. Ghaemmaghami, A. Khademhosseini, Delivery strategies to control inflammatory response: modulating M1–M2 polarization in tissue engineering applications, *J. Control. Release* 240 (2016) 349–363.
- [152] D. Hachim, S.T. LoPresti, C.C. Yates, B.N. Brown, Shifts in macrophage phenotype at the biomaterial interface via IL-4 eluting coatings are associated with improved implant integration, *Biomaterials* 112 (2017) 95–107.
- [153] H. Knopf-Marques, S. Singh, S.S. Htwe, L. Wolfova, R. Buffa, J. Bacharouche, G.-G. Francius, J.-C. Voegel, P. Schaaf, A.M. Ghaemmaghami, Immunomodulation with self-crosslinked polyelectrolyte multilayer-based coatings, *Biomacromolecules* 17 (6) (2016) 2189–2198.
- [154] J.M.H. Liu, X. Zhang, S. Joe, X. Luo, L.D. Shea, Evaluation of biomaterial scaffold delivery of IL-33 as a localized immunomodulatory agent to support cell transplantation in adipose tissue, *J. Immunol. Regen. Med.* 1 (2018) 1–12.
- [155] F. Zhao, B. Lei, X. Li, Y. Mo, R. Wang, D. Chen, X. Chen, Promoting in vivo early angiogenesis with sub-micrometer strontium-contained bioactive microspheres through modulating macrophage phenotypes, *Biomaterials* 178 (2018) 36–47.
- [156] J. Chen, M. Li, C. Yang, X. Yin, K. Duan, J. Wang, B. Feng, Macrophage phenotype switch by sequential action of immunomodulatory cytokines from hydrogel layers on titania nanotubes, *Colloids Surf. B* 163 (2018) 336–345.
- [157] M. Takewaki, M. Kajiyu, K. Takeda, S. Sasaki, S. Motoike, N. Komatsu, S. Matsuda, K. Ouhara, N. Mizuno, T. Fujita, MSC/ECM cellular complexes induce periodontal tissue regeneration, *J. Dent. Res.* 96 (9) (2017) 984–991.
- [158] V. Bunpetch, Z.-Y. Zhang, X. Zhang, S. Han, P. Zongyou, H. Wu, O. Hong-Wei, Strategies for MSC expansion and MSC-based microtissue for bone regeneration, *Biomaterials* 196 (2019) 67–79.
- [159] H. Xia, X. Li, W. Gao, X. Fu, R.H. Fang, L. Zhang, K. Zhang, Tissue repair and regeneration with endogenous stem cells, *Nat. Rev. Mater.* 3 (7) (2018) 174–193.
- [160] J. Pajarinen, T. Lin, E. Gibon, Y. Kohno, M. Maruyama, K. Nathan, L. Lu, Z. Yao, S.B. Goodman, Mesenchymal stem cell-macrophage crosstalk and bone healing, *Biomaterials* 196 (2019) 80–89.
- [161] K.M. Hotchkiss, N.M. Clark, R. Olivares-Navarrete, Macrophage response to hydrophilic biomaterials regulates MSC recruitment and T-helper cell populations, *Biomaterials* 182 (2018) 202–215.
- [162] E. Seebach, H. Freischmidt, J. Holschbach, J. Fellenberg, W. Richter, Mesenchymal stroma cells trigger early attraction of M1 macrophages and endothelial cells into fibrin hydrogels, stimulating long bone healing without long-term engraftment, *Acta Biomater.* 10 (11) (2014) 4730–4741.
- [163] L. Saldaña, F. Bensiamar, G. Vallés, F.J. Mancebo, E. García-Rey, N. Vilaboa, Immunoregulatory potential of mesenchymal stem cells following activation by macrophage-derived soluble factors, *Stem Cell Res. Ther.* 10 (1) (2019) 1–15.
- [164] K. Hyvärinen, M. Holopainen, V. Skirdenko, H. Ruhanen, P. Lehenkari, M. Korhonen, R. Kälälä, S. Laitinen, E. Kerkelä, Mesenchymal stromal cells and their extracellular vesicles enhance the anti-inflammatory phenotype of regulatory macrophages by downregulating the production of interleukin (IL)-23 and IL-22, *Front. Immunol.* 9 (2018) 771.
- [165] X. Qiu, S. Liu, H. Zhang, B. Zhu, Y. Su, C. Zheng, R. Tian, M. Wang, H. Kuang, X. Zhao, Mesenchymal stem cells and extracellular matrix scaffold promote muscle regeneration by synergistically regulating macrophage polarization toward the M2 phenotype, *Stem Cell Res. Ther.* 9 (1) (2018) 1–15.
- [166] D.P. Vasconcelos, M. Costa, I.F. Amaral, M.A. Barbosa, A.P. Aguiar, J.N. Barbosa, Modulation of the inflammatory response to chitosan through M2 macrophage polarization using pro-resolution mediators, *Biomaterials* 37 (2015) 116–123.
- [167] C. Yin, Q. Zhao, W. Li, Z. Zhao, J. Wang, T. Deng, P. Zhang, K. Shen, Z. Li, Y. Zhang, Biomimetic anti-inflammatory nano-capsule serves as a cytokine blocker and M2 polarization inducer for bone tissue repair, *Acta Biomater.* 102 (2020) 416–426.
- [168] J. Shi, X. Zhang, L. Jiang, L. Zhang, Y. Dong, A.C. Midgley, D. Kong, S. Wang, Regulation of the inflammatory response by vascular grafts modified with Aspirin-Triggered Resolvin D1 promotes blood vessel regeneration, *Acta Biomater.* 97 (2019) 360–373.

- [169] B. Saleh, H.K. Dhaliwal, R. Portillo-Lara, E. Shirzaei Sani, R. Abdi, M.M. Amiji, N. Annabi, Local immunomodulation using an adhesive hydrogel loaded with miRNA-laden nanoparticles promotes wound healing, *Small* 15 (36) (2019) 1902232.
- [170] S.H. Taj, W. Kho, M. Aswendt, F.M. Collmann, C. Green, J. Adamczak, A. Tennstaedt, M. Hoehn, Dynamic modulation of microglia/macrophage polarization by miR-124 after focal cerebral ischemia, *J. Neuroimmune Pharmacol.* 11 (4) (2016) 733–748.
- [171] J. Zheng, Y. Kong, X. Hu, Z. Li, Y. Li, Y. Zhong, X. Wei, J. Ling, MicroRNA-enriched small extracellular vesicles possess odonto-immunomodulatory properties for modulating the immune response of macrophages and promoting odontogenesis, *Stem Cell Res. Ther.* 11 (1) (2020) 517.
- [172] X. Wu, Y. Dai, Y. Yang, C. Huang, X. Meng, B. Wu, J. Li, Emerging role of microRNAs in regulating macrophage activation and polarization in immune response and inflammation, *Immunology* 148 (3) (2016) 237–248.
- [173] T. Desgeorges, G. Caratti, R. Mounier, J. Tuckermann, B. Chazaud, Glucocorticoids shape macrophage phenotype for tissue repair, *Front. Immunol.* 10 (2019) 1591.
- [174] M. Kumar, P. Gupta, S. Bhattacharjee, S.K. Nandi, B.B. Mandal, Immunomodulatory injectable silk hydrogels maintaining functional islets and promoting anti-inflammatory M2 macrophage polarization, *Biomaterials* 187 (2018) 1–17.
- [175] S.P. Hui, D.Z. Sheng, K. Sugimoto, A. Gonzalez-Rajal, S. Nakagawa, D. Hesselson, K. Kikuchi, Zebrafish regulatory T cells mediate organ-specific regenerative programs, *Dev. Cell* 43 (6) (2017) 659–672.
- [176] P. Soroosh, T.A. Doherty, W. Duan, A.K. Mehta, H. Choi, Y.F. Adams, Z. Mikulski, N. Khorram, P. Rosenthal, D.H. Broide, Lung-resident tissue macrophages generate Foxp3+ regulatory T cells and promote airway tolerance, *J. Exp. Med.* 210 (4) (2013) 775–788.
- [177] M. Watanabe, Y. Lu, M. Breen, R.J. Hodes, B7-CD28 co-stimulation modulates central tolerance via thymic clonal deletion and Treg generation through distinct mechanisms, *Nat. Commun.* 11 (1) (2020) 1–14.
- [178] Q. Wen, Y. Kong, H.Y. Zhao, Y.Y. Zhang, T.T. Han, Y. Wang, L.P. Xu, X.H. Zhang, X.J. Huang, G-CSF-induced macrophage polarization and mobilization may prevent acute graft-versus-host disease after allogeneic hematopoietic stem cell transplantation, *Bone Marrow Transplant* 54 (9) (2019) 1419–1433.
- [179] M. Li, X. Zhang, X. Zheng, D. Lian, Z.-X. Zhang, W. Ge, J. Yang, C. Vladau, M. Suzuki, D. Chen, Immune modulation and tolerance induction by RelB-silenced dendritic cells through RNA interference, *J. Immunol.* 178 (9) (2007) 5480–5487.
- [180] M.D. McHugh, J. Park, R. Uhrich, W. Gao, D.A. Horwitz, T.M. Fahmy, Paracrine co-delivery of TGF- β and IL-2 using CD4-targeted nanoparticles for induction and maintenance of regulatory T cells, *Biomaterials* 59 (2015) 172–181.
- [181] M. Panduro, C. Benoist, D. Mathis, Treg cells limit IFN- γ production to control macrophage accrual and phenotype during skeletal muscle regeneration, *Proc. Natl. Acad. Sci. USA* 115 (11) (2018) E2585–E2593.
- [182] S.G. Zheng, J. Wang, P. Wang, J.D. Gray, D.A. Horwitz, IL-2 is essential for TGF- β to convert naive CD4+CD25– cells to CD25+Foxp3+ regulatory T cells and for expansion of these cells, *J. Immunol.* 178 (4) (2007) 2018–2027.
- [183] K. Freudenberg, N. Lindner, S. Dohnke, A.I. Garbe, S. Schallenberg, K. Kretschmer, Critical role of TGF- β and IL-2 receptor signaling in Foxp3 induction by an inhibitor of DNA methylation, *Front. Immunol.* 9 (2018) 125.
- [184] T.R. Malek, A. Yu, V. Vincek, P. Scibelli, L. Kong, CD4 regulatory T cells prevent lethal autoimmunity in IL-2R β -deficient mice: implications for the nonredundant function of IL-2, *Immunity* 17 (2) (2002) 167–178.
- [185] S.P. Gadani, J.T. Walsh, I. Smirnov, J. Zheng, J. Kipnis, The glia-derived alarmin IL-33 orchestrates the immune response and promotes recovery following CNS injury, *Neuron* 85 (4) (2015) 703–709.
- [186] D. Kolodin, N. van Panhuys, C. Li, A.M. Magnuson, D. Cipolletta, C.M. Miller, A. Wagers, R.N. Germain, C. Benoist, D. Mathis, Antigen- and cytokine-driven accumulation of regulatory T cells in visceral adipose tissue of lean mice, *Cell Metab.* 21 (4) (2015) 543–557.
- [187] D.C. Nascimento, P.H. Melo, A.R. Pineros, R.G. Ferreira, D.F. Colón, P.B. Donate, F.V. Castanheira, A. Gozzi, P.G. Czaikoski, W. Niedbala, IL-33 contributes to sepsis-induced long-term immunosuppression by expanding the regulatory T cell population, *Nat. Commun.* 8 (1) (2017) 1–14.
- [188] C. Schiering, T. Krausgruber, A. Chomka, A. Fröhlich, K. Adelman, E.A. Wohlfert, J. Pott, T. Griseri, J. Bollrath, A.N. Hegazy, The alarmin IL-33 promotes regulatory T-cell function in the intestine, *Nature* 513 (7519) (2014) 564–568.
- [189] A. Vasanthakumar, K. Moro, A. Xin, Y. Liao, R. Gloury, S. Kawamoto, S. Fagarasan, L.A. Mielke, S. Afshar-Sterle, S.L. Masters, The transcriptional regulators IRF4, BATF and IL-33 orchestrate development and maintenance of adipose tissue-resident regulatory T cells, *Nat. Immunol.* 16 (3) (2015) 276–285.
- [190] Q. Huang, X. Ma, Y. Wang, Z. Niu, R. Wang, F. Yang, M. Wu, G. Liang, P. Rong, H. Wang, D.C. Harris, W. Wang, Q. Cao, IL-10 producing type 2 innate lymphoid cells prolong islet allograft survival, *EMBO Mol. Med.* 12 (11) (2020) e12305.
- [191] K. Kawai, M. Uchiyama, J. Hester, F. Issa, IL-33 drives the production of mouse regulatory T cells with enhanced *in vivo* suppressive activity in skin transplantation, *Am. J. Transplant.* (2020), doi:10.1111/ajt.16266.
- [192] A.J. Lam, K.N. MacDonald, A.M. Pesenacker, S.C. Juvet, K.A. Morishita, B. Bressler, C. iGemoMed, J.G. Pan, S.S. Sidhu, J.D. Rioux, M.K. Levings, Innate control of tissue-reparative human regulatory T cells, *J. Immunol.* 202 (8) (2019) 2195–2209.
- [193] W. Kuswanto, D. Burzyn, M. Panduro, K.K. Wang, Y.C. Jang, A.J. Wagers, C. Benoist, D. Mathis, Poor repair of skeletal muscle in aging mice reflects a defect in local, interleukin-33-dependent accumulation of regulatory T cells, *Immunity* 44 (2) (2016) 355–367.
- [194] C. Roux, G. Saviane, J. Pini, N. Belaid, G. Dhib, C. Voha, L. Ibanez, A. Boutin, N.M. Mazure, A. Wakkach, C. Blin-Wakkach, M. Rouleau, Immunosuppressive mesenchymal stromal cells derived from human-induced pluripotent stem cells induce human regulatory T cells *in vitro* and *in vivo*, *Front. Immunol.* 8 (2017) 1991.
- [195] S. Cosenza, K. Toupet, M. Maumus, P. Luz-Crawford, O. Blanc-Brude, C. Jorgensen, D. Noel, Mesenchymal stem cells-derived exosomes are more immunosuppressive than microparticles in inflammatory arthritis, *Theranostics* 8 (5) (2018) 1399–1410.