

Interactive Assignment 5

6 pages

Homework problem 7.1 p 1

Homework problem 7.4 p 3

Yves '68ATTU

Put the ODE in standard form:

$$y'' + \frac{3z}{1-z^2} y' + \frac{1}{1-z^2} y = 0$$

thus $p(z) = -\frac{3z}{1-z^2}$ and $q(z) = \frac{1}{1-z^2}$

$p(0)=0$ and $q(0)=1$ then $z=0$ is an ordinary point, we can then represent the solutions in terms of a power series:

$$y(z) = \sum_{n=0}^{\infty} a_n z^n$$

$$y'(z) = \sum_{n=0}^{\infty} n a_n z^{n-1}$$

$$y''(z) = \sum_{n=0}^{\infty} n(n-1) a_n z^{n-2}$$

Plug back these into the initial ODE gives:

$$(1-z^2) \sum_{n=0}^{\infty} n(n-1) a_n z^{n-2} - 3z \sum_{n=0}^{\infty} n a_n z^{n-1} + \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n z^{n-2} - \sum_{n=0}^{\infty} n(n-1) a_n z^n - 3 \sum_{n=0}^{\infty} n a_n z^n + \sum_{n=0}^{\infty} a_n z^n = 0$$

Reindexing the first term:

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n - \sum_{n=0}^{\infty} n(n-1) a_n z^n - 3 \sum_{n=0}^{\infty} n a_n z^n + \sum_{n=0}^{\infty} a_n z^n = 0$$

Chapter 7- Problem 7.1

Collecting the coefficients for same power of z :

$$\sum_{n=0}^{\infty} [(n+2)(n+1)a_{n+2} - (n^2 + 2n - d)a_n] z^n = 0$$

Discarding the degenerate solution for $z^n = 0$ yields the recurrent relation:

$$a_{n+2} = \frac{n^2 + 2n - d}{(n+1)(n+2)} a_n$$

Note that for $d = n(n+2)$ all the a_m terms for $m > n$ are 0!
The solution in this case is the polynomial of degree n : $y(z) = \sum_{k=0}^n a_k z^k$

For $n=2$, $d=8$, take $a_1=0$ and $a_2 = \frac{(-8)}{2+1} a_0 = -4a_0$
 $a_0 \neq 0$

$$\text{Therefore for } n=2, \text{ we get } y(z) = U_2(z) = a_0 - 4a_0 z^2 \\ = a_0 (1 - 4z^2)$$

For $n=3$, $d=3(3+2)=15$, take $a_0=0$ and $a_1 \neq 0$ to generate a power series with odd terms

$$a_{d+2} = a_3 = \frac{3-15}{(1+2)(1+1)} a_1 = -\frac{12}{6} a_1 = -2a_1$$

$$\text{Therefore for } n=3 \quad y(z) = U_3(z) = a_1 (z - 2z^3)$$

Chapter 7 - Problem 7.4

By the change of variable $x = z - a$ and using the chain rule:

$$\frac{df}{dz} = \frac{df}{dx} \frac{dx}{dz} = \frac{df}{dx} \cdot 1 = \frac{df}{dx}$$

$$\frac{d^2f}{dz^2} = \frac{dx}{dz} \frac{d}{dx} \left(\frac{df}{dz} \right) = 1 \cdot \frac{d}{dx} \left(\frac{df}{dx} \right) = \frac{d^2f}{dx^2}$$

Therefore the ODE becomes:

$$\frac{d^2f}{dz^2} + 2x \frac{df}{dz} + 4f = 0$$

The solution can be represented at any point x as:

$$f(x) = \sum_{n=0}^{\infty} a_n x^n \quad \text{since there is no singular point.}$$

$$f'(x) = \sum_{n=0}^{\infty} n a_n x^{n-1} \quad \text{and} \quad f''(x) = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

Substitution of these derivatives into the ODE gives

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 2x \sum_{n=0}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\Rightarrow \sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + (2n+4) a_n] x^n = 0$$

$$\Rightarrow \sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + 2(n+2) a_n] x^n = 0$$

Chapter 7 - Problem 7.4

Avoiding the degenerate solution $x=0$, for each n

$$(n+2)(n+1)a_{n+2} + 2(n+2)a_n = 0$$

for all n :
$$a_{n+2} = -\frac{2}{n+1}a_n$$

For even n : $n=2p$
$$\begin{aligned} a_{2p} &= -\frac{2}{2p-1}a_{2p-2} = \frac{(-2)}{2p-1} \frac{(-2)}{(2p-3)} a_{2p-4} \\ &= \frac{(-2)(-2)(-2)(-2)}{(2p-1)(2p-3)(2p-5)(2p-7)} a_{2p-8} \\ &= \dots = \frac{(-2)^{p+1}}{(2p-1)(2p-3)\dots 3} a_2 \\ &= \frac{(-2)^{p+2}}{(2p-1)(2p-3)\dots 3} a_0 \\ &= \frac{(-2)^p 2p (2p-2)(2p-4)\dots 2}{(2p)!} a_0 \\ &= \frac{(-2)^p 2^p p!}{(2p)!} a_0 = \frac{(-4)^p p!}{(2p)!} a_0 \end{aligned}$$

Shifting the indices:

$$a_n = \frac{-2}{n-1} a_{n-2}$$

And for odd n :
$$a_{2p-1} = \frac{-2}{2p-1-1} a_{2p-3} = \frac{-2}{2(p-1)} a_{2p-3}$$

Lecture 7 - Problem 7.4

We have

$$\begin{aligned} a_{2p+1} &= \frac{(-2)}{2p} a_{2p-1} = \frac{(-1)}{p} a_{2p-1} = \frac{(-1)(-1)}{p(p-1)} a_{2p-3} \\ &= \dots = \frac{(-1) \dots (-1)}{p(p-1)(p-2) \dots 2 \cdot 1} a_1 = \frac{(-1)^p}{p!} a_1 \end{aligned}$$

We want to generate two independent solutions: So for one case set $a_1 = 0$ then $a_{2p+1} = 0$ and $a_{2p} = \frac{(-1)^p p!}{2p!}$

This independent solution is of the form:

$$\begin{aligned} f(x) &= \sum_{p=0}^{\infty} a_{2p} x^{2p} = \sum_{p=0}^{\infty} \frac{(-1)^p p!}{2p!} a_0 \cdot x^{2p} \\ &= a_0 \sum_{p=0}^{\infty} \frac{(-1)^p p!}{2p!} x^{2p} \end{aligned}$$

For the second case set $a_0 = 0$ then $a_{2p} = 0$ and $a_{2p+1} = \frac{(-1)^p}{p!} a_1$.

which yields:

$$\begin{aligned} f(x) &= \sum_{p=0}^{\infty} a_{2p+1} x^{2p+1} = \sum_{p=0}^{\infty} \frac{(-1)^p}{p!} a_1 x^{2p+1} \\ &= a_1 x \sum_{p=0}^{\infty} \frac{(-1)^p}{p!} x^{2p} \\ &= a_1 x \sum_{p=0}^{\infty} \frac{(-x^2)^p}{p!} = a_1 x e^{-x^2} \end{aligned}$$

Chapter 7 - Problem 7.4

Substitute back in the two proposed independent solutions $x = z - a$, the general solution of

$$\frac{d^2 f}{dz^2} + z(z-a) \frac{df}{dz} + 4f = 0$$

is therefore

$$f(z, \alpha) = A(z-a) e^{-(z-a)^2} + B \sum_{m=0}^{\infty} \frac{(-4)^m m!}{2m!} (z-a)^{2m}$$