

# Small-molecule fluorophores and fluorescent probes for bioimaging

Takuya Terai · Tetsuo Nagano

Received: 7 August 2012 / Revised: 31 January 2013 / Accepted: 31 January 2013 / Published online: 15 February 2013  
© Springer-Verlag Berlin Heidelberg 2013

**Abstract** Fluorescent compounds based on synthetic small molecules are powerful tools to visualize biological events in living cells and organisms. Ever since the discovery of organic fluorescent compounds in the late nineteenth century, efforts have been made to “see” the behaviors of specific biomolecules in living systems by using these dyes as labels. Also, following the development of fluorescent  $\text{Ca}^{2+}$  indicators in the 1980s, many fluorescent probes or biosensors, which are defined as molecules that show a change in fluorescence properties in the presence of their target molecule, have been reported and applied in biological research. Today, a variety of probes are available that target metal ions, pH, enzyme activities, and signaling molecules. In this review, we first consider the history of organic fluorescent molecules and discuss their utility for labeling biomolecules and staining cells. Then, we review recent progress in small-molecule fluorescent probes for metal ions and reactive oxygen species, focusing on representative work in each category. Finally, we briefly discuss attempts to create novel kinds of probes, including hybrids of small molecules and genetically encoded proteins, with the potential to overcome some of the limitations of current probes.

**Keywords** Fluorescence · Imaging ·  $\text{Ca}^{2+}$  signaling · Reactive oxygen species · Protein labeling

## Introduction

It is said that humans rely on vision for approximately 80 % of all the information they receive from the external world.

Hence, as the proverb “seeing is believing” suggests, an appeal to the eye should be most effective and informative even in the world of science. Unfortunately, however, our body and cells generally do not supply enough visual information detectable with the naked eye to allow us to understand specific biological events, such as cell signaling and protein expression. Therefore, tools that allow us to “see into the body” or “see into cells” are essential not only for basic biology research, but also for the diagnosis and treatment of diseases. Suitable methods must meet several conditions. First, the technique should be sensitive enough to visualize biological compounds at physiological concentrations, which in most cases range from nanomolars to micromolars. Second, the method should have sufficient spatial and temporal resolution to analyze dynamic cell signaling processes. Third, the method should not be invasive, i.e., detection should be possible from outside of the samples. Last, but not the least, it should rely on instruments that are not too special or expensive. At least for cells and dissected tissues, fluorescence imaging is an ideal methodology that satisfies all these requirements, compared with other technologies based on radioactivity, bioluminescence, electromagnetism, and electrochemistry. This is the reason why fluorescence live imaging is coming into widespread use in cell biology. Another advantage of the technology that we should emphasize here is that the fluorescence signal of a molecule can be drastically modulated by design, so probes that exhibit *activation* of the signal, not just *accumulation* of the compound, can be exploited, as we will describe later.

Needless to say, to perform fluorescence imaging, fluorescent molecules must be present in the sample. Although some researchers have used internal or endogenous fluorescent molecules in cells [11,12,38,122,133], including tryptophan, NAD(P)H, and flavins, in most cases, external fluorophores are added to the samples, either chemically or genetically. Before the middle of the 1990s, when fusion of green fluorescent protein [131] was introduced as a labeling

T. Terai · T. Nagano (✉)  
Graduate School of Pharmaceutical Sciences,  
The University of Tokyo, Tokyo, Japan  
e-mail: tlong@mol.f.u-tokyo.ac.jp  
URL: <http://www.f.u-tokyo.ac.jp/~tlong/en/>

tool for proteins in general [19,145], organic fluorescent molecules were the only choices for fluorophores. Although fluorescent proteins are currently widely used, small-molecule fluorescent compounds are still important as both labeling agents and activatable sensors because (1) they are applicable to any sample, including the human body, (2) they are relatively inexpensive and easier to handle, and (3) they can generally provide high signal-to-noise ratios (up to over 1,000) as a result of ingenious chemical design (vide infra).

Therefore, in this review, we focus on organic small-molecule fluorophores and their use as labels and sensors for bioimaging. Probes using fluorescent proteins [26,109,165], quantum dots [98,134], lanthanide ions [13,90,138], and other new platforms are outside the scope of this review. Also, due to space limitation, we describe only representative small-molecule probes here. Various monographs and review articles provide detailed coverage of fluorescence spectroscopy [58,83], fluorophores [47,51,96,97], and fluorescent probes [63,64,77,139,160] for readers who require further information.


### Small-molecule fluorophores

Although phenomena ascribed to fluorescence had been observed for water infusion from a plant [111] and for some kinds of rock in the early modern period, the first fluorescent organic molecule to be specifically identified was

quinine sulfate (Fig. 1a), which was reported by Sir John Herschel in 1845 [54]. Strictly speaking, this is not quite correct because the word “fluorescence” was not coined until 1852, when George Stokes published a long paper on the subject [136]. Since the discovery of quinine sulfate, which emits blue fluorescence when irradiated with ultraviolet (UV) light, many other organic compounds have been found or created to fluoresce in a variety of colors. Fluorescein (Fig. 1b) [4] and rhodamine (Fig. 1c) [18] derivatives, which were first reported as early as the late nineteenth century, are especially important, as they are among the representative platforms currently employed in fluorescent labels and probes used for bioimaging. BODIPY dyes (Fig. 1d) [96] and cyanine dyes (Fig. 1e) [97] are also often used for this purpose.

Despite this long history, structural modifications of organic fluorescent dyes still remain to be explored. For example, fluorescein derivatives in which the carboxyl group is replaced with other functional groups, termed TokyoGreens (TGs), were not created until the twenty-first century (Fig. 1b) [149]. Bright and photostable dyes with red to near-infrared (NIR) fluorescence are especially in demand for in vivo imaging in animals and humans, owing to the transparency of tissues in this wavelength region [97,157]. This issue has been addressed by the development of new dyes with additional ring systems [147,163] and by adding heteroatoms [36,79] to existing fluorophores, as well as by the modification of cyanine dyes [127]. For example, our group has replaced the oxygen atom of TGs with a

**Fig. 1** Chemical structures of representative organic fluorescent molecules. **a** Quinine sulfate, **b** fluorescein and 2-Me TG [149], **c** rhodamine B, **d** 1,3,5,7-tetramethyl-8-phenyl-BODIPY, **e** indocyanine green, **f** 2-Me TM [36], and **g** a photochromic rhodamine [41]. The color of the molecules roughly represents the emission color



dimethylsilyl group to obtain novel fluorescent molecules with red emission, TokyoMagentas (TMs; Fig. 1f) [36]. Compared to fluorescein, 2-Me TM has absorption and emission maxima at around 90 nm longer wavelength. Interestingly, the dye exhibited a large hypsochromic shift of the absorption spectrum upon protonation at weakly acidic pH. Another class of fluorescent dyes that has recently attracted attention is those suitable for two-photon imaging [73,82,99] and for super-resolution imaging. Super-resolution imaging is a collective term that refers to methodologies used to overcome the diffraction limit of conventional light microscopy [59]. Fluorescent molecules that undergo reversible photoswitching [27,41,53] or irreversible photoactivation [8,89] are required for some of these imaging technologies, including stochastic optical reconstruction microscopy (STORM) [125] and photoactivated localization microscopy (PALM) [10]. For instance, Stefan Hell and others reported a photochromic rhodamine derivative whose fluorescence is turned on after UV irradiation and turned off again by thermal relaxation (Fig. 1g) [41].

Due to the difficulty of predicting the photophysical properties of molecules under physiological conditions a priori, most of the novel fluorescent molecules were discovered serendipitously [4,18] or by screening multiple compounds [75,127]. With the aim of accelerating rational and efficient development, many efforts to explain [86,115,123] and predict [87] the optical properties of fluorescent molecules by quantum calculation have been made in recent years, with partial success. It may eventually be possible to combine organic and computational chemistry to obtain new fluorescent scaffolds with superior properties to classical ones.

### Fluorescence imaging with labeled molecules and staining agents


A few decades after the initial development of organic fluorescent molecules, fluorescence microscopy was invented and

applied to biological research. Fluorescence microscopes were first reported by Reichert and Heimstädt in 1911 and by Lehmann in 1913, as an outgrowth of Köhler's UV microscope [37]. They used a carbon arc with appropriate filters as a source of UV light and observed the transmitted fluorescence. The first biological samples to be observed included plants, bacteria, and animal tissues that spontaneously fluoresce. Meanwhile, von Prowazek studied the binding of dyes to cells in order to make them fluorescent in 1914 [152]. This should be regarded as the first attempt to achieve fluorescence staining of cells. Another important development was made in 1942 by Coons, who covalently labeled antibodies with fluorescein isocyanate and used them for tissue imaging with a fluorescence microscope [28]. This was the beginning of immunofluorescence, which is still widely used today.

Since those early works, a wide range of fluorescent labeling agents and cell staining agents has been developed and applied in biological and medical studies. For example, fluorescent dyes that selectively localize and stain a cellular target organelle, such as mitochondria, lysosomes, endoplasmic reticulum, or Golgi apparatus, are now commercially available [63]. Perhaps the most widely used are dyes for nuclear staining, such as 4',6-diamidino-2-phenylindole (DAPI) [67] and Hoechst 33258 [85], which emit strong fluorescence when bound to DNA. More recent examples of organelle-targeted dyes include a series of styryl fluorophores reported by Young-Tae Chang [124] and a mitochondrial surface-specific fluorescent molecule reported by Motonari Uesugi [68]. Notably, these dyes were developed by combinatorial approaches as described in the succeeding sections.

In the last decade, *in vivo* imaging of tumors and other lesions with antibodies [6,117] and peptides [21,69] labeled with fluorescent dyes has attracted attention. For these applications, our laboratory has developed Si-substituted rhodamine-based NIR fluorescent labeling agents, including SiR700 [78]. When used to label tumor-specific antibodies, they are superior to conventional cyanine-based agents in terms of photostability. As shown in Fig. 2, we demonstrated

**Fig. 2** *In vivo* tumor imaging with SiR700-labeled anti-tenascin-C antibody [78]. **a** Representative illustration of the imaging agent. **b** Fluorescence image of a mouse xenograft tumor model (prepared with HKBMM cells) before antibody injection. The white circle indicates the position of the implanted tumor. **c** Image of the same mouse 24 h after antibody injection. The images were taken on a Pearl Impulse Imager with a 700-nm channel



that such a dye–antibody conjugate could clearly visualize mouse tumors *in vivo*. Other remarkable recent reports on fluorescence imaging of specific tissues *in vivo* include nerve detection by Quyen Nguyen [159] and artery labeling by Prakash Kara [128].


### Activatable fluorescent probes

As mentioned earlier, one of the advantages of fluorescence imaging is that the signal of a molecule can be drastically modulated. In this review, those molecules whose own optical properties are altered in response to the target chemical species or environments are defined as “fluorescent probes” (Fig. 3a). There are several rationales for the design of fluorescent probes, including photoinduced electron transfer (PeT) [16,146,149], intramolecular (or internal) charge transfer [31], Förster (or fluorescence) resonance energy transfer (FRET) [42,109], and intramolecular spirocyclization [24,65]. Schematic representations of PeT-based and FRET-based probes are shown in Fig. 3b, c, respectively. PeT is the transfer of an electron to (or from) the fluorophore at the excited state from (or to) an electron donor (or acceptor) moiety that is introduced near the fluorophore. If this process occurs, the excited state of the fluorescent molecule is quenched, resulting in low fluorescence intensity. In the case of activatable fluorescent probes, PeT is cancelled by chemical reaction of the probe with the target molecule, with consequent recovery of strong fluorescence (Fig. 3b). The kinetics and thermodynamics of PeT were formulated by Rudolph Marcus [101] and Dieter Rehm

and Albert Weller [121], respectively. Early work on PeT-based fluorescent probes included the pH probes developed by A. Prasanna de Silva and others [31]. On the other hand, FRET is a nonradiative transfer of energy between two fluorescent molecules, one being the donor and the other being the acceptor. If this occurs, fluorescence of the donor molecule is weakened, and instead, fluorescence from the acceptor molecule is observed even though the donor is irradiated. For efficient FRET, the spectral overlap, as well as the distance, between the donor and acceptor chromophore is important [109]. When, for example, the linker between the two fluorophores is cleaved by a proteolytic enzyme, FRET is cancelled and the donor fluorescence is restored (Fig. 3c). In contrast to PeT-based probes, which only show a change of fluorescence intensity, FRET-based probes show a change in the shape of the excitation/emission spectra in response to the target molecule. As is well known, this feature allows for ratiometric measurement, which is highly advantageous for precise quantification of the analyte. Representative FRET-based small-molecule probes include a  $\beta$ -lactamase probe developed by Roger Tsien’s group [166].

Because the number as well as the scope of fluorescent probes based on small molecules is continuously expanding, it is not possible to cover all of them in this short article. Instead, we will briefly describe probes for the following two categories: metal ions and reactive oxygen (nitrogen) species. Some important areas that are not reviewed herein include slow and fast membrane potential probes [39,48,49,106,153], enzyme activity probes [7,120,126], and pH indicators [112,148,158].

**Fig. 3** Development of fluorescent probes. **a** Schematic representation of fluorescent probes, **b** design of probes based on PeT, and **c** design of probes based on FRET. For details, see the references mentioned in the text



## Probes for metal ions

### Probes for $\text{Ca}^{2+}$

The classical, and still probably the most common, target of fluorescent probes is the  $\text{Ca}^{2+}$  ion. Needless to say,  $\text{Ca}^{2+}$  as a second messenger plays pivotal roles in many types of cells, including neurons [15], eggs [34], and myocytes [9]. In 1980s, Roger Tsien and coworkers first reported a series of selective fluorescent probes for  $\text{Ca}^{2+}$ , including quin2 [144], fura-2 [50], fluo-3 [107], and rhod-2 [107] (Fig. 4a, b). These epoch-making probes demonstrated the power of fluorescence live cell imaging to biologists in every field. Even today, some of those dyes are still used to provide novel scientific insights. Importantly, the probes, together with the newer ones (vide infra), share the same chelator moiety called 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid (BAPTA). BAPTA was developed by Tsien as a novel chelator with improved properties over the previously known ethylene glycol-bis(2-aminoethyl ether)-*N,N,N',N'*-tetraacetic acid (EGTA; Fig. 4c) [144]. Although EGTA itself had high selectivity for  $\text{Ca}^{2+}$  over  $\text{Mg}^{2+}$ , only a small fraction of EGTA exists in an unprotonated state at physiological pH due to the high  $\text{pK}_a$  of the amine moieties (8.96 and 9.58). To overcome this problem, Tsien improved the chelator by replacing the methylene groups connecting nitrogen and oxygen of EGTA with aromatic rings. This modification lowered the  $\text{pK}_a$  values of the BAPTA amines to 6.36 and 5.47 and allowed  $\text{Ca}^{2+}$  to be well-chelated in the physiological pH range. Also, it made the sensor relatively insensitive to small fluctuations in pH.

After 1990, novel fluorescent probes for  $\text{Ca}^{2+}$  such as fluo-4 [44], Oregon Green 488 BAPTA-1 [95], and a two-photon probe [74] were developed and applied for imaging cells and tissues. For more information, see other references [63,137]. In 2011, two NIR fluorescent probes with large


activation ratios in the presence of  $\text{Ca}^{2+}$  were reported independently by our group [35] (Fig. 4d) and Koji Suzuki's group [104]. These NIR probes should be useful not only for monitoring  $\text{Ca}^{2+}$  fluctuation in tissues, but also for multicolor imaging in combination with visible-range fluorescent probes.

### Probes for $\text{K}^+$ and $\text{Na}^+$

$\text{K}^+$  is the most abundant metal ion inside cells and plays essential roles in many physiological events, such as cardiac and neuronal excitability, cellular ionic homeostasis, and cell proliferation [130]. The first fluorescent probe for  $\text{K}^+$ , potassium-binding benzofuran isophthalate (PBFI) (Fig. 4e), incorporating an 18-crown-6 base chelator, was reported by Tsien during the course of studies aimed at a  $\text{Na}^+$ -selective probe [108]. Although PBFI is commercially available [63], its biological applications are relatively limited, in contrast to the case of the  $\text{Ca}^{2+}$  probes, due to the low fluorescence quantum yield and poor selectivity for  $\text{K}^+$  over  $\text{Na}^+$ . To address this problem, a novel  $\text{K}^+$ -selective cryptand-based chelator, 2-triazacryptand[2,2,3]-1-(2-methoxyethoxy)benzene (TAC), was recently synthesized by Huarui He [52], and several fluorescent probes using TAC as a recognition moiety have been reported [57,114,119], including TAC-Red by Alan Verkman (Fig. 4f) [119]. These probes possess improved selectivity for  $\text{K}^+$  and can monitor changes of  $\text{K}^+$  concentration outside the cell, where  $\text{Na}^+$  is present in excess.

$\text{Na}^+$  is the most abundant metal ion in extracellular fluids. Flux of  $\text{Na}^+$  into cells induces membrane depolarization, and abnormality of  $\text{Na}^+$  channels is associated with human diseases [5,162]. As mentioned previously, the first fluorescent probe for  $\text{Na}^+$  in living cells, sodium-binding benzofuran isophthalate (SBFI), was reported by Tsien in 1989 [108]. Although it is relatively selective for  $\text{Na}^+$  and is suitable for ratiometric imaging, its excitation wavelengths are in the UV

**Fig. 4** Chemical structures of fluorescent probes and chelators for metal ions: **a** fura-2 [50], **b** fluo-3 [107], **c** EGTA, **d** CaSiR-1 [35], **e** PBFI [108], and **f** TAC-Red [119]





region and its brightness is limited. As improved versions, CoroNa Green [105] and CoroNa Red [63] with visible excitation wavelengths have been developed and commercialized. However, CoroNa Green has the drawback of fast leakage from live cells [105]. Very recently, Christophe Lamy solved this problem by encapsulating the probe in a dendrimer to prolong its intracellular retention while retaining the response to  $\text{Na}^+$  [84]. Though the conjugate has to be delivered into cells by microinjection or electroporation due to its membrane impermeability, improved retention will be beneficial for imaging experiments.

#### Probes for other metal ions

Many metal ions other than  $\text{Ca}^{2+}$ ,  $\text{Na}^+$ , and  $\text{K}^+$  can currently be visualized by the use of specific fluorescent probes [24,33,143]. Among them, perhaps the most popular target is  $\text{Zn}^{2+}$ . Zinc is an essential metal for life and is the second most abundant transition metal in humans. The biological functions of  $\text{Zn}^{2+}$  have been well-established for the protein-bound form, but the functions of free or loosely bound (labile, mobile)  $\text{Zn}^{2+}$  are less clear [72]. In order to address this issue, many fluorescent probes for  $\text{Zn}^{2+}$  have been developed, including 6-methoxy-(8-*p*-toluenesulfonamido)quinoline (TSQ) [43], zinpyrs (reported by Stephen Lippard's group) [14], and ZnAFs (reported from our group) [56]. These probes have been reviewed in detail elsewhere [62,72,116].

Copper is also an essential metal for life, mainly due to its redox activity, and it is a popular target of fluorescent probes [24,33]. It is known that the brain needs quite high levels of copper, but at the same time, mishandling of neuronal copper stores is associated with various pathological conditions, including Menkes and Wilson's diseases [17]. In 2011, Christopher Chang developed a novel BODIPY-based fluorescent probe for  $\text{Cu}^+$  and discovered that pools of copper in neurons undergo a dynamic redistribution that is dependent on calcium signaling [32].

#### Probes for reactive oxygen (nitrogen) species

Reactive oxygen species (ROS) is a collective term for activated oxygen species, including superoxide ( $\text{O}_2^{\cdot-}$ ), hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), hydroxyl radical ( $\cdot\text{OH}$ ), and others (Fig. 5a). They are important signaling molecules, which regulate a wide range of physiological functions. However, overproduction of ROS results in oxidative stress and is involved in the pathogenesis of many diseases [29]. To analyze the dynamic functions of ROS in living systems *in situ*, many fluorescent probes based on small molecules have been developed. The classical probes are chemically reduced forms of fluorescent dyes, such as 2',7'-dichlorodihydrofluorescein (DCFH;

Fig. 5b) [25] and dihydrorhodamine 123 [81], which are oxidized by ROS to afford strongly fluorescent products. Although they are widely used in cell imaging, auto-oxidation and lack of specificity among ROS are major drawbacks. To address these problems, probes that fluoresce only in the presence of specific ROS have recently been developed, some of which are described in the succeeding section. More comprehensive reviews of previously reported ROS probes are available [45,113,156].

Reactive chemical species analogous to ROS but containing nitrogen are known as reactive nitrogen species (RNS; Fig. 5a). The most representative RNS is nitric oxide, which is involved in many physiological and pathological processes [76]. In the last two decades, many fluorescent probes for RNS have been developed by us and others [46,91,113], but they will not be reviewed here.


#### Probes for hydrogen peroxide

$\text{H}_2\text{O}_2$  is a relatively mild ROS and has attracted intense interest recently because of its involvement in signal transduction by reversible oxidation of proteins, in addition to its established cytotoxic effects [150]. Most of the reported fluorescent probes for  $\text{H}_2\text{O}_2$  have esters of aryl boronates as the recognition moieties; these are converted to phenols by  $\text{H}_2\text{O}_2$  with a concomitant increase of fluorescence intensity. To our knowledge, this type of probe was first developed by Lee-Chiang Lo in 2003 [93] and vigorously improved by Christopher Chang and others [92]. A representative example, PF1, is shown in Fig. 5c [20]. Probes based on other chemistries, including ROS-mediated cleavage of benzenesulfonate groups, have been developed as well [100]. Recently, our group reported a novel probe incorporating benzil as a reactive moiety, NBzF [1]. Because benzil has a high reduction potential, the fluorescence of the probe was significantly quenched by PeT from the fluorophore to the benzil moiety, and the PeT was cancelled selectively and rapidly by reaction with  $\text{H}_2\text{O}_2$  (Fig. 5d). These specific probes are expected to be useful in understanding the roles of  $\text{H}_2\text{O}_2$  in living systems.

#### Probes for hypochlorous acid

Hypochlorous acid (HOCl) is a powerful oxidant that is produced by myeloperoxidase from  $\text{H}_2\text{O}_2$  and  $\text{Cl}^-$  in certain types of cells, including activated phagocytes. Under physiological conditions, approximately half of HOCl is present as the hypochlorite anion ( $\text{OCl}^-$ ) form. It is involved in killing microorganisms and also in many brain diseases [161]. The first practical fluorescent probe for HOCl, HySOx, was developed by our group in 2007 [70], using rhodamine spirocyclization as the rationale (Fig. 5e). With this probe, we could visualize the generation of HOCl in

**Fig. 5** ROS and fluorescent probes used to detect them.  
**a** Schematic relationship of various ROS (red) and RNS (blue). Chemical structures and reactions of **b** DCFH [25], **c** PF1 [20], **d** NBzF [1], and **e** HySOx [70]



phagosomes in real time. Following this work, many specific probes for HOCl have been reported based on different chemical reactions [22,23,129] and applied for bioimaging.

### Novel approaches for the future

In the last part of this review, we briefly describe two new approaches that can potentially overcome the limitations of

current small molecule-based fluorescent probes: combinatorial probe development and the use of protein/peptide tag technology.

### Combinatorial development of probes

As described previously, small molecule-based fluorescent probes can be developed rationally by incorporating specific reactive moieties into fluorescent molecules. However, in

some cases, it is hardly possible to design a suitable chemical reaction or a recognition moiety for the target. For example, if one wants to detect cells of a specific kind, it would be very difficult to decide what target to select. To overcome this limitation, a combinatorial or diversity-oriented approach is gaining increasing attention [68,124,151]. The power of this strategy was fully demonstrated in a paper by Young-Tae Chang and others, who screened a rosamine library they had previously reported [2] and found a fluorescent compound that selectively stains pluripotent stem cells [60]. His group also prepared a BODIPY library and obtained fluorescent probes for dopamine [88] and fructose [164]. A similar approach was reported by Seung Bum Park and others [75]; they synthesized a library of compounds and screened them to find novel fluorescent molecules with excellent photophysical properties. The works on fluorogenic enzyme substrates by Jean-Louis Reymond [154] and Dalibor Sames [40] may also be classified into this category.

#### Use of protein/peptide tags


Although many small-molecule fluorescent probes provide a good activation ratio with high target specificity in living cells, undesired localization and retention properties of the probe often become major concerns. For example, some fluorescein-based probes are readily removed from cells [61,105], while rhodamine-based probes tend to localize in mitochondria [80]. Chemical modifications of probes, such as conjugation of chelators [61] or targeting moieties [3], are

applicable in some cases, but the need for more general methodologies to control probe localization at will is clear. To address this issue, the use of tag technologies in combination with small-molecule fluorescent probes is becoming popular. Here, a tag technology corresponds to a method to couple small functional molecules (fluorescent dyes, in most cases) to genetically encoded proteins/peptides that are designed to specifically bind, either covalently or noncovalently, to molecules with characteristic chemical motifs. So far, a few tag technologies have been established, including FLAsH/ReAsH system [103], SNAP-tag [71], and HaloTag [94]. In addition, several new technologies are being developed [55,110,118].

A representative combination probe was developed by Tsien's group in 2007; they created a fluorescent  $\text{Ca}^{2+}$  probe with a biarsenical moiety that selectively binds to tetracysteine-tagged membrane proteins (Fig. 6a) and used it to monitor local  $\text{Ca}^{2+}$  dynamics [141]. Similar strategies have been reported by other groups to direct  $\text{Ca}^{2+}$  probes [66,102], a  $\text{Zn}^{2+}$  probe [140] (Fig. 6b), and a  $\text{H}_2\text{O}_2$  probe [135] to selected organelles in order to measure local changes of the analyte.

Moreover, the scope for combining proteins with small-molecule fluorophores is not limited to these areas. For example, researchers have used solvent-sensitive (alternatively called environment-sensitive) fluorophores to monitor enzyme activities [132] or conformational changes [142] of proteins or peptides, and this approach has also been applied to monitor ion concentration [155], though these applications are beyond the scope of the present review.

**Fig. 6** Fluorescent probes with labeling moieties directed to specific tags. **a** A biarsenical  $\text{Ca}^{2+}$  probe that binds to tetracysteine-conjugated proteins [141]. **b** A  $\text{Zn}^{2+}$  probe that labels SNAP-tag fusion proteins [140]





## Concluding remarks

It has been about 150 years since the discovery of organic small-molecule fluorophores. Throughout this time, numerous fluorescent molecules have been synthesized and applied in imaging of biomolecules, cells, and organisms. Introduction of novel imaging platforms, including in vivo NIR imaging, super-resolution imaging, and multiphoton imaging, has always been accompanied by the development/discovery of appropriate organic fluorescent labeling dyes that play an essential role in enabling the technologies. Importantly, the fluorescence properties of organic molecules can be rationally regulated by means of several mechanisms, allowing chemists to develop activatable fluorescent probes. At present, many fluorescent probes are available, directed to a range of targets. Without doubt, they have made significant contributions to our understanding of dynamic and complicated processes in biological systems. As mentioned previously, recently developed novel approaches should complement the conventional methodology for probes based on small-molecule fluorophores. These and other innovations are expected to spur the development of biological sciences and medicine.

**Acknowledgments** We thank Dr. Toru Komatsu for the helpful comments on the manuscript. We are also grateful for the financial support from the Ministry of Education, Culture, Sports, Science and Technology of Japan (grant no. 22000006 to T.N. and T.T. and grant no. 23651231 to T.T.). T.T. was also supported by the Cosmetology Research Foundation and Mochida Memorial Foundation for Medical and Pharmaceutical Research, Japan.

## References

- Abo M, Urano Y, Hanaoka K, Terai T, Komatsu T, Nagano T (2011) Development of a highly sensitive fluorescence probe for hydrogen peroxide. *J Am Chem Soc* 133:10629–10637
- Ahn Y-H, Lee J-S, Chang Y-T (2007) Combinatorial rosamine library and application to in vivo glutathione probe. *J Am Chem Soc* 129:4510–4511
- Baek NY, Heo CH, Lim CS, Masanta G, Cho BR, Kim HM (2012) A highly sensitive two-photon fluorescent probe for mitochondrial zinc ions in living tissue. *Chem Commun* 48:4546–4548
- Baeyer A (1871) Ueber eine neue Klasse von Farbstoffen. *Ber Dtsch Chem Ges* 4:555–558
- Baker MD, Wood JN (2001) Involvement of Na<sup>+</sup> channels in pain pathways. *Trends Pharmacol Sci* 22:27–31
- Ballou B, Ernst LA, Waggoner AS (2005) Fluorescence imaging of tumors in vivo. *Curr Med Chem* 12:795–805
- Baruch A, Jeffery DA, Bogoy M (2004) Enzyme activity—it's all about image. *Trends Cell Biol* 14:29–35
- Belov VN, Wurm CA, Boyarskiy VP, Jakobs S, Hell SW (2010) Rhodamines NN: a novel class of caged fluorescent dyes. *Angew Chem Int Ed* 49:3520–3523
- Bers DM (2008) Calcium cycling and signaling in cardiac myocytes. *Annu Rev Physiol* 70:23–49
- Betzig E, Patterson GH, Sougrat R, Lindwasser OW, Olenych S, Bonifacino JS, Davidson MW, Lippincott-Schwartz J, Hess HF (2006) Imaging intracellular fluorescent proteins at nanometer resolution. *Science* 313:1642–1645
- Brancaleon L, Durkin AJ, Tu JH, Menaker G, Fallon GD, Kollias N (2001) In vivo fluorescence spectroscopy of nonmelanoma skin cancer. *Photochem Photobiol* 73:178–183
- Brookner CK, Follen M, Boiko I, Galvan J, Thomsen S, Malpica A, Suzuki S, Lotan R, Richards-Kortum R (2000) Autofluorescence patterns in short-term cultures of normal cervical tissue. *Photochem Photobiol* 71:730–736
- Bünzli J-CG (2010) Lanthanide luminescence for biomedical analyses and imaging. *Chem Rev* 110:2729–2755
- Burdette SC, Walkup GK, Springler B, Tsien RY, Lippard SJ (2001) Fluorescent sensors for Zn<sup>2+</sup> based on fluorescein platform: synthesis, properties and intracellular distribution. *J Am Chem Soc* 123:7831–7841
- Burgoyne RD (2007) Neuronal calcium sensor proteins: generating diversity in neuronal Ca<sup>2+</sup> signaling. *Nat Rev Neurosci* 8:182–193
- Callan JF, de Silva AP, Magri DC (2005) Luminescent sensors and switches in the early 21st century. *Tetrahedron* 61:8551–8588
- Camakaris J, Voskoboinik I, Mercer JF (1999) Molecular mechanisms of copper homeostasis. *Biochem Biophys Res Commun* 261:225–232
- Ceresole M (1888) Production of new red coloring-matter. *US Patent* 377,349
- Chalfie M, Tu Y, Euskirchen G, Ward WW, Prasher DC (1994) Green fluorescent protein as a marker for gene expression. *Science* 263:802–805
- Chang MCY, Pralle A, Isacoff EY, Chang CJ (2004) A selective, cell-permeable optical probe for hydrogen peroxide in living cells. *J Am Chem Soc* 126:15392–15393
- Chen X, Conti PS, Moats RA (2004) In vivo near-infrared fluorescence imaging of integrin  $\alpha_v\beta_3$  in brain tumor xenografts. *Cancer Res* 64:8009–8014
- Chen X, Lee K-A, Ha E-M, Lee KM, Seo YY, Choi HK, Kim HN, Kim MJ, Cho C-S, Lee SY, Lee W-J, Yoon J (2011) A specific and sensitive method for detection of hypochlorous acid for the imaging of microbe-induced HOCl production. *Chem Commun* 47:4373–4375
- Chen S, Lu J, Sun C, Ma H (2010) A highly specific ferrocene-based fluorescent probe for hypochlorous acid and its application to cell imaging. *Analyst* 135:577–582
- Chen X, Pradhan T, Wang F, Kim JS, Yoon J (2012) Fluorescent chemosensors based on spiroring-opening of xanthenes and related derivatives. *Chem Rev* 112:1910–1956
- Chen X, Zhong Z, Xu Z, Chen L, Wang Y (2010) 2',7'-Dichlorodihydrofluorescein as a fluorescent probe for reactive oxygen species measurement: forty years of application and controversy. *Free Radic Res* 44:587–604
- Chudakov DM, Matz MV, Lukyanov S, Lukyanov KA (2010) Fluorescent proteins and their applications in imaging living cells and tissues. *Physiol Rev* 90:1103–1163
- Conley NR, Biteen JS, Moerner WE (2008) Cy3–Cy5 covalent heterodimers for single-molecule photoswitching. *J Phys Chem B* 112:11878–11880
- Coons AH, Creech HJ, Jones RN, Berliner E (1942) The demonstration of pneumococcal antigen in tissues by the use of fluorescent antibody. *J Immunol* 45:159–170
- D'Aurèaux B, Toledano MB (2007) ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis. *Nat Rev Mol Cell Biol* 8:813–824
- de Silva AP, Gunaratne HQN, Gunnlaugsson T, Huxley AJM, McCoy CP, Rademacher JP, Rice TE (1997) Signaling recognition events with fluorescent sensors and switches. *Chem Rev* 97:1515–1566

31. de Silva AP, Rupasinghe RAD (1985) A new class of fluorescent pH indicators based on photo-induced electron transfer. *J Chem Soc Chem Commun* (23):1669–1670
32. Dodani SC, Domaille DW, Nam CI, Miller EW, Finney LA, Vogt S, Chang CJ (2011) Calcium-dependent copper redistributions in neuronal cells revealed by a fluorescent copper sensor and X-ray fluorescence microscopy. *Proc Natl Acad Sci USA* 108:5980–5985
33. Domaille DW, Que EL, Chang CJ (2008) Synthetic fluorescent sensors for studying the cell biology of metals. *Nat Chem Biol* 4:168–175
34. Ducibella T, Fissore R (2008) The roles of  $\text{Ca}^{2+}$ , downstream protein kinases, and oscillatory signaling in regulating fertilization and the activation of development. *Dev Biol* 315:257–279
35. Egawa T, Hanaoka K, Koide Y, Ujita S, Takahashi N, Ikegaya Y, Matsuki N, Terai T, Ueno T, Komatsu T, Nagano T (2011) Development of a far-red to near-infrared fluorescence probe for calcium ion and its application to multicolor neuronal imaging. *J Am Chem Soc* 133:14157–14159
36. Egawa T, Koide Y, Hanaoka K, Komatsu T, Terai T, Nagano T (2011) Development of a fluorescein analogue, TokyoMagenta, as a novel scaffold for fluorescence probes in red region. *Chem Commun* 47:4162–4164
37. Ellinger P (1940) Fluorescence microscope in biology. *Biol Rev* 15:323–347
38. Eng J, Lynch RM, Balaban RS (1989) Nicotinamide adenine dinucleotide fluorescence spectroscopy and imaging of isolated cardiac myocytes. *Biophys J* 55:621–630
39. Fluhler E, Burnham VG, Loew LM (1985) Spectra, membrane binding, and potentiometric responses of new charge shift probes. *Biochem* 24:5749–5755
40. Foemming MK, Sames D (2007) Harnessing functional plasticity of enzymes: a fluorogenic probe for imaging  $17\alpha$ -HSD10 dehydrogenase, an enzyme involved in Alzheimer's and Parkinson's diseases. *J Am Chem Soc* 129:14518–14522
41. Fölling J, Belov V, Kunetsky R, Medda R, Schönle A, Egner A, Eggeling C, Bossi B, Hell SW (2007) Photochromic rhodamines provide nanoscopy with optical sectioning. *Angew Chem Int Ed* 46:6266–6270
42. Förster T (1959) Transfer mechanism of electronic excitation. *Discuss Faraday Soc* 27:7–17
43. Frederickson CJ, Kasarskis EJ, Ringo D, Frederickson RE (1987) A quinoline fluorescence method for visualizing and assaying the histochemically reactive zinc (bouton zinc) in the brain. *J Neurosci Methods* 20:91–103
44. Gee KR, Brown KA, Chen W-NU, Bishop-Stewart J, Gray D, Johnson I (2000) Chemical and physiological characterization of fluo-4  $\text{Ca}^{2+}$ -indicator dyes. *Cell Calcium* 27:97–106
45. Gomes A, Fernandes E, Lima JLFC (2005) Fluorescence probes used for detection of reactive oxygen species. *J Biochem Biophys Methods* 65:45–80
46. Gomes A, Fernandes E, Lima JLFC (2006) Use of fluorescence probes for detection of reactive nitrogen species: a review. *J Fluoresc* 16:119–139
47. Gonçalves MST (2009) Fluorescent labeling of biomolecules with organic probes. *Chem Rev* 109:190–212
48. González JE, Tsien RY (1997) Improved indicators of cell membrane potential that use fluorescence resonance energy transfer. *Chem Biol* 4:269–277
49. Grinvald A, Fine A, Farber IC, Hildesheim R (1983) Fluorescence monitoring of electrical responses from small neurons and their processes. *Biophys J* 42:195–198
50. Grynkiewicz G, Poenie M, Tsien RY (1985) A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J Biol Chem* 260:3440–3450
51. Hawe A, Sutter M, Jiskoot W (2007) Extrinsic fluorescent dyes as tools for protein characterization. *Pharm Res* 25:1487–1499
52. He H, Mortellaro MA, Leiner MJP, Fraatz RJ, Tusa JK (2003) A fluorescent sensor with high selectivity and sensitivity for potassium in water. *J Am Chem Soc* 125:1468–1469
53. Heilemann M, van de Linde S, Mukherjee A, Sauer M (2009) Super-resolution imaging with small organic fluorophores. *Angew Chem Int Ed* 48:6903–6908
54. Herschel Sir JFW (1845) On a case of superficial colour presented by a homogeneous liquid internally colorless. *Phil Trans Roy Soc London* 135:143–145
55. Hirabayashi K, Hanaoka K, Shimonishi M, Terai T, Komatsu T, Ueno T, Nagano T (2011) Selective two-step labeling of proteins with an off/on fluorescent probe. *Chem Eur J* 17:14763–14771
56. Hirano T, Kikuchi K, Urano Y, Nagano T (2002) Improvement and biological applications of fluorescent probes for zinc, ZnAFs. *J Am Chem Soc* 124:6555–6562
57. Hirata T, Terai T, Komatsu T, Hanaoka K, Nagano T (2011) Development of a potassium ion-selective fluorescent sensor based on 3-styrylated BODIPY. *Bioorg Med Chem Lett* 21:6090–6093
58. Hof M, Hutterer R, Fidler V (eds) (2010) Fluorescence spectroscopy in biology: advanced methods and their applications to membranes, proteins, DNA, and cells. Springer, Berlin
59. Huang B, Bates M, Zhuang X (2009) Super-resolution fluorescence microscopy. *Annu Rev Biochem* 78:993–1016
60. Im C-N, Kang N-Y, Ha H-H, Bi X, Lee JJ, Park S-J, Lee SY, Vendrell M, Kim YK, Lee J-S, Li J, Y-H FB, Ng H-H, Yun S-W, Chang Y-T (2010) A fluorescent rosamine compound selectively stains pluripotent stem cells. *Angew Chem Int Ed* 49:7497–7500
61. Izumi S, Urano Y, Hanaoka K, Terai T, Nagano T (2009) A simple and effective strategy to increase the sensitivity of fluorescence probes in living cells. *J Am Chem Soc* 131:10189–10200
62. Jiang P, Guo Z (2004) Fluorescent detection of zinc in biological systems: recent development on the design of chemosensors and biosensors. *Coord Chem Rev* 248:205–229
63. Johnson I, Spence MTZ (eds) (2010) The molecular probes handbook, 11th edn. Life Technologies, Carlsbad
64. Juskowiak B (2011) Nucleic acid-based fluorescent probes and their analytical potential. *Anal Bioanal Chem* 399:3157–3176
65. Kamiya M, Asanuma D, Kuranaga E, Takeishi A, Sakabe M, Miura M, Nagano T, Urano Y (2011)  $\beta$ -Galactosidase fluorescence probe with improved cellular accumulation based on a spirocyclized rhodol scaffold. *J Am Chem Soc* 133:12960–12963
66. Kamiya M, Johnsson K (2010) Localizable and highly sensitive calcium indicator based on a BODIPY fluorophore. *Anal Chem* 82:6472–6479
67. Kapuscinski J (1995) DAPI: a DNA-specific fluorescent probe. *Biotech Histochem* 70:220–233
68. Kawazoe Y, Shimogawa H, Sato A, Uesugi M (2011) A mitochondrial surface-specific fluorescent probe activated by bioconversion. *Angew Chem Int Ed* 50:5478–5481
69. Ke S, Wen X, Gurfinkel M, Chamsangavej C, Wallace S, Seavick-Muraca EM, Li C (2003) Near-infrared optical imaging of epidermal growth factor receptor in breast cancer xenografts. *Cancer Res* 63:7870–7875
70. Kenmoku S, Urano Y, Kojima H, Nagano T (2007) Development of a highly specific rhodamine-based fluorescence probe for hypochlorous acid and its application to real-time imaging of phagocytosis. *J Am Chem Soc* 129:7313–7318
71. Keppler A, Gendrezig S, Gronemeyer T, Pick H, Vogel H, Johnsson K (2003) A general method for the covalent labeling of fusion proteins with small molecules in vivo. *Nat Biotechnol* 21:86–89

72. Kikuchi K, Komatsu K, Nagano T (2004) Zinc sensing for cellular application. *Curr Opin Chem Biol* 8:182–191
73. Kim HM, Cho BR (2009) Two-photon probes for intracellular free metal ions, acidic vesicles, and lipid rafts in live tissues. *Acc Chem Res* 42:863–872
74. Kim HM, Kim BR, Hong JH, Park J-S, Lee KJ, Cho BR (2007) A two-photon fluorescent probe for calcium waves in living tissue. *Angew Chem Int Ed* 46:7445–7448
75. Kim E, Koh M, Ryu J, Park SB (2008) Combinatorial discovery of full-color-tunable emissive fluorescent probes using a single core skeleton, 1,2-dihydropyrrolo[3,4- $\beta$ ]indolizin-3-one. *J Am Chem Soc* 130:12206–12207
76. Kim PKM, Zamora R, Petrosko P, Billiar TR (2001) The regulatory role of nitric oxide in apoptosis. *Int Immunopharmacol* 1:1421–1441
77. Kiyose K, Kojima H, Nagano T (2008) Functional near-infrared fluorescent probes. *Chem Asian J* 3:506–515
78. Koide Y, Urano Y, Hanaoka K, Piao W, Kusakabe M, Saito N, Terai T, Okabe T, Nagano T (2012) Development of NIR fluorescent dyes based on Si-rhodamine for in vivo imaging. *J Am Chem Soc* 134:5029–5031
79. Koide Y, Urano Y, Hanaoka K, Terai T, Nagano T (2011) Evolution of group 14 rhodamines as platforms for near-infrared fluorescence probes utilizing photoinduced electron transfer. *ACS Chem Biol* 6:600–608
80. Koide Y, Urano Y, Kenmoku S, Kojima H, Nagano T (2007) Design and synthesis of fluorescent probes for selective detection of highly reactive oxygen species in mitochondria of living cells. *J Am Chem Soc* 129:10324–10325
81. Kooy NW, Royall JA, Ischiropoulos H, Beckman JS (1994) Peroxynitrite-mediated oxidation of dihydrorhodamine 123. *Free Radic Biol Med* 16:149–156
82. Krishna TR, Parent M, Werts MHV, Moreaux L, Gmouh S, Charpak S, Caminade A-M, Majoral J-P, Blanchard-Desce M (2006) Water-soluble dendrimeric two-photon tracers for in vivo imaging. *Angew Chem Int Ed* 45:4645–4648
83. Lakowicz JR (2006) Principles of fluorescence spectroscopy, 3rd edn. Springer, New York
84. Lamy CM, Sallin O, Loussert C, Chatton J-Y (2012) Sodium sensing in neurons with a dendrimer-based nanoprobe. *ACS Nano* 6:1176–1187
85. Latt SA, Stetten G (1976) Spectral studies on 33258 Hoechst and related bisbenzimidazole dyes useful for fluorescent detection of deoxyribonucleic acid synthesis. *J Histochem Cytochem* 24:24–33
86. Lazarides T, McCormick TM, Wilson KC, Lee S, McCamant DW, Eisenberg R (2011) Sensitizing the sensitizer: the synthesis and photophysical study of Bodipy-Pt(II)(diimine)(dithiolate) conjugates. *J Am Chem Soc* 133:350–364
87. Le Guennic B, Maurya O, Jacquemin D (2012) Aza-boron-dipyrromethene dyes: TD-DFT benchmarks, spectral analysis and design of original near-IR structures. *Phys Chem Chem Phys* 14:157–164
88. Lee J-S, Kim HK, Feng S, Vendrell M, Chang Y-T (2011) Accelerating fluorescent sensor discovery: unbiased screening of a diversity-oriented BODIPY library. *Chem Commun* 47:2339–2341
89. Lee HD, Lord SJ, Iwanaga S, Zhan K, Xie H, Williams JC, Wang H, Bowman GR, Goley ED, Shapiro L, Twieg RJ, Rao J, Moerner WE (2010) Superresolution imaging of targeted proteins in fixed and living cells using photoactivatable organic fluorophores. *J Am Chem Soc* 132:15099–15101
90. Li F, Yang H, Hu H (2010) Luminescent rare earth complexes as chemosensors and bioimaging probes. In: Huang C (ed) Rare earth coordination chemistry: fundamentals and applications. Wiley (Asia), Singapore, pp 529–570
91. Lim MH, Wong BA, Pitcock WH Jr, Mokshagundam D, Baik M-H, Lippard SJ (2006) Direct nitric oxide detection in aqueous solution by copper(II) fluorescein complexes. *J Am Chem Soc* 128:14364–14373
92. Lippert AR, van de Bittner GC, Chang CJ (2011) Boronate oxidation as a bioorthogonal reaction approach for studying the chemistry of hydrogen peroxide in living systems. *Acc Chem Res* 44:793–804
93. Lo LC, Chu CY (2003) Development of highly selective and sensitive probes for hydrogen peroxide. *Chem Commun* (21):2728–2729
94. Los GV, Encell LP, McDougall MG, Hartzell DD, Karassina N, Zimprich C, Wood MG, Learish R, Ohana RF, Urh M, Simpson D, Mendez J, Zimmerman K, Otto P, Vidugiris G, Zhu J, Darzins A, Klaubert DH, Bulleit RB, Wood KV (2008) HaloTag: a novel protein labeling technology for cell imaging and protein analysis. *ACS Chem Biol* 3:373–382
95. Losonczy A, Makara JK, Magee JC (2008) Compartmentalized dendritic plasticity and input feature storage in neurons. *Nature* 452:436–441
96. Loudet A, Burgess K (2007) BODIPY dyes and their derivatives: syntheses and spectroscopic properties. *Chem Rev* 107:4891–4932
97. Luo S, Zhang E, Su Y, Cheng T, Shi C (2011) A review of NIR dyes in cancer targeting and imaging. *Biomaterials* 32:7127–7138
98. Ma Q, Su X (2011) Recent advances and applications in QDs-based sensors. *Analyst* 136:4883–4893
99. Maçôas E, Marcelo G, Pinto S, Cañeque T, Cuadro AM, Vaquerob JJ, Martinhoa JMG (2011) A V-shaped cationic dye for nonlinear optical bioimaging. *Chem Commun* 47:7374–7376
100. Maeda H, Fukuyasu Y, Yoshida S, Fukuda M, Saeki K, Matsuno H, Yamauchi Y, Yoshida K, Hirata K, Miyamoto K (2004) Fluorescent probes for hydrogen peroxide based on a non-oxidative mechanism. *Angew Chem Int Ed* 43:2389–2391
101. Marcus RA (1993) Electron transfer reactions in chemistry: theory and experiment (Nobel lecture). *Angew Chem Int Ed Engl* 32:1111–1121
102. Marks KM, Rosinov M, Nolan GP (2004) In vivo targeting of organic calcium sensors via genetically selected peptides. *Chem Biol* 11:347–356
103. Martin BR, Giepmans BNG, Adams SR, Tsien RY (2005) Mammalian cell-based optimization of the biarsenical-binding tetracycline motif for improved fluorescence and affinity. *Nat Biotechnol* 23:1308–1314
104. Matsui A, Umezawa K, Shindo Y, Fujii T, Citterio D, Oka K, Suzuki K (2011) A near-infrared fluorescent calcium probe: a new tool for intracellular multicolour  $\text{Ca}^{2+}$  imaging. *Chem Commun* 47:10407–10409
105. Meier SD, Kovalchuk Y, Rose CR (2006) Properties of the new fluorescent  $\text{Na}^+$  indicator CoroNa Green: comparison with SBFI and confocal  $\text{Na}^+$  imaging. *J Neurosci Methods* 155:251–259
106. Miller EW, Lin JY, Frady EP, Steinbach PA, Kristan WB Jr, Tsien RY (2012) Optically monitoring voltage in neurons by photoinduced electron transfer through molecular wires. *Proc Natl Acad Sci USA* 109:2114–2119
107. Minta A, Kao JPY, Tsien RY (1989) Fluorescent indicators for cytosolic calcium based on rhodamine and fluorescein chromophores. *J Biol Chem* 264:8181–8178
108. Minta A, Tsien RY (1989) Fluorescent indicators for cytosolic sodium. *J Biol Chem* 264:19449–19457
109. Miyawaki A (2011) Development of probes for cellular functions using fluorescent proteins and fluorescence resonance energy transfer. *Annu Rev Biochem* 80:357–373



110. Mizukami S, Watanabe S, Hori Y, Kikuchi K (2009) Covalent protein labeling based on noncatalytic  $\beta$ -lactamase and a designed FRET substrate. *J Am Chem Soc* 131:5016–5017
111. Muyskens M (2006) The fluorescence of *Lignum nephriticum*: a flash back to the past and a simple demonstration of natural substance fluorescence. *J Chem Educ* 83:765–768
112. Myochin T, Kiyose K, Hanaoka K, Kojima H, Terai T, Nagano T (2011) Rational design of ratiometric near-infrared fluorescent pH probes with various  $pK_a$  values, based on aminocyanine. *J Am Chem Soc* 133:3401–3409
113. Nagano T (2009) Bioimaging probes for reactive oxygen species and reactive nitrogen species. *J Clin Biochem Nutr* 45:111–124
114. Namkung W, Padmawar P, Mills AD, Verkman AS (2008) Cell-based fluorescence screen for  $K^+$  channels and transporters using an extracellular triazacryptand-based  $K^+$  sensor. *J Am Chem Soc* 130:7794–7795
115. Niu Y, Peng Q, Deng C, Gao X, Shuai Z (2010) Theory of excited state decays and optical spectra: application to polyatomic molecules. *J Phys Chem A* 114:7817–7831
116. Nolan EM, Lippard SJ (2009) Small-molecule fluorescent sensors for investigating zinc metalloneurochemistry. *Acc Chem Res* 42:193–203
117. Ogawa M, Kosaka N, Choyke PL, Kobayashi H (2009) In vivo molecular imaging of cancer with a quenching near-infrared fluorescent probe using conjugates of monoclonal antibodies and indocyanine green. *Cancer Res* 69:1268–1272
118. Ojida A, Honda K, Shinmi D, Kiyonaka S, Mori Y, Hamachi I (2006) Oligo-Asp tag/Zn(II) complex probe as a new pair for labeling and fluorescence imaging of proteins. *J Am Chem Soc* 128:10452–10459
119. Padmawar P, Yao X, Bloch O, Manley GT, Verkman AS (2005)  $K^+$  waves in brain cortex visualized using a long wavelength  $K^+$ -sensing fluorescent indicator. *Nat Methods* 2:825–827
120. Razgulin A, Ma N, Rao J (2011) Strategies for in vivo imaging of enzyme activity: an overview and recent advances. *Chem Soc Rev* 40:4186–4216
121. Rehm D, Weller A (1970) Kinetics of fluorescence quenching by electron and H-atom transfer. *Israel J Chem* 8:259–271
122. Reinert KC, Dunbar RL, Gao W, Chen G, Ebner TJ (2004) Flavoprotein autofluorescence imaging of neuronal activation in the cerebellar cortex in vivo. *J Neurophysiol* 92:199–211
123. Robinson D, Besley NA, O'Shea P, Hirst JD (2011) Di-8-ANEPPS emission spectra in phospholipid/cholesterol membranes: a theoretical study. *J Phys Chem B* 115:4160–4167
124. Rosania GR, Lee JW, Ding L, Yoon H-S, Chang Y-T (2003) Combinatorial approach to organelle-targeted fluorescent library based on the styryl scaffold. *J Am Chem Soc* 125:1130–1131
125. Rust MJ, Bates M, Zhuang X (2006) Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM). *Nat Methods* 3:793–795
126. Sakabe M, Asanuma D, Kamiya M, Iwatate RJ, Hanaoka K, Terai T, Nagano T, Urano Y (2013) Rational design of highly sensitive fluorescence probes for protease and glycosidase based on precisely controlled spirocyclization. *J Am Chem Soc* 135:409–414
127. Samanta A, Vendrell M, Dasa R, Chang Y-T (2010) Development of photostable near-infrared cyanine dyes. *Chem Commun* 46:7406–7408
128. Shen Z, Lu Z, Chhatbar PY, O'Herron P, Kara P (2012) An artery-specific fluorescent dye for studying neurovascular coupling. *Nat Methods* 9:273–276
129. Shepherd J, Hilderbrand SA, Waterman P, Heinecke JW, Weissleder R, Libby P (2007) A fluorescent probe for the detection of myeloperoxidase activity in atherosclerosis-associated macrophages. *Chem Biol* 14:1221–1231
130. Shieh CC, Coghlan M, Sullivan JP, Gopalakrishnan M (2000) Potassium channels: molecular defects, diseases, and therapeutic opportunities. *Pharmacol Rev* 52:557–593
131. Shimomura O, Johnson FH, Saiga Y (1962) Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusa, *Aequorea*. *J Cell Comp Physiol* 59:223–239
132. Shultz MD, Janes KA, Lauffenburger DA, Imperiali B (2005) A multiplexed homogeneous fluorescence-based assay for protein kinase activity in cell lysates. *Nat Methods* 2:277–284
133. Skala MC, Riching KM, Gendron-Fitzpatrick A, Eickhoff J, Eliceiri KW, White JG, Ramanujam M (2007) In vivo multiphoton microscopy of NADH and FAD redox states, fluorescence lifetimes, and cellular morphology in precancerous epithelia. *Proc Natl Acad Sci USA* 104:10494–10499
134. Somers RC, Bawendi MG, Nocera DG (2007) CdSe nanocrystal based chem-/bio- sensors. *Chem Soc Rev* 36:579–591
135. Srikun D, Albers AE, Nam CI, Iavarone AT, Chang CJ (2010) Organelle-targetable fluorescent probes for imaging hydrogen peroxide in living cells via SNAP-tag protein labeling. *J Am Chem Soc* 132:4455–4465
136. Stokes GG (1845) On the change of refrangibility of light. *Phil Trans Roy Soc London* 142:463–562
137. Takahashi A, Camacho P, Lechleiter JD, Herman B (1999) Measurement of intracellular calcium. *Physiol Rev* 79:1089–1125
138. Terai T, Kikuchi K, Urano Y, Kojima H, Nagano T (2012) A long-lived luminescent probe to sensitively detect arylamine *N*-acetyltransferase (NAT) activity of cells. *Chem Commun* 48:2234–2236
139. Terai T, Nagano T (2008) Fluorescent probes for bioimaging applications. *Curr Opin Chem Biol* 12:515–521
140. Tomat E, Nolan EM, Jaworski J, Lippard SJ (2008) Organelle-specific zinc detection using zinpyr-labeled fusion proteins in live cells. *J Am Chem Soc* 130:15776–15777
141. Tour O, Adams SR, Kerr RA, Meijer RM, Sejnowski TJ, Tsien RW, Tsien RY (2007) Calcium Green FAsH as a genetically targeted small-molecule calcium indicator. *Nat Chem Biol* 3:423–431
142. Touthkine A, Kraynov V, Hahn K (2003) Solvent-sensitive dyes to report protein conformational changes in living cells. *J Am Chem Soc* 125:4132–4145
143. Trapani V, Farruggia G, Marraccini C, Iotti S, Cittadinia A, Wolf FI (2010) Intracellular magnesium detection: imaging a brighter future. *Analyst* 135:1855–1866
144. Tsien RY (1980) New calcium indicators and buffers with high selectivity against magnesium and protons: design, synthesis, and properties of prototype structures. *Biochem* 19:2396–2404
145. Tsien RY (1998) The green fluorescent protein. *Annu Rev Biochem* 67:509–544
146. Ueno T, Urano Y, Setsukinai K, Takakusa H, Kojima H, Kikuchi K, Ohkubo K, Fukuzumi S, Nagano T (2004) Rational principles for modulating fluorescence properties of fluorescein. *J Am Chem Soc* 126:14079–14085
147. Umezawa K, Nakamura Y, Makino H, Citterio D, Suzuki K (2008) Bright, color-tunable fluorescent dyes in the visible-near-infrared region. *J Am Chem Soc* 130:1550–1551
148. Urano Y, Asanuma D, Hama Y, Koyama Y, Barrett T, Kamiya M, Nagano T, Watanabe T, Hasegawa A, Choyke PL, Kobayashi H (2009) Selective molecular imaging of viable cancer cells with pH-activatable fluorescence probes. *Nat Med* 15:104–109
149. Urano Y, Kamiya M, Kanda K, Ueno T, Hirose K, Nagano T (2005) Evolution of fluorescein as a platform for finely tunable fluorescence probes. *J Am Chem Soc* 127:4888–4894
150. Veal EA, Day AM, Morgan BA (2007) Hydrogen peroxide sensing and signaling. *Mol Cell* 26:1–14

151. Vendrell M, Zhai D, Er JC, Chang Y-T (2012) Combinatorial strategies in fluorescent probe development. *Chem Rev* 112:4391–4420
152. von Prowazek S (1914) Über Fluoreszenz der Zellen. *Kleinwelt* 6:37–40
153. Waggoner AS (1979) Dye indicators of membrane potential. *Ann Rev Biophys Bioeng* 8:47–68
154. Wahler D, Badalassi F, Crotti P, Reymond J-L (2002) Enzyme fingerprints of activity, and stereo- and enantioselectivity from fluorogenic and chromogenic substrate arrays. *Chem Eur J* 8:3211–3228
155. Walkup GK, Imperiali B (1997) Fluorescent chemosensors for divalent zinc based on zinc finger domains. Enhanced oxidative stability, metal binding affinity, and structural and functional characterization. *J Am Chem Soc* 119:3443–3450
156. Wardman P (2007) Fluorescent and luminescent probes for measurement of oxidative and nitrosative species in cells and tissues: progress, pitfalls, and prospects. *Free Radic Biol Med* 43:995–1022
157. Weissleder R, Ntziachristos V (2003) Shedding light onto live molecular targets. *Nat Med* 9:123–128
158. Whitaker JE, Haugland RP, Prendergast FG (1991) Spectral and photophysical studies of benzo[c]xanthene dyes: dual emission pH sensors. *Anal Biochem* 194:330–344
159. Whitney MA, Crisp JL, Nguyen LT, Friedman B, Gross LA, Steinbach P, Tsien RY, Nguyen QT (2011) Fluorescent peptides highlight peripheral nerves during surgery in mice. *Nat Biotechnol* 29:352–356
160. Wysockia LM, Lavis LD (2011) Advances in the chemistry of small molecule fluorescent probes. *Curr Opin Chem Biol* 15:752–759
161. Yap YW, Whiteman M, Cheung NS (2007) Chlorinative stress: an under appreciated mediator of neurodegeneration? *Cell Signal* 19:219–228
162. Yu FH, Catterall WA (2003) Overview of the voltage-gated sodium channel family. *Genome Biol* 4:207.1–207.7
163. Yuan L, Lin W, Yang Y, Chen H (2012) A unique class of near-infrared functional fluorescent dyes with carboxylic-acid-modulated fluorescence ON/OFF switching: rational design, synthesis, optical properties, theoretical calculations, and applications for fluorescence imaging in living animals. *J Am Chem Soc* 134:1200–1211
164. Zhai D, Lee S-C, Vendrell M, Leong LP, Chang Y-T (2012) Synthesis of a novel BODIPY library and its application in the discovery of a fructose sensor. *ACS Comb Sci* 14:81–84
165. Zhang J, Campbell RE, Ting AY, Tsien RY (2002) Creating new fluorescent probes for cell biology. *Nat Rev Mol Cell Biol* 3:906–918
166. Zlokarnik G, Negulescu PA, Knapp TE, Mere L, Burres N, Feng L, Whitney M, Roemer K, Tsien RY (1998) Quantitation of transcription and clonal selection of single living cells with  $\beta$ -lactamase as reporter. *Science* 279:84–88