

INVITED REVIEW

Disease modelling using human iPSCs

Stormy J. Chamberlain*

Department of Genetics and Genome Sciences, University of Connecticut Health Center, Farmington, CT, USA

*To whom correspondence should be addressed at: Department of Genetics and Genome Sciences, University of Connecticut Health Center, 400 Farmington Ave, Farmington, CT 06030-6403, USA. Tel.: 860-679-4433; Fax: 860-679-8345; Email: chamberlain@uchc.edu

Introduction to iPSC Technology

Human embryonic stem cells (hESCs) paved the way for the development of induced pluripotent stem cells (iPSCs). First derived and described by James Thomson in 1998, hESCs demonstrated that human pluripotent stem cells could be cultivated and could differentiate into all three germ layers, albeit as part of a teratoma in a mouse (1). Importantly, the development of hESC lines established media recipes and cell culture techniques comprising the cornerstone for iPSC derivation (1). Furthermore, they enabled researchers to gain experience culturing these somewhat peculiar and difficult stem cells and establish differentiation protocols (1–5). The lack of patient-specific hESCs prompted focus to be shifted to establishing banks of hESCs that would be representative of entire populations (6). In parallel, the use of somatic cell nuclear transfer (SCNT) or cloning to generate human patient-specific stem cell lines was developed (7). Neither approach ameliorated the ethical issues surrounding the use of sensitive human tissues (in vitro derived embryos and human oocytes) and the possibility that the process would be used to create genetically-modified human beings.

In 2006, Shinya Yamanaka rocked the stem cell world by reporting the first iPSCs from murine fibroblasts (8). Built upon previous experiments that identified transcription factors differentially expressed between pluripotent stem cells and differentiated cells (9), Takahashi and Yamanaka tested the ability of 20 factors *en masse* to convert somatic cells into pluripotent stem cells. After narrowing the number of required factors down to four—known widely as the Yamanaka four: OCT4, SOX2, KLF4 and C-MYC—he and James Thomson independently reported the reprogramming of human fibroblasts into pluripotent stem cells (10,11). Although additional factors were added to the mix, the community has largely settled on the Yamanaka

four plus or minus LIN28 as being the most efficient way to reprogram human somatic cells into iPSCs.

The development of human iPSCs from patient-specific samples has revolutionized the study of human genetic disorders over the past 10 years. The next 10 years are certain to see more rapid development of targeted therapeutics for some of these disorders, thanks to this technological advance.

State of the iPSC Art

Improvements to the process of epigenetic reprogramming into iPSCs followed rapidly due to the fervour behind the exciting technology. Different cell types permissive for reprogramming were tested to improve reprogramming efficiency, reduce the number of factors required for reprogramming, or to reduce the invasiveness of obtaining patient samples. Fibroblasts derived from dermal punch biopsies were first used to derive iPSCs (10,11). Owing to previously banked fibroblasts from individuals with genetic disorders, iPSCs could be derived from many disorders without seeking additional skin biopsies. Nonetheless, obtaining a skin punch biopsy from a new or rare patient to establish fibroblast lines and derive iPSCs is somewhat invasive. Neural stem cells and renal tubular cells were shown to only require one or two reprogramming factors, but it was not feasible to obtain neural stem cells from most patients (12,13). Keratinocytes were reported to be non-invasive and easily obtainable from patients by plucking hairs, but the establishment of keratinocyte cultures from the hair follicle samples proved to be challenging, and the reprogramming was difficult to reproduce early on (14). Eventually, peripheral blood mononuclear cells (PBMCs) were shown to be both easily obtained from patients and easily reprogrammed, largely solving the somatic cell source problem (15). Recent reports even demonstrated the

It has been 10 years since Shinya Yamanaka rocked the scientific community by reporting the first derivation of induced pluripotent stem cells (iPSCs). This review highlights some of the accomplishments made in iPSC models of genetic disorders, focusing mainly on neural and cardiac disease.

Received: June 23, 2016. Revised: June 23, 2016. Accepted: June 27, 2016

© The Author (2016). Published by Oxford University Press.

All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

ability to reprogram renal epithelial cells from urine (16) and dental pulp stem cells from deciduous teeth (17).

Improvements to the delivery of the reprogramming factors also quickly followed. The field quickly replaced retroviral vectors, which required replication to transduce somatic cells, with lentiviral vectors (11). Worry about insertional mutation due to viral integration led to the development of the STEMCCA vector in which all reprogramming factors were included on a single lentiviral vector that could be excised using Cre-recombinase (18). Provision of the reprogramming factors as mRNAs (19) or proteins (20,21) was a non-integrating answer to the viral integration issue. These approaches were not adopted widely due to the difficulty of generating sufficient quantities of high-quality mRNAs or proteins and/or the difficulty delivering them into somatic cells. Finally, Sendai virus vectors (22) and episomal vectors (23) have met widespread adoption because they meet the criteria of being both non-integrating and easy to use. Both reprogramming systems are commercially available, making the process accessible to many labs and reproducible.

The Promises and Pitfalls of iPSC Disease Models

The promise of iPSCs as a clinical and research tool was apparent in two different ways. First, translational research applications for pluripotent stem cells were rooted in regenerative medicine with the idea that stem cells could be differentiated and implanted into individuals to cure injury or disease. Thus, one immediate application for iPSCs was autologous stem cell transplants. **If one could generate pristine iPSCs from any given patient, then they could provide their own tissue to repair/replace injured ones.** Second, many recognized that **the genetics of the iPSCs reflected the genetics of the patient.** Thus, iPSCs could be used to model human genetic disorders. Since it was still relatively difficult to use homologous recombination in hESCs, iPSCs provided a more direct way to study cells harbouring a particular genetic disorder, including those difficult to generate by homologous recombination.

Early studies of iPSCs focused on identifying differences between hESCs and iPSCs to identify *bona fide* iPSCs and/or ascertain the quality of individual iPSC clones (24). These studies turned toward epigenetic analyses, since the difference between the iPSCs and their somatic counterparts were known to be epigenetic and variances were suspected to lie somewhere in between those extremes. Indeed, one study reported that iPSCs retained the epigenetic signature of their starting somatic cells, which in turn, affects their differentiation potential (25). Another study reported a realm of pluripotency, in which hESCs occupied a small niche, and iPSCs occupied a larger, but overlapping niche (26).

While these epigenetic studies focused on distinguishing iPSCs from hESCs, they also pointed to larger epigenetic changes that distinguish somatic cells from pluripotent stem cells. This has an impact on the use of iPSC for disease modelling. The altering of epigenetic states between somatic cells and iPSCs may obscure the retention of disease state for disorders that involve epigenetic modification (i.e. imprinting disorders or X-linked disorders in females) (27) or for disorders with mixed aetiology in which other factors such as the environment or epigenetic alterations may play a role. This possibility remains a warning for those using iPSCs to model disorders with unknown aetiologies. With this limitation aside, iPSCs are beginning to hit their stride at models for human disease.

iPSC Models of Human Neural Disorders

The earliest iPSC models of human disease were reported from patients with neurological disorders. This may have been due to the fact that the earliest hESC differentiation protocols were focused on neurons. Early differentiation protocols utilized developmental principles to differentiate and pattern neurons (28). They usually involved embryoid body (EB) intermediates, which emulates gastrulation to direct some tissue down the ectodermal lineage (2,29). More recent neural differentiation protocols capitalized on dual SMAD inhibition using a combination of Noggin or the small molecule LDN193189 along with SB431542 or Noggin alone to restrict differentiation to ectodermal tissues (30). Serum-free media formulated to encourage the growth of neural cell types and substrates required for neural cell adhesion were used to preferentially encourage the growth and expansion of neural ectoderm in both approaches. Morphogens such as retinoic acid and sonic hedgehog are used to caudalize and ventralize the neural ectoderm, respectively.

Beginning in 2010, Marius Wernig and Thomas Sudhof published a series of papers reporting the direct conversion of mouse and human fibroblasts into neurons using transcription factors. Combinations of BRN2, ASCL1 and MYT1L plus NEUROD1 in humans were used to convert fibroblasts into neurons. Ultimately, the addition of only one transcription factor, NGN2, was shown to convert hESCs and human iPSCs into neurons. Following neural induction, in vitro-derived human neurons require maturation to develop synaptic activity and mature action potentials. This process seems to take approximately 7 weeks, except for iN cells, which are reported to take a week. Studies of dendritic spine morphology and densities may take even longer. Nonetheless, in vitro differentiated neurons demonstrate typical electrophysiological properties, and may even be amenable to studies of synaptic plasticity.

Neurodegenerative disorders such as amyotrophic lateral sclerosis (ALS) (31–38), Huntington disease (39–41), Parkinson's disease (PD) (42–47), and Alzheimer disease (48–50) are among the most studied disorders using iPSCs. iPSC-derived neurons from these disorders have been reported to have altered neuronal morphology, including neurite degeneration, reduced soma size, and disturbed mitochondrial morphology. Protein aggregate formation has been observed in neurons, as has increased cellular stress. The ability of in vitro-derived neurons to recapitulate cellular phenotypes is remarkable since these disorders usually present in aged individuals, and the neurons represent fetal neurons. A neurodegenerative disorder that affects young children, spinal muscular atrophy (SMA), has also been modelled by using iPSCs by multiple groups (51–53). Cellular phenotypes associated with this disorder include hyperexcitability, smaller soma size, incomplete synapses, and a reduction in cytoplasmic and nuclear aggregate structures, called gems. iPSCs from individuals with familial dysautonomia, a fatal peripheral neuropathy, have also been made (54). Upon differentiation into neural crest precursors and peripheral neurons, differentiation and migration defects were observed. Table 1 summarizes many of the neural disorders that have been modelled using iPSCs and lists the neural phenotypes identified for each disorder.

Tremendous progress has also been made modelling neurodevelopmental disorders using pluripotent stem cells. Induced pluripotent stem cells have been derived from individuals with Rett (55–57), Fragile X (58–60), Timothy (61,62), Dravet (63,64), Prader-Willi (65–67), Angelman (65), Dup15q (68), 15q11.2 micro-deletion (69), and Phelan McDermid syndromes (70). For many

Table 1. iPSC models of neural disorders

Disorder	Genes involved	Neural phenotypes	Refs
Neurodegenerative			
ALS	TDP43, SOD1, C9ORF72, VAPB, FUS, sporadic	Neuronal death, neurite degeneration, reduced soma size, and disturbed mitochondrial morphology, altered excitability	(31–38)
SMA	SMN1	Increased apoptosis, altered axonal transport, hyperexcitability	(51–53)
Parkinsons' Disease	PARKIN1, PARKIN2, PINK, SNCA, LRRK2	α -synuclein aggregates, increased nitrosative stress, increased endoplasmic reticulum associated degradation products	(42–47)
Huntington Disease	HTT	Elevated glutamate evoked responses, higher lysosomal activity, increased caspase activity	(39–41,99,100)
Alzheimer Disease	APP, PS1, PS2	Increased A β 40, phosphorylated Tau, and active GSK3 β	(48–50)
Familial Dysautonomia	IKBKAP	Defective differentiation, migration	(54)
Neurodevelopmental			
Rett syndrome	MECP2	Reduced dendritic spine density, smaller soma size, altered electrophysiology, alterations in Ca ²⁺ influx and fewer synapses	(55–57,101)
Fragile X syndrome	FMR1	Fewer synapses, reduced neurite length, increased Ca ²⁺ transients	(58–60)
Angelman syndrome	UBE3A, 15q11.2-q13 deletion	Neural phenotypes not studied other than gene expression	(65)
Prader-Willi syndrome	15q11.2-q13 deletion	Neural phenotypes not studied other than gene expression	(65–67,102)
Dup15q syndrome	15q11.2-q13 duplication	Neural phenotypes not studied other than gene expression	(68)
Idiopathic autism	TRPC6, CHD8, CNTNAP2	Altered neuronal migration, reduced dendritic length/branching, reduced dendritic spine density	(79,80)
Dravet syndrome	SNC1A	Altered Na ⁺ current density, spontaneous activity	(63,64)
Timothy syndrome	CACNA1	Altered Ca ²⁺ signaling, dendritic retraction	(61, 62,103)
Phelan-McDermid syndrome	SHANK3	Fewer synapses, altered electrophysiology	(70)
Schizophrenia	22q11 deletion, DISC1, sporadic	Aberrant migration, altered WNT signaling, and increased oxidative stress in neural progenitor cells, increased catecholamine expression in neurons	(72–76,104)
Bipolar disorder	Sporadic, but familial	Decreased proliferation of neural stem cells, mitochondrial defects, hyperexcitability	(77,78)
15q11.2 deletion syndrome	CYFIP, TUBGCP5, NIPA1, NIPA2	Altered dendritic morphology	(69)
Hereditary spastic paraplegia	SPAST, ATL1, SPG11	Motor neurons, forebrain glutamatergic neurons	(81–83)

of these disorders, forebrain glutamatergic neurons have been generated, although other neuronal subtypes are also being investigated. As these disorders are thought to influence synaptic activity, electrophysiology and morphological analyses of iPSC-derived neurons have been employed to study these disorders. **Morphological cellular phenotypes observed have included small soma size, fewer synapses, lower synaptic protein levels, and reduced neurite length.** Electrophysiological and functional phenotypes observed have included **abnormal calcium signaling, altered sodium current density, prolonged action potential patterns, and altered excitatory synaptic activity** (Table 1). These phenotypes are likely the tip of the iceberg for neurodevelopmental disorders. Our understanding of neural development and neuronal subtype specification is increasing and is being translated into differentiation protocols. This will enable the dissection of cell autonomous and non-cell autonomous phenotypes, as well as cellular phenotypes that require a precise balance of neuronal subtypes. Furthermore, the development of assays adapted for cultured neurons—for instance, chemically induced synaptic plasticity—is expanding. Thus, we can expect a reciprocal expansion of electrophysiological phenotypes.

Other neural disorders that have been modelled using human iPSCs include the psychiatric disorders, schizophrenia (71–76) and bipolar disorder (77,78); idiopathic autism spectrum disorder (79,80); and hereditary spastic paraplegia (81–83). Although schizophrenia and autism spectrum disorder do not have known genetic causes in most affected individuals, they have high heritability. iPSCs have been derived either from one of the rare known genetic subtypes of these disorders, from families with a clear autosomal dominant inheritance pattern, or from an individual with severe early onset disease. Thus illustrating the ability of iPSCs to model disorders of unknown genetic etiologies. iPSCs derived from individuals with a few different forms of hereditary spastic paraplegia are chipping away at the variety of different cellular phenotypes associated with several phenotypically similar disorders.

iPSC Models of Inherited Heart Disease

Like their neural counterparts, differentiation of iPSCs into cardiomyocytes follows embryonic developmental pathways. Also like their neural counterparts, cardiomyocyte differentiation can be accomplished using embryoid body-mediated and

Table 2. iPSC models of cardiac disorders

Disorder	Genes	Cardiomyocyte phenotypes	Ref
Classic cardiomyopathies			
HCM	MYH7, MYBPC3	Cellular enlargement, contractile arrhythmia, dysregulated Ca ²⁺ cycling, increased expression of HCM genes, Cellular enlargement, myofibrillar disarray	(84–86)
DCM	TNNT2, DES, TTN, RBM20	Altered Ca ²⁺ handling, impaired response to β -adrenergic stress, decreased contractility, disorganized sarcomeres, abnormal α -actinin distribution in sarcomeres, Abnormal aggregation of DES, Abnormal aggregation of DES	(87–91)
ARVC	PKP2	Cellular enlargement, altered Ca ²⁺ handling, increased adipogenicity, reduced localization of cell surface desmosomal proteins, metabolic defects, and abnormal (PPAR- γ) activation	(92–94)
Channelopathies			
Long QT syndrome	KCNH2, KCNQ1, CACNA1C	Prolonged action potentials, early afterdepolarizations, susceptibility to arrhythmias, altered Ca ²⁺ handling,	(103,105–110)
CVPT	RYR2, CASQ2	Delayed afterdepolarizations, altered Ca ²⁺ handling, disorganized myofibrils, contractile arrhythmia	(111,112)
Mitochondrial disorders			
Barth syndrome	TAZ	Sparse/irregular sarcomeres, contractile abnormalities	(113)
Friedrich's ataxia	FXN	Impaired Ca ²⁺ handling, cellular enlargement, impaired β -adrenergic response	(114)
CPT II	CPT2	Not determined	(115)
Other cardiac disease-associated rare genetic disorders			
LEOPARD syndrome	PTPN11	Cellular enlargement, increased sarcomeric organization, nuclear NFATC4 localization	(116)
Pompe disease	GAA	Glycogen-filled lysosomes	(117–120)
Lamin A/C mutations	LMNA	Accelerated senescence and apoptosis	(121)

monolayer-based protocols. **In both cases, iPSCs are first coaxed into accepting a mesodermal identity, and then specified and matured into cardiac cells.** BMP, TGF β /activin/NODAL, WNT, and FGF signalling are all important for differentiation into myocardial cells. Differentiated cardiomyocytes express TNNT2 and MEF2C and exhibit intact sarcomeres and spontaneous rhythmic contraction. iPSC-derived cardiomyocytes also require *in vitro* maturation, and despite best efforts, still display phenotypes and gene expression profiles most similar to fetal cardiomyocytes. Continued improvement to differentiation protocols, including 3D tissue engineering strategies will benefit these *in vitro* models of cardiac development.

iPSC models of the three major classes of pure genetic cardiomyopathies have been developed: hypertrophic cardiomyopathy (HCM; 84–86), dilated cardiomyopathy (DCM; 87–91), and arrhythmogenic right ventricular cardiomyopathy (ARVC; 92–94). These three types of cardiomyopathy typically affect the whole beating heart, presenting unique challenges for these disorders. Additionally, iPSC models of other genetic disorders resulting in heart disease have been created. These include channelopathies such as long QT syndrome and catecholaminergic polymorphic ventricular tachycardia (CPVT; 103,105–110); mitochondrial disorders such as Barth syndrome (113), Friedrich's ataxia (114), and carnitine palmitoyltransferase II deficiency (CPT II; 115); and other rare genetic conditions associated with cardiac disease, such as LEOPARD syndrome (116), Pompe disease (117–120), and laminopathies (121). Significant progress has been made, and iPSC models have contributed to an understanding of the mechanistic underpinnings of cardiac disease. Table 2 summarizes the iPSC models of cardiomyopathies or cardiac associated disorders as well as their cellular phenotypes.

Genetic forms of HCM, DCM, and ARVC can be caused by mutations in structural proteins that make up cardiac muscle.

Genetic forms of HCM have been modelled by iPSCs derived from individuals with mutations in MYH7 and MYBPC3 (84–86). iPSCs have been derived from patients with DCM caused by mutations in TTN, DES, TNNT2 and RBM20 (87–91). ARVC has been modelled using iPSCs harbouring mutations in PKP2 (92–94). iPSC-derived cardiomyocytes from individuals with these disorders show structural abnormalities such as cellular enlargement, sarcomere and/or myofibrillar disorganization, and mislocalized or aberrantly aggregated structural proteins. Importantly, they also show functional deficits such as dysregulated calcium (Ca²⁺) handling, contractile dysfunction, and impaired responses to β -adrenergic or mechanical stress (Table 2).

Studies in cardiomyocytes generated from iPSC models of other genetic disorders involving cardiac dysfunction (i.e. Barth syndrome, Friedrich's ataxia, CPT II, LEOPARD syndrome, and Pompe disease) show many overlapping features with HCM, DCM, and ARVC, including cellular enlargement, altered Ca²⁺ handling, and contractile abnormalities (Table 2). As expected, each of these disorders also frequently show phenotypic features unique to their genetic etiologies, as well as non-cardiac features in other iPSC-derivatives.

Other Genetic Disorders

Many other genetic disorders have been modelled using patient-specific iPSCs in the past 10 years. iPSC models of Duchenne's Muscular Dystrophy (41,95) have been generated and differentiated into myocytes and myotubes. iPSCs from individuals with juvenile diabetes have been differentiated into insulin producing cells (41). Hepatocytes derived from an individual with familial hypercholesterolemia were deficient in LDL-C uptake and had elevated secretion of a lipidated protein (96–98). Kotton et al. even generated more than 100 disease-

specific iPSC lines from individuals with a variety of different diseases that affect the lung, including cystic fibrosis, α -1 antitrypsin deficiency, scleroderma, and sickle-cell disease (18). Furthermore, they were able to differentiate them into definitive endoderm, a developmental progenitor to lung tissue. Although there may be some disorders that are difficult to model through this epigenetic reprogramming (i.e. some imprinting disorders), the technology is broadly applicable to almost any human genetic disorder, limited mainly by the ability to readily differentiate the cells into disease-relevant cell types.

Looking Forward

In the 10 years since the first discovery of epigenetic reprogramming into iPSCs, work has been done by many different groups to model a variety of genetic disorders, in part because of its ease, accessibility, and commercialization. With many such disease models in existence, new technologies such as direct conversion to differentiated cell types, 3D culture systems, and clustered regularly interspersed repeats (CRISPR) further expand the possibilities for iPSC-based disease modelling. For example, direct lineage conversion of iPSCs into neurons as described above, circumvents the slow pace of neural development. While this might not be the most developmentally correct differentiation strategy, it may facilitate the use of iPSC-derived neurons for high-throughput drug screening. 3D cultures enable the study of phenotypes that are inaccessible or undetectable in monolayer or colony-based cultures. For instance, 3D cardiac microtissues harbouring mutations in *TTN* were required to uncover the disease-related deficits in contractile function (90). Using CRISPR or related technologies, causative mutations have been genetically corrected for ALS, Parkinson's disease, and many other disorders. This enables comparison of isogenic control samples and greater confidence in cellular phenotypes of disease.

Acknowledgements

The author regrets that not all iPSC models of human disease could be discussed here. Overlooked papers or findings were not intentional.

Conflict of Interest statement. None declared.

Funding

The author acknowledges the following funding sources: NICHD/NIH (R01HD068730); the Connecticut Regenerative Medicine Research Fund (09SCAUCHC14 and 14-SCDIS-UCHC-01), and the Raymond and Beverly Sackler foundation.

References

1. Thomson, J.A., Itskovitz-Eldor, J., Shapiro, S.S., Waknitz, M.A., Swiergiel, J.J., Marshall, V.S. and Jones, J.M. (1998) Embryonic stem cell lines derived from human blastocysts. *Science*, **282**, 1145–1147.
2. Li, X.J. and Zhang, S.C. (2006) In vitro differentiation of neural precursors from human embryonic stem cells. *Methods Mol. Biol.*, **331**, 177–1.
3. Xue, T., Cho, H.C., Akar, F.G., Tsang, S.Y., Jones, S.P., Marban, E., Tomaselli, G.F. and Li, R.A. (2005) Functional integration of electrically active cardiac derivatives from

- genetically engineered human embryonic stem cells with quiescent recipient ventricular cardiomyocytes: insights into the development of cell-based pacemakers. *Circulation*, **111**, 11–20.
4. Yan, Y., Yang, D., Zarnowska, E.D., Du, Z., Werbel, B., Valliere, C., Pearce, R.A., Thomson, J.A. and Zhang, S.C. (2005) Directed differentiation of dopaminergic neuronal subtypes from human embryonic stem cells. *Stem Cells*, **23**, 781–790.
5. Li, X.J., Du, Z.W., Zarnowska, E.D., Pankratz, M., Hansen, L.O., Pearce, R.A. and Zhang, S.C. (2005) Specification of motoneurons from human embryonic stem cells. *Nat Biotechnol.*, **23**, 215–221.
6. Healy, L., Hunt, C., Young, L. and Stacey, G. (2005) The UK Stem Cell Bank: its role as a public research resource centre providing access to well-characterised seed stocks of human stem cell lines. *Adv. Drug Deliv. Rev.*, **57**, 1981–1988.
7. Tachibana, M., Amato, P., Sparman, M., Gutierrez, N.M., Tippner-Hedges, R., Ma, H., Kang, E., Fulati, A., Lee, H.S., Sritanaudomchai, H., et al. (2013) Human embryonic stem cells derived by somatic cell nuclear transfer. *Cell*, **153**, 1228–1238.
8. Takahashi, K. and Yamanaka, S. (2006) Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*, **126**, 663–676.
9. Takahashi, K., Ichisaka, T. and Yamanaka, S. (2006) Identification of genes involved in tumor-like properties of embryonic stem cells. *Methods Mol. Biol.*, **329**, 449–458.
10. Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K. and Yamanaka, S. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell*, **131**, 861–872.
11. Yu, J., Vodyanik, M.A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J.L., Tian, S., Nie, J., Jonsdottir, G.A., Ruotti, V., Stewart, R., et al. (2007) Induced pluripotent stem cell lines derived from human somatic cells. *Science*, **318**, 1917–1920.
12. Montserrat, N., Ramirez-Bajo, M.J., Xia, Y., Sancho-Martinez, I., Moya-Rull, D., Miquel-Serra, L., Yang, S., Nivet, E., Cortina, C., Gonzalez, F., et al. (2012) Generation of induced pluripotent stem cells from human renal proximal tubular cells with only two transcription factors, OCT4 and SOX2. *J. Biol. Chem.*, **287**, 24131–24138.
13. Kim, J.B., Zaehres, H., Wu, G., Gentile, L., Ko, K., Sebastiano, V., Arauzo-Bravo, M.J., Ruau, D., Han, D.W., Zenke, M., et al. (2008) Pluripotent stem cells induced from adult neural stem cells by reprogramming with two factors. *Nature*, **457**, 646–650.
14. Aasen, T., Raya, A., Barrero, M.J., Garreta, E., Consiglio, A., Gonzalez, F., Vassena, R., Bilic, J., Pekarik, V., Tiscornia, G., et al. (2008) Efficient and rapid generation of induced pluripotent stem cells from human keratinocytes. *Nat Biotechnol.*, **26**, 1276–1284.
15. Staerk, J., Dawlaty, M.M., Gao, Q., Maetzel, D., Hanna, J., Sommer, C.A., Mostoslavsky, G. and Jaenisch, R. (2010) Reprogramming of human peripheral blood cells to induced pluripotent stem cells. *Cell Stem Cell*, **7**, 20–24.
16. Xue, Y., Cai, X., Wang, L., Liao, B., Zhang, H., Shan, Y., Chen, Q., Zhou, T., Li, X., Hou, J., et al. (2013) Generating a non-integrating human induced pluripotent stem cell bank from urine-derived cells. *PLoS One*, **8**, e70573.
17. Yan, X., Qin, H., Qu, C., Tuan, R.S., Shi, S. and Huang, G.T. (2010) iPS cells reprogrammed from human mesenchymal-like stem/progenitor cells of dental tissue origin. *Stem Cells Dev.*, **19**, 469–480.

18. Somers, A., Jean, J.C., Sommer, C.A., Omari, A., Ford, C.C., Mills, J.A., Ying, L., Sommer, A.G., Jean, J.M., Smith, B.W., et al. (2010) Generation of transgene-free lung disease-specific human induced pluripotent stem cells using a single excisable lentiviral stem cell cassette. *Stem Cells*, **28**, 1728–1740. doi: 1710.1002/stem.1495.
19. Warren, L., Manos, P.D., Ahfeldt, T., Loh, Y.H., Li, H., Lau, F., Ebina, W., Mandal, P.K., Smith, Z.D., Meissner, A., et al. (2010) Highly efficient reprogramming to pluripotency and directed differentiation of human cells with synthetic modified mRNA. *Cell*, **7**, 618–630. Epub 2010 Sep 2030.
20. Kim, D., Kim, C.H., Moon, J.I., Chung, Y.G., Chang, M.Y., Han, B.S., Ko, S., Yang, E., Cha, K.Y., Lanza, R., et al. (2009) Generation of human induced pluripotent stem cells by direct delivery of reprogramming proteins. *Cell Stem Cell*, **4**, 472–476.
21. Zhou, H., Wu, S., Joo, J.Y., Zhu, S., Han, D.W., Lin, T., Trauger, S., Bien, G., Yao, S., Zhu, Y., et al. (2009) Generation of induced pluripotent stem cells using recombinant proteins. *Cell Stem Cell*, **4**, 381–384.
22. Seki, T., Yuasa, S. and Fukuda, K. (2012) Generation of induced pluripotent stem cells from a small amount of human peripheral blood using a combination of activated T cells and Sendai virus. *Nature Protocols*, **7**, 718–728.
23. Yu, J., Hu, K., Smuga-Otto, K., Tian, S., Stewart, R., Slukvin, I.I. and Thomson, J.A. (2009) Human induced pluripotent stem cells free of vector and transgene sequences. *Science*, **324**, 797–801.
24. Stadtfeld, M. and Hochedlinger, K. (2010) Induced pluripotency: history, mechanisms, and applications. *Genes Dev.*, **24**, 2239–2263.
25. Kim, K., Zhao, R., Doi, A., Ng, K., Unternaehrer, J., Cahan, P., Huo, H., Loh, Y.H., Aryee, M.J., Lensch, M.W., et al. (2011) Donor cell type can influence the epigenome and differentiation potential of human induced pluripotent stem cells. *Nat. Biotechnol.*, **29**, 1117–1119.
26. Guenther, M.G., Frampton, G.M., Soldner, F., Hockemeyer, D., Mitalipova, M., Jaenisch, R. and Young, R.A. (2010) Chromatin structure and gene expression programs of human embryonic and induced pluripotent stem cells. *Cell Stem Cell*, **7**, 249–257.
27. Rugg-Gunn, P.J., Ferguson-Smith, A.C. and Pedersen, R.A. (2007) Status of genomic imprinting in human embryonic stem cells as revealed by a large cohort of independently derived and maintained lines. *Hum. Mol. Genet.*, **16 Spec No. 2**, R243–R251.
28. Zhang, S.C., Wernig, M., Duncan, I.D., Brustle, O. and Thomson, J.A. (2001) In vitro differentiation of transplantable neural precursors from human embryonic stem cells. *Nat. Biotechnol.*, **19**, 1129–1133.
29. Johnson, M.A., Weick, J.P., Pearce, R.A. and Zhang, S.C. (2007) Functional neural development from human embryonic stem cells: accelerated synaptic activity via astrocyte coculture. *J. Neurosci.*, **27**, 3069–3077.
30. Germain, N.D., Banda, E.C., Becker, S., Naegele, J.R. and Gabel, L.B. (2013) Derivation and Isolation of NKX2.1-Positive Basal Forebrain Progenitors from Human Embryonic Stem Cells. *Stem Cells Dev.*, **22**, 1477–1489.
31. de Boer, A.S. and Eggan, K. (2015) A perspective on stem cell modeling of amyotrophic lateral sclerosis. *Cell Cycle*, **14**, 3679–3688.
32. Dimos, J.T., Rodolfa, K.T., Niakan, K.K., Weisenthal, L.M., Mitsumoto, H., Chung, W., Croft, G.F., Saphier, G., Leibel, R., Goland, R., et al. (2008) Induced pluripotent stem cells generated from patients with ALS can be differentiated into motor neurons. *Science*, **321**, 1218–1221.
33. Mitne-Neto, M., Machado-Costa, M., Marchetto, M.C., Bengtson, M.H., Joazeiro, C.A., Tsuda, H., Bellen, H.J., Silva, H.C., Oliveira, A.S., Lazar, M., et al. (2011) Downregulation of VAPB expression in motor neurons derived from induced pluripotent stem cells of ALS8 patients. *Hum. Mol. Genet.*, **20**, 3642–3652.
34. Bilican, B., Serio, A., Barmada, S.J., Nishimura, A.L., Sullivan, G.J., Carrasco, M., Phatnani, H.P., Puddifoot, C.A., Story, D., Fletcher, J., et al. (2012) Mutant induced pluripotent stem cell lines recapitulate aspects of TDP-43 proteinopathies and reveal cell-specific vulnerability. *Proc. Natl Acad. Sci. U S A*, **109**, 5803–5808.
35. Donnelly, C.J., Zhang, P.W., Pham, J.T., Haeusler, A.R., Mistry, N.A., Vidsensky, S., Daley, E.L., Poth, E.M., Hoover, B., Fines, D.M., et al. (2013) RNA toxicity from the ALS/FTD C9ORF72 expansion is mitigated by antisense intervention. *Neuron*, **80**, 415–428.
36. Wainger, B.J., Kiskinis, E., Mellin, C., Wiskow, O., Han, S.S., Sandoe, J., Perez, N.P., Williams, L.A., Lee, S., Boulting, G., et al. (2014) Intrinsic membrane hyperexcitability of amyotrophic lateral sclerosis patient-derived motor neurons. *Cell Rep.*, **7**, 1–11.
37. Chen, H., Qian, K., Du, Z., Cao, J., Petersen, A., Liu, H., Blackburn, L.W.t., Huang, C.L., Errigo, A., Yin, Y., et al. (2014) Modeling ALS with iPSCs reveals that mutant SOD1 misregulates neurofilament balance in motor neurons. *Cell Stem Cell*, **14**, 796–809.
38. Sances, S., Bruijn, L.I., Chandran, S., Eggan, K., Ho, R., Klim, J.R., Livesey, M.R., Lowry, E., Macklis, J.D., Rushton, D., et al. (2016) Modeling ALS with motor neurons derived from human induced pluripotent stem cells. *Nat. Neurosci.*, **16**, 542–553.
39. Juopperi, T.A., Kim, W.R., Chiang, C.H., Yu, H., Margolis, R.L., Ross, C.A., Ming, G.L. and Song, H. (2012) Astrocytes generated from patient induced pluripotent stem cells recapitulate features of Huntington's disease patient cells. *Mol. Brain*, **5**, 17.
40. Zhang, N., An, M.C., Montoro, D. and Ellerby, L.M. (2010) Characterization of Human Huntington's Disease Cell Model from Induced Pluripotent Stem Cells. *PLoS Curr.*, **2**, RRN1193.
41. Park, I.H., Arora, N., Huo, H., Maherali, N., Ahfeldt, T., Shimamura, A., Lensch, M.W., Cowan, C., Hochedlinger, K. and Daley, G.Q. (2008) Disease-specific induced pluripotent stem cells. *Cell*, **134**, 877–886.
42. Jiang, H., Ren, Y., Yuen, E.Y., Zhong, P., Ghaedi, M., Hu, Z., Azabdaftari, G., Nakaso, K., Yan, Z. and Feng, J. (2012) Parkin controls dopamine utilization in human midbrain dopaminergic neurons derived from induced pluripotent stem cells. *Nat. Commun.*, **3**, 668.
43. Sanchez-Danes, A., Richaud-Patin, Y., Carballo-Carbajal, I., Jimenez-Delgado, S., Caig, C., Mora, S., Di Guglielmo, C., Ezquerro, M., Patel, B., Giralt, A., et al. (2012) Disease-specific phenotypes in dopamine neurons from human iPS-based models of genetic and sporadic Parkinson's disease. *EMBO Mol. Med.*, **4**, 380–395.
44. Imaizumi, Y., Okada, Y., Akamatsu, W., Koike, M., Kuzumaki, N., Hayakawa, H., Nihira, T., Kobayashi, T., Ohyama, M., Sato, S., et al. (2012) Mitochondrial dysfunction associated with increased oxidative stress and alpha-synuclein accumulation in PARK2 iPSC-derived neurons and postmortem brain tissue. *Mol. Brain*, **5**, 35.

45. Soldner, F., Laganier, J., Cheng, A.W., Hockemeyer, D., Gao, Q., Alagappan, R., Khurana, V., Golbe, L.I., Myers, R.H., Lindquist, S., et al. (2011) Generation of isogenic pluripotent stem cells differing exclusively at two early onset Parkinson point mutations. *Cell*, **146**, 318–331.
46. Nguyen, H.N., Byers, B., Cord, B., Shcheglovitov, A., Byrne, J., Gujar, P., Kee, K., Schule, B., Dolmetsch, R.E., Langston, W., et al. (2011) LRRK2 mutant iPSC-derived DA neurons demonstrate increased susceptibility to oxidative stress. *Cell Stem Cell*, **8**, 267–280.
47. Seibler, P., Graziotto, J., Jeong, H., Simunovic, F., Klein, C. and Krainc, D. (2011) Mitochondrial Parkin recruitment is impaired in neurons derived from mutant PINK1 induced pluripotent stem cells. *J. Neurosci.*, **31**, 5970–5976.
48. Kondo, T., Asai, M., Tsukita, K., Kutoku, Y., Ohsawa, Y., Sunada, Y., Imamura, K., Egawa, N., Yahata, N., Okita, K., et al. (2013) Modeling Alzheimer's disease with iPSCs reveals stress phenotypes associated with intracellular Abeta and differential drug responsiveness. *Cell Stem Cell*, **12**, 487–496.
49. Israel, M.A., Yuan, S.H., Bardy, C., Reyna, S.M., Mu, Y., Herrera, C., Hefferan, M.P., Van Gorp, S., Nazor, K.L., Boscolo, F.S., et al. (2012) Probing sporadic and familial Alzheimer's disease using induced pluripotent stem cells. *Nature*, **482**, 216–220.
50. Yagi, T., Ito, D., Okada, Y., Akamatsu, W., Nihei, Y., Yoshizaki, T., Yamanaka, S., Okano, H. and Suzuki, N. (2011) Modeling familial Alzheimer's disease with induced pluripotent stem cells. *Hum. Mol. Genet.*, **20**, 4530–4539.
51. Ebert, A.D., Yu, J., Rose, F.F., Jr., Mattis, V.B., Lorson, C.L., Thomson, J.A. and Svendsen, C.N. (2009) Induced pluripotent stem cells from a spinal muscular atrophy patient. *Nature*, **457**, 277–280.
52. Yoshida, M., Kitaoka, S., Egawa, N., Yamane, M., Ikeda, R., Tsukita, K., Amano, N., Watanabe, A., Morimoto, M., Takahashi, J., et al. (2015) Modeling the early phenotype at the neuromuscular junction of spinal muscular atrophy using patient-derived iPSCs. *Stem Cell Reports*, **4**, 561–568.
53. Arbab, M., Baars, S. and Geijsen, N. (2014) Modeling motor neuron disease: the matter of time. *Trends Neurosci.*, **37**, 642–652.
54. Lee, G., Papapetrou, E.P., Kim, H., Chambers, S.M., Tomishima, M.J., Fasano, C.A., Ganat, Y.M., Menon, J., Shimizu, F., Viale, A., et al. (2009) Modelling pathogenesis and treatment of familial dysautonomia using patient-specific iPSCs. *Nature*, **461**, 402–406.
55. Marchetto, M.C., Carromeu, C., Acab, A., Yu, D., Yeo, G.W., Mu, Y., Chen, G., Gage, F.H. and Muotri, A.R. (2010) A model for neural development and treatment of Rett syndrome using human induced pluripotent stem cells. *Cell*, **143**, 527–539.
56. Ricciardi, S., Ungaro, F., Hambrock, M., Rademacher, N., Stefanelli, G., Brambilla, D., Sessa, A., Magagnotti, C., Bachi, A., Giarda, E., et al. (2012) CDKL5 ensures excitatory synapse stability by reinforcing NGL-1-PSD95 interaction in the postsynaptic compartment and is impaired in patient iPSC-derived neurons. *Nat. Cell Biol.*, **14**, 911–923.
57. Cheung, A.Y., Horvath, L.M., Carrel, L. and Ellis, J. (2012) X-chromosome inactivation in rett syndrome human induced pluripotent stem cells. *Front Psychiat.*, **3**, 24.
58. Sheridan, S.D., Theriault, K.M., Reis, S.A., Zhou, F., Madison, J.M., Daheron, L., Loring, J.F. and Haggarty, S.J. (2011) Epigenetic characterization of the FMR1 gene and aberrant neurodevelopment in human induced pluripotent stem cell models of fragile X syndrome. *PLoS One*, **6**, e26203.
59. Liu, J., Koscielska, K.A., Cao, Z., Hulsizer, S., Grace, N., Mitchell, G., Nacey, C., Githinji, J., McGee, J., Garcia-Arocena, D., et al. (2012) Signaling defects in iPSC-derived fragile X pre-mutation neurons. *Hum. Mol. Genet.*, **21**, 3795–3805.
60. Doers, M.E., Musser, M.T., Nichol, R., Berndt, E.R., Baker, M., Gomez, T.M., Zhang, S.C., Abbeduto, L. and Bhattacharyya, A. (2014) iPSC-derived forebrain neurons from FXS individuals show defects in initial neurite outgrowth. *Stem Cells Dev.*, **23**, 1777–1787.
61. Pasca, S.P., Portmann, T., Voineagu, I., Yazawa, M., Shcheglovitov, A., Pasca, A.M., Cord, B., Palmer, T.D., Chikahisa, S., Nishino, S., et al. (2011) Using iPSC-derived neurons to uncover cellular phenotypes associated with Timothy syndrome. *Nat. Med.*, **17**, 1657–1662.
62. Tian, Y., Voineagu, I., Pasca, S.P., Won, H., Chandran, V., Horvath, S., Dolmetsch, R.E. and Geschwind, D.H. (2014) Alteration in basal and depolarization induced transcriptional network in iPSC derived neurons from Timothy syndrome. *Genome Med.*, **6**, 75.
63. Liu, Y., Lopez-Santiago, L.F., Yuan, Y., Jones, J.M., Zhang, H., O'Malley, H.A., Patino, G.A., O'Brien, J.E., Rusconi, R., Gupta, A., et al. (2013) Dravet syndrome patient-derived neurons suggest a novel epilepsy mechanism. *Ann. Neurol.*, **74**, 128–139.
64. Jiao, J., Yang, Y., Shi, Y., Chen, J., Gao, R., Fan, Y., Yao, H., Liao, W., Sun, X.F. and Gao, S. (2013) Modeling Dravet syndrome using induced pluripotent stem cells (iPSCs) and directly converted neurons. *Hum. Mol. Genet.*, **22**, 4241–4252.
65. Chamberlain, S.J., Chen, P.F., Ng, K.Y., Bourgeois-Rocha, F., Lemtiri-Chlieh, F., Levine, E.S. and Lalande, M. (2010) Induced pluripotent stem cell models of the genomic imprinting disorders Angelman and Prader-Willi syndromes. *Proc. Natl Acad. Sci. U S A.*, **107**, 17668–17673.
66. Stelzer, Y., Sagi, I., Yanuka, O., Eiges, R. and Benvenisty, N. (2014) The noncoding RNA IPW regulates the imprinted DLK1-DIO3 locus in an induced pluripotent stem cell model of Prader-Willi syndrome. *Nat. Genet.*, **46**, 551–557.
67. Yang, J., Cai, J., Zhang, Y., Wang, X., Li, W., Xu, J., Li, F., Guo, X., Deng, K., Zhong, M., et al. (2010) Induced pluripotent stem cells can be used to model the genomic imprinting disorder Prader-Willi syndrome. *J. Biol. Chem.*, **285**, 40303–40311.
68. Germain, N.D., Chen, P.F., Plocik, A.M., Glatt-Deeley, H., Brown, J., Fink, J.J., Bolduc, K.A., Robinson, T.M., Levine, E.S., Reiter, L.T., et al. (2014) Gene expression analysis of human induced pluripotent stem cell-derived neurons carrying copy number variants of chromosome 15q11-q13.1. *Mol. Autism*, **5**, 44.
69. Das, D.K., Tapias, V., D'Aiuto, L., Chowdari, K.V., Francis, L., Zhi, Y., Ghosh, B.A., Surti, U., Tischfield, J., Sheldon, M., et al. (2015) Genetic and morphological features of human iPSC-derived neurons with chromosome 15q11.2 (BP1-BP2) deletions. *Mol. Neuropsychiatry*, **1**, 116–123.
70. Shcheglovitov, A., Shcheglovitova, O., Yazawa, M., Portmann, T., Shu, R., Sebastiano, V., Krawisz, A., Froehlich, W., Bernstein, J.A., Hallmayer, J.F., et al. (2013) SHANK3 and IGF1 restore synaptic deficits in neurons from 22q13 deletion syndrome patients. *Nature*, **503**, 267–271.
71. Brennand, K., Savas, J.N., Kim, Y., Tran, N., Simone, A., Hashimoto-Torii, K., Beaumont, K.G., Kim, H.J., Topol, A., Ladrón, I., et al. (2015) Phenotypic differences in hiPSC NPCs derived from patients with schizophrenia. *Mol. Psychiatry*, **20**, 361–8.

72. Murai, K., Sun, G., Ye, P., Tian, E., Yang, S., Cui, Q., Sun, G., Trinh, D., Sun, O., Hong, T., et al. (2016) The TLX-miR-219 cascade regulates neural stem cell proliferation in neurodevelopment and schizophrenia iPSC model. *Nat. Commun.*, **7**, 10965.
73. Topol, A., English, J.A., Flaherty, E., Rajarajan, P., Hartley, B.J., Gupta, S., Desland, F., Zhu, S., Goff, T., Friedman, L., et al. (2015) Increased abundance of translation machinery in stem cell-derived neural progenitor cells from four schizophrenia patients. *Transl Psychiatry*, **5**, e662.
74. Topol, A., Zhu, S., Tran, N., Simone, A., Fang, G. and Brennand, K.J. (2015) Altered WNT Signaling in Human Induced Pluripotent Stem Cell Neural Progenitor Cells Derived from Four Schizophrenia Patients. *Biol Psychiatry*, **78**, e29–e34.
75. Hook, V., Brennand, K.J., Kim, Y., Toneff, T., Funkelstein, L., Lee, K.C., Ziegler, M. and Gage, F.H. (2014) Human iPSC neurons display activity-dependent neurotransmitter secretion: aberrant catecholamine levels in schizophrenia neurons. *Stem Cell Reports*, **3**, 531–538.
76. Pedrosa, E., Sandler, V., Shah, A., Carroll, R., Chang, C., Rockowitz, S., Guo, X., Zheng, D. and Lachman, H.M. (2011) Development of patient-specific neurons in schizophrenia using induced pluripotent stem cells. *J Neurogenet.*, **25**, 88–103.
77. Madison, J.M., Zhou, F., Nigam, A., Hussain, A., Barker, D.D., Nehme, R., van der Ven, K., Hsu, J., Wolf, P., Fleishman, M., et al. (2015) Characterization of bipolar disorder patient-specific induced pluripotent stem cells from a family reveals neurodevelopmental and mRNA expression abnormalities. *Mol. Psychiatry*, **20**, 703–717.
78. Mertens, J., Wang, Q.W., Kim, Y., Yu, D.X., Pham, S., Yang, B., Zheng, Y., Diffenderfer, K.E., Zhang, J., Soltani, S., et al. (2015) Differential responses to lithium in hyperexcitable neurons from patients with bipolar disorder. *Nature*, **527**, 95–99.
79. Mariani, J., Coppola, G., Zhang, P., Abyzov, A., Provini, L., Tomasini, L., Amenduni, M., Szekely, A., Palejev, D., Wilson, M., et al. (2015) FOXG1-Dependent Dysregulation of GABA/Glutamate Neuron Differentiation in Autism Spectrum Disorders. *Cell*, **162**, 375–390.
80. Griesi-Oliveira, K., Acab, A., Gupta, A.R., Sunaga, D.Y., Chailangkarn, T., Nicol, X., Nunez, Y., Walker, M.F., Murdoch, J.D., Sanders, S.J., et al. (2015) Modeling non-syndromic autism and the impact of TRPC6 disruption in human neurons. *Mol. Psychiatry*, **20**, 1350–1365.
81. Mishra, H.K., Prots, I., Havlicek, S., Kohl, Z., Perez-Branguli, F., Boerstler, T., Anneser, L., Minakaki, G., Wend, H., Hampl, M., et al. (2016) GSK3ss-dependent dysregulation of neurodevelopment in SPG11-patient iPSC model. *Ann. Neurol.*, doi: 10.1002/ana.24633.
82. Zhu, P.P., Denton, K.R., Pierson, T.M., Li, X.J. and Blackstone, C. (2014) Pharmacologic rescue of axon growth defects in a human iPSC model of hereditary spastic paraplegia SPG3A. *Hum. Mol. Genet.*, **23**, 5638–5648.
83. Denton, K.R., Lei, L., Grenier, J., Rodionov, V., Blackstone, C. and Li, X.J. (2014) Loss of spastin function results in disease-specific axonal defects in human pluripotent stem cell-based models of hereditary spastic paraplegia. *Stem Cells*, **32**, 414–423.
84. Han, L., Li, Y., Tcho, J., Kaplan, A.D., Lin, B., Li, Y., Mich-Basso, J., Lis, A., Hassan, N., London, B., et al. (2014) Study familial hypertrophic cardiomyopathy using patient-specific induced pluripotent stem cells. *Cardiovasc. Res.*, **104**, 258–269.
85. Tanaka, A., Yuasa, S., Mearini, G., Egashira, T., Seki, T., Kodaira, M., Kusumoto, D., Kuroda, Y., Okata, S., Suzuki, T., et al. (2014) Endothelin-1 induces myofibrillar disarray and contractile vector variability in hypertrophic cardiomyopathy-induced pluripotent stem cell-derived cardiomyocytes. *J. Am. Heart. Assoc.*, **3**, e001263.
86. Lan, F., Lee, A.S., Liang, P., Sanchez-Freire, V., Nguyen, P.K., Wang, L., Han, L., Yen, M., Wang, Y., Sun, N., et al. (2013) Abnormal calcium handling properties underlie familial hypertrophic cardiomyopathy pathology in patient-specific induced pluripotent stem cells. *Cell Stem Cell*, **12**, 101–113.
87. Sun, N., Yazawa, M., Liu, J., Han, L., Sanchez-Freire, V., Abilez, O.J., Navarrete, E.G., Hu, S., Wang, L., Lee, A., et al. (2012) Patient-specific induced pluripotent stem cells as a model for familial dilated cardiomyopathy. *Sci. Transl Med.*, **4**, 130ra147.
88. Tse, H.F., Ho, J.C., Choi, S.W., Lee, Y.K., Butler, A.W., Ng, K.M., Siu, C.W., Simpson, M.A., Lai, W.H., Chan, Y.C., et al. (2013) Patient-specific induced-pluripotent stem cells-derived cardiomyocytes recapitulate the pathogenic phenotypes of dilated cardiomyopathy due to a novel DES mutation identified by whole exome sequencing. *Hum. Mol. Genet.*, **22**, 1395–1403.
89. Hinson, J.T., Chopra, A., Nafissi, N., Polacheck, W.J., Benson, C.C., Swist, S., Gorham, J., Yang, L., Schafer, S., Sheng, C.C., et al. (2015) HEART DISEASE. Titin mutations in iPSC cells define sarcomere insufficiency as a cause of dilated cardiomyopathy. *Science*, **349**, 982–986.
90. Burridge, P.W., Diecke, S., Matsa, E., Sharma, A., Wu, H. and Wu, J.C. (2016) Modeling Cardiovascular Diseases with Patient-Specific Human Pluripotent Stem Cell-Derived Cardiomyocytes. *Methods Mol. Biol.*, **1353**, 119–130.
91. Wyles, S.P., Li, X., Hrstka, S.C., Reyes, S., Oommen, S., Beraldi, R., Edwards, J., Terzic, A., Olson, T.M. and Nelson, T.J. (2016) Modeling structural and functional deficiencies of RBM20 familial dilated cardiomyopathy using human induced pluripotent stem cells. *Hum. Mol. Genet.*, **25**, 254–265.
92. Caspi, O., Huber, I., Gepstein, A., Arbel, G., Maizels, L., Boulos, M. and Gepstein, L. (2013) Modeling of arrhythmogenic right ventricular cardiomyopathy with human induced pluripotent stem cells. *Circ. Cardiovasc. Genet.*, **6**, 557–568.
93. Ma, D., Wei, H., Lu, J., Ho, S., Zhang, G., Sun, X., Oh, Y., Tan, S.H., Ng, M.L., Shim, W., et al. (2013) Generation of patient-specific induced pluripotent stem cell-derived cardiomyocytes as a cellular model of arrhythmogenic right ventricular cardiomyopathy. *Eur. Heart J.*, **34**, 1122–1133.
94. Kim, C., Wong, J., Wen, J., Wang, S., Wang, C., Spiering, S., Kan, N.G., Forcales, S., Puri, P.L., Leone, T.C., et al. (2013) Studying arrhythmogenic right ventricular dysplasia with patient-specific iPSCs. *Nature*, **494**, 105–110.
95. Lin, B., Li, Y., Han, L., Kaplan, A.D., Ao, Y., Kalra, S., Bett, G.C., Rasmusson, R.L., Denning, C. and Yang, L. (2015) Modeling and study of the mechanism of dilated cardiomyopathy using induced pluripotent stem cells derived from individuals with Duchenne muscular dystrophy. *Dis. Model. Mech.*, **8**, 457–466.
96. Si-Tayeb, K., Idriss, S., Champon, B., Caillaud, A., Pichelin, M., Arnaud, L., Lemarchand, P.L., May, C., Zibara, K. and Cariou, B., (2016) Urine-sample-derived human induced pluripotent stem cells as a model to study PCSK9-mediated

- autosomal dominant hypercholesterolemia. *Dis. Model Mech.*, **9**, 81–90.
97. Fattahi, F., Asgari, S., Pournasr, B., Seifinejad, A., Totonchi, M., Taei, A., Aghdami, N., Salekdeh, G.H. and Baharvand, H. (2013) Disease-corrected hepatocyte-like cells from familial hypercholesterolemia-induced pluripotent stem cells. *Mol. Biotechnol.*, **54**, 863–873.
 98. Cayo, M.A., Cai, J., DeLaForest, A., Noto, F.K., Nagaoka, M., Clark, B.S., Coltery, R.F., Si-Tayeb, K. and Duncan, S.A. (2012) JD induced pluripotent stem cell-derived hepatocytes faithfully recapitulate the pathophysiology of familial hypercholesterolemia. *Hepatology*, **56**, 2163–2171.
 99. Camnasio, S., Delli Carri, A., Lombardo, A., Grad, I., Mariotti, C., Castucci, A., Rozell, B., Lo Riso, P., Castiglioni, V., Zuccato, C., et al. (2012) The first reported generation of several induced pluripotent stem cell lines from homozygous and heterozygous Huntington's disease patients demonstrates mutation related enhanced lysosomal activity. *Neurobiol. Dis.*, **46**, 41–51.
 100. Consortium, H.D.i. (2012) Induced pluripotent stem cells from patients with Huntington's disease show CAG-repeat-expansion-associated phenotypes. *Cell Stem Cell*, **11**, 264–278.
 101. Farra, N., Zhang, W.B., Pasceri, P., Eubanks, J.H., Salter, M.W. and Ellis, J. (2012) Rett syndrome induced pluripotent stem cell-derived neurons reveal novel neurophysiological alterations. *Mol. Psychiatry*, **17**, 1261–1271.
 102. Martins-Taylor, K., Hsiao, J.S., Chen, P.F., Glatt-Deeley, H., De Smith, A.J., Blakemore, A.I., Lalande, M. and Chamberlain, S.J. (2014) Imprinted expression of UBE3A in non-neuronal cells from a Prader-Willi syndrome patient with an atypical deletion. *Hum. Mol. Genet.*, **23**, 2364–2373.
 103. Yazawa, M., Hsueh, B., Jia, X., Pasca, A.M., Bernstein, J.A., Hallmayer, J. and Dolmetsch, R.E. (2011) Using induced pluripotent stem cells to investigate cardiac phenotypes in Timothy syndrome. *Nature*, **471**, 230–234.
 104. Brennand, K., Savas, J.N., Kim, Y., Tran, N., Simone, A., Hashimoto-Torii, K., Beaumont, K.G., Kim, H.J., Topol, A., Ladrán, I., et al. (2015) Phenotypic differences in hiPSC NPCs derived from patients with schizophrenia. *Mol Psychiatry*, **20**, 361–368.
 105. Bellin, M., Casini, S., Davis, R.P., D'Aniello, C., Haas, J., Ward-van Oostwaard, D., Tertoolen, L.G., Jung, C.B., Elliott, D.A., Welling, A., et al. (2013) Isogenic human pluripotent stem cell pairs reveal the role of a KCNH2 mutation in long-QT syndrome. *Embo J.*, **32**, 3161–3175.
 106. Moretti, A., Bellin, M., Welling, A., Jung, C.B., Lam, J.T., Bott-Flugel, L., Dorn, T., Goedel, A., Hohnke, C., Hofmann, F., et al. (2010) Patient-Specific Induced Pluripotent Stem-Cell Models for Long-QT Syndrome. *N. Engl J. Med.*, **6**, 407–411.
 107. Itzhaki, I., Maizels, L., Huber, I., Zwi-Dantsis, L., Caspi, O., Winterstern, A., Feldman, O., Gepstein, A., Arbel, G., Hammerman, H., et al. (2011) Modelling the long QT syndrome with induced pluripotent stem cells. *Nature*, **471**, 225–229.
 108. Matsa, E., Rajamohan, D., Dick, E., Young, L., Mellor, I., Staniforth, A. and Denning, C. (2011) Drug evaluation in cardiomyocytes derived from human induced pluripotent stem cells carrying a long QT syndrome type 2 mutation. *Eur. Heart J.*, **32**, 952–962.
 109. Lahti, A.L., Kujala, V.J., Chapman, H., Koivisto, A.P., Pekkanen-Mattila, M., Kerkela, E., Hyttinen, J., Kontula, K., Swan, H., Conklin, B.R., et al. (2012) Model for long QT syndrome type 2 using human iPSCs demonstrates arrhythmogenic characteristics in cell culture. *Dis. Model Mech.*, **5**, 220–230.
 110. Egashira, T., Yuasa, S., Suzuki, T., Aizawa, Y., Yamakawa, H., Matsushashi, T., Ohno, Y., Tohyama, S., Okata, S., Seki, T., et al. (2012) Disease characterization using LQTS-specific induced pluripotent stem cells. *Cardiovasc. Res.*, **95**, 419–429.
 111. Jung, C.B., Moretti, A., Mederos y Schnitzler, M., Iop, L., Storch, U., Bellin, M., Dorn, T., Ruppenthal, S., Pfeiffer, S., Goedel, A., et al. (2012) Dantrolene rescues arrhythmogenic RYR2 defect in a patient-specific stem cell model of catecholaminergic polymorphic ventricular tachycardia. *EMBO Mol. Med.*, **4**, 180–191.
 112. Kujala, K., Paavola, J., Lahti, A., Larsson, K., Pekkanen-Mattila, M., Viitasalo, M., Lahtinen, A.M., Toivonen, L., Kontula, K., Swan, H., et al. (2012) Cell model of catecholaminergic polymorphic ventricular tachycardia reveals early and delayed afterdepolarizations. *PLoS One*, **7**, e44660.
 113. Wang, G., McCain, M.L., Yang, L., He, A., Pasqualini, F.S., Agarwal, A., Yuan, H., Jiang, D., Zhang, D., Zangi, L., et al. (2014) Modeling the mitochondrial cardiomyopathy of Barth syndrome with induced pluripotent stem cell and heart-on-chip technologies. *Nat. Med.*, **20**, 616–623.
 114. Hick, A., Wattenhofer-Donze, M., Chintawar, S., Tropel, P., Simard, J.P., Vaucamps, N., Gall, D., Lambot, L., Andre, C., Reutenauer, L., et al. (2013) Neurons and cardiomyocytes derived from induced pluripotent stem cells as a model for mitochondrial defects in Friedreich's ataxia. *Dis. Model Mech.*, **6**, 608–621.
 115. Yasuno, T., Osafune, K., Sakurai, H., Asaka, I., Tanaka, A., Yamaguchi, S., Yamada, K., Hitomi, H., Arai, S., Kurose, Y., et al. (2014) Functional analysis of iPSC-derived myocytes from a patient with carnitine palmitoyltransferase II deficiency. *Biochem. Biophys. Res. Commun.*, **448**, 175–181.
 116. Carvajal-Vergara, X., Sevilla, A., D'Souza, S.L., Ang, Y.S., Schaniel, C., Lee, D.F., Yang, L., Kaplan, A.D., Adler, E.D., Rozov, R., et al. (2010) Patient-specific induced pluripotent stem-cell-derived models of LEOPARD syndrome. *Nature*, **465**, 808–812.
 117. Sato, Y., Kobayashi, H., Higuchi, T., Shimada, Y., Era, T., Kimura, S., Eto, Y., Ida, H. and Ohashi, T. (2015) Disease modeling and lentiviral gene transfer in patient-specific induced pluripotent stem cells from late-onset Pompe disease patient. *Mol. Ther. Methods Clin. Dev.*, **2**, 15023.
 118. Raval, K.K., Tao, R., White, B.E., De Lange, W.J., Koonce, C.H., Yu, J., Kishnani, P.S., Thomson, J.A., Mosher, D.F., Ralphe, J.C., et al. (2015) Pompe disease results in a Golgi-based glycosylation deficit in human induced pluripotent stem cell-derived cardiomyocytes. *J. Biol. Chem.*, **290**, 3121–3136.
 119. Huang, H.P., Chen, P.H., Hwu, W.L., Chuang, C.Y., Chien, Y.H., Stone, L., Chien, C.L., Li, L.T., Chiang, S.C., Chen, H.F., et al. (2011) Human Pompe disease-induced pluripotent stem cells for pathogenesis modeling, drug testing and disease marker identification. *Hum. Mol. Genet.*, **20**, 4851–4864.
 120. Higuchi, T., Kawagoe, S., Otsu, M., Shimada, Y., Kobayashi, H., Hirayama, R., Eto, K., Ida, H., Ohashi, T., Nakauchi, H., et al. (2014) The generation of induced pluripotent stem cells (iPSCs) from patients with infantile and late-onset types of Pompe disease and the effects of treatment with acid-alpha-glucosidase in Pompe's iPSCs. *Mol. Genet. Metab.*, **112**, 44–48.
 121. Siu, C.W., Lee, Y.K., Ho, J.C., Lai, W.H., Chan, Y.C., Ng, K.M., Wong, L.Y., Au, K.W., Lau, Y.M., Zhang, J., et al. (2012) Modeling of lamin A/C mutation premature cardiac aging using patient-specific induced pluripotent stem cells. *Aging (Albany NY)*, **4**, 803–822.