

A Review of the Complex Logarithm

Li Yueheng, s2306706

The University of Edinburgh

13th February 2022

- 1 Definition and basic properties
- 2 Branches of the logarithm and branch cuts
- 3 Holomorphicity gained and continuity lost

Definition of the complex logarithm

Definition (1.7.1 on p.16, Gratwick)

For $z \in \mathbb{C} - \{0\}$, define its *logarithm* by

$$\log(z) := \{w \in \mathbb{C} : \exp(w) = z\}$$

Definition of the complex logarithm

Definition (1.7.1 on p.16, Gratwick)

For $z \in \mathbb{C} - \{0\}$, define its *logarithm* by

$$\log(z) := \{w \in \mathbb{C} : \exp(w) = z\}$$

Remark

The complex logarithm is a multivalued function. We call any element of the set $\log(z)$ a logarithm of z .

Basic properties of the complex logarithm

Proposition (1.7.3 on p.16, Gratwick)

Let $z, w \in \mathbb{C} - \{0\}$, and $z = re^{i\theta}$ in exponential form. Then

$$\textcircled{1} \log(z) = \{\ln r + i\theta + i2\pi k : k \in \mathbb{Z}\} \quad (\text{Depiction of the set } \log(z))$$

Basic properties of the complex logarithm

Proposition (1.7.3 on p.16, Gratwick)

Let $z, w \in \mathbb{C} - \{0\}$, and $z = re^{i\theta}$ in exponential form. Then

- ① $\log(z) = \{\ln r + i\theta + i2\pi k : k \in \mathbb{Z}\}$ (Depiction of the set $\log(z)$)
- ② $\log(zw) = \log(z) + \log(w)$ (Law of multiplication)

Basic properties of the complex logarithm

Proposition (1.7.3 on p.16, Gratwick)

Let $z, w \in \mathbb{C} - \{0\}$, and $z = re^{i\theta}$ in exponential form. Then

- ① $\log(z) = \{\ln r + i\theta + i2\pi k : k \in \mathbb{Z}\}$ (Depiction of the set $\log(z)$)
- ② $\log(zw) = \log(z) + \log(w)$ (Law of multiplication)
- ③ $\log(1/z) = -\log(z)$ (Law of reciprocal)

Basic properties of the complex logarithm

Proposition (1.7.3 on p.16, Gratwick)

Let $z, w \in \mathbb{C} - \{0\}$, and $z = re^{i\theta}$ in exponential form. Then

- ① $\log(z) = \{\ln r + i\theta + i2\pi k : k \in \mathbb{Z}\}$ (Depiction of the set $\log(z)$)
- ② $\log(zw) = \log(z) + \log(w)$ (Law of multiplication)
- ③ $\log(1/z) = -\log(z)$ (Law of reciprocal)

Remark

These statements should be understood as equalities between sets.

Sketch of proof

1. Let $w = a + ib$ and solve the equation

$$\exp(w) = \exp(a + ib) = re^{i\theta} = z$$

to find all $w \in \log(z)$.

Sketch of proof

1. Let $w = a + ib$ and solve the equation

$$\exp(w) = \exp(a + ib) = re^{i\theta} = z$$

to find all $w \in \log(z)$.

2. The law of multiplication follows directly from our depiction of the logarithm.

Sketch of proof

1. Let $w = a + ib$ and solve the equation

$$\exp(w) = \exp(a + ib) = re^{i\theta} = z$$

to find all $w \in \log(z)$.

2. The law of multiplication follows directly from our depiction of the logarithm.
3. Notice that $\log(1) = \{i2\pi k : k \in \mathbb{Z}\}$ and use the law of multiplication.

Branches of the logarithm

Definition (1.7.9 on p.17, Gratwick)

A *branch* of the logarithm is defined as a function $\text{Log}_\phi : \mathbb{C} - \{0\} \rightarrow \mathbb{C}$,

$$\text{Log}_\phi(z) := \ln |z| + i\text{Arg}_\phi(z)$$

where $\text{Arg}_\phi(z) \in \arg(z) \cap (\phi, \phi + 2\pi]$ is a properly defined real number.

Branches of the logarithm

Definition (1.7.9 on p.17, Gratwick)

A *branch* of the logarithm is defined as a function $\text{Log}_\phi : \mathbb{C} - \{0\} \rightarrow \mathbb{C}$,

$$\text{Log}_\phi(z) := \ln |z| + i\text{Arg}_\phi(z)$$

where $\text{Arg}_\phi(z) \in \arg(z) \cap (\phi, \phi + 2\pi]$ is a properly defined real number.

Example

The *principal branch* of the logarithm is the function $\text{Log} : \mathbb{C} - \{0\} \rightarrow \mathbb{C}$ defined as

$$\text{Log}(z) := \text{Log}_{-\pi}(z) = \ln |z| + i\text{Arg}(z)$$

Branch cuts and the cut plane

Definition (1.7.7 on p.17, Gratwick)

A *branch cut* L is a subset removed from the complex plane \mathbb{C} so that a holomorphic branch of a multivalued function may be defined on $D = \mathbb{C} - L$, the cut plane.

Branch cuts and the cut plane

Definition (1.7.7 on p.17, Gratwick)

A *branch cut* L is a subset removed from the complex plane \mathbb{C} so that a holomorphic branch of a multivalued function may be defined on $D = \mathbb{C} - L$, the cut plane.

Remark

Half-lines are an important class of branch cuts, so we introduce the following notation: For $z_0 \in \mathbb{C}$ and $\phi \in \mathbb{R}$, define

$$L_{z_0, \phi} = \{z \in \mathbb{C} : z = z_0 + re^{i\phi}, r \geq 0\}$$

and define its cut plane $D_{z_0, \phi} := \mathbb{C} - L_{z_0, \phi}$. In particular, $D_\phi := D_{0, \phi}$.

Holomorphicity gained

Theorem (1.7.10 on p.18, Gratwick)

A brunch of the logarithm Log_ϕ is holomorphic on the cut plane D_ϕ , and $\text{Log}'_\phi(z) = 1/z$ in D_ϕ .

Holomorphicity gained

Theorem (1.7.10 on p.18, Gratwick)

A brunch of the logarithm Log_ϕ is holomorphic on the cut plane D_ϕ , and $\text{Log}'_\phi(z) = 1/z$ in D_ϕ .

Proof.

A sketch:

1. Verify that $\text{Log}_\phi(z) = \ln(|z|) + i\text{Arg}_\phi(z)$ has continuously differentiable real and imaginary parts on D_ϕ .

Holomorphicity gained

Theorem (1.7.10 on p.18, Gratwick)

A branch of the logarithm Log_ϕ is holomorphic on the cut plane D_ϕ , and $\text{Log}'_\phi(z) = 1/z$ in D_ϕ .

Proof.

A sketch:

1. Verify that $\text{Log}_\phi(z) = \ln(|z|) + i\text{Arg}_\phi(z)$ has continuously differentiable real and imaginary parts on D_ϕ .
2. Check that the Cauchy-Riemann equations are satisfied on D_ϕ . These two steps combined show that Log_ϕ is holomorphic on D_ϕ .

Holomorphicity gained

Theorem (1.7.10 on p.18, Gratwick)

A branch of the logarithm Log_ϕ is holomorphic on the cut plane D_ϕ , and $\text{Log}'_\phi(z) = 1/z$ in D_ϕ .

Proof.

A sketch:

1. Verify that $\text{Log}_\phi(z) = \ln(|z|) + i\text{Arg}_\phi(z)$ has continuously differentiable real and imaginary parts on D_ϕ .
2. Check that the Cauchy-Riemann equations are satisfied on D_ϕ . These two steps combined show that Log_ϕ is holomorphic on D_ϕ .
3. Differentiate $z = \exp(\text{Log}_\phi(z))$ using the chain rule to compute the derivative of Log_ϕ on D_ϕ .



Continuity lost

Since $\text{Arg}_\phi(z)$ takes a jump when z crosses $L_{0,\phi}$, it isn't continuous there, and hence isn't $\text{Log}_\phi(z) = \ln(|z|) + i\text{Arg}_\phi(z)$.

Continuity lost

Since $\text{Arg}_\phi(z)$ takes a jump when z crosses $L_{0,\phi}$, it isn't continuous there, and hence isn't $\text{Log}_\phi(z) = \ln(|z|) + i\text{Arg}_\phi(z)$.

Example

Notice that

$$\text{Arg}_0(1) = 2\pi$$

while

$$\text{Arg}_0(e^{i\varepsilon}) = \varepsilon$$

Thus $2\pi = \text{Arg}_0(1) = \text{Arg}_0(\lim_{\varepsilon \rightarrow 0} e^{i\varepsilon}) \neq \lim_{\varepsilon \rightarrow 0} \text{Arg}_0(e^{i\varepsilon}) = 0$

To conclude

- 1 The complex logarithm is a multivalued function.

To conclude

- ① The complex logarithm is a multivalued function.
- ② The complex logarithm "inherits" some properties of the real one, but just formally!

To conclude

- 1 The complex logarithm is a multivalued function.
- 2 The complex logarithm "inherits" some properties of the real one, but just formally!
- 3 By cutting the complex plane, we may define branches of the logarithm, which are *functions*.

To conclude

- ① The complex logarithm is a multivalued function.
- ② The complex logarithm "inherits" some properties of the real one, but just formally!
- ③ By cutting the complex plane, we may define branches of the logarithm, which are *functions*.
- ④ Branches lose continuity on the branch cuts, but gain holomorphicity elsewhere.

To conclude

- ① The complex logarithm is a multivalued function.
- ② The complex logarithm "inherits" some properties of the real one, but just formally!
- ③ By cutting the complex plane, we may define branches of the logarithm, which are *functions*.
- ④ Branches lose continuity on the branch cuts, but gain holomorphicity elsewhere.

1. Gratwick, R. (2022). Honours Complex Variables Lecture Notes 2021–2022. Accessed February 13th, 2022.