

# **Tsiatis (2006) Chapters 10–11**

**Asymptotic Statistics 2025 Summer Reading Group**

---

Naoki Eguchi

August 18, 2025

Faculty of Medicine, Kyoto University

# Today's agenda

- Recap: Notation, Coarsened data, IF geometry, AIPWCC
- Chapter 10: Two-level missingness  $\Rightarrow$  efficient AIPWCC & DR
- *Skip (brief mention only)*: monotone coarsening details, censoring
- Generalization idea (operators  $L$ ,  $M$ ,  $M^{-1}$ ) for later use
- Chapter 11: (two representations; how to compute)
- Implementation scheme

## Recap

---

## Coarsened data and Semiparametric statistics

- Full data  $Z$ ; observed data  $\{C, G_C(Z)\}$  under CAR (coarsening at random).  
 $\Rightarrow \forall r, P(C = r \mid Z) = \pi\{r, G_r(Z)\}$  (coarsening depends only on observed data.)
- When  $C = \infty$ , the data are completely observed (equal to full data).
- In semiparametric statistics, we study the influence function, an element of  $\Lambda^\perp$ .
  - Full-data tangent space:  $\mathcal{H}^F$ ; full-data nuisance tangent space:  $\Lambda_F \subset \mathcal{H}^F$ .
    - (Observed) tangent space:  $\mathcal{H}$ ; nuisance tangent space:  $\Lambda = \Lambda_\psi \oplus \Lambda_\eta$  ( $\Lambda_\psi \perp \Lambda_\eta$ ).
- **Theorem 8.3:** all observed-data influence functions can be written as

$$\varphi\{C, G_C(Z)\} = \left[ \frac{\mathbf{1}(C = \infty)}{\pi(\infty, Z, \psi_0)} \varphi^F(Z) + L_2\{C, G_C(Z)\} \right] - \Pi([\cdot] \mid \Lambda_\psi)$$

where  $\varphi^F(Z)$  : full-data IF,  $L_2\{C, G_C(Z)\} \in \Lambda_2$  (augmentation space)

## **10. Improving Efficiency & Double Robustness with Coarsened Data**

---

# The optimal (variance-minimizing) observed-data IF

## Theorem 10.1

- For fixed full-data IF  $\varphi_F(Z) \in (\text{IF})_F$ , the optimal choice is

$$L_2\{C, G_C(Z)\} = -\Pi \left[ \frac{\mathbf{1}(C = \infty) \varphi_F(Z)}{\pi(\infty, Z, \psi_0)} \mid \Lambda_2 \right].$$

- Hence the optimal observed-data influence function is given by

$$\frac{\mathbf{1}(C = \infty) \varphi_F(Z)}{\pi(\infty, Z, \psi_0)} - \Pi \left[ \frac{\mathbf{1}(C = \infty) \varphi_F(Z)}{\pi(\infty, Z, \psi_0)} \mid \Lambda_2 \right].$$

- *Remark 1* : If we **know**  $\psi_0$ , we can choose optimal IF.

Also, we do not have to estimate  $\psi$  because  $\Lambda_\psi \subset \Lambda_2$  (already subtracted !)

## Generalized by linear operator

### Definition 1

- Define the linear operator  $\mathcal{J} : \mathcal{H}^F \rightarrow \mathcal{H}$

$$h_F \mapsto \mathcal{J}(h_F) = \frac{\mathbf{1}(C = \infty)}{\pi(\infty, Z, \psi_0)} h_F(Z) - \Pi \left[ \frac{\mathbf{1}(C = \infty)}{\pi(\infty, Z, \psi_0)} h_F(Z) \mid \Lambda_2 \right].$$

- This operator turns full-data IF into efficient observed-data IF, so we can create efficient observed-data IF space.

### Definition 2

- The class of *double-robust* observed-data influence functions is defined by

$$(\text{IF})_{\text{DR}} = \{ \mathcal{J}(\varphi_F) : \varphi_F(Z) \in (\text{IF})_F \}.$$

- $(\text{IF})^F = \varphi^F(Z) + \mathcal{T}^{F\perp}$  is a linear variety in  $\mathcal{H}^F$ .
- Since  $\mathcal{J}$  is **linear** operator,  $(\text{IF})_{\text{DR}} = \mathcal{J}\{(\text{IF})^F\} = \mathcal{J}(\varphi^F) + \mathcal{J}(\mathcal{T}^{F\perp})$  in  $\mathcal{H}$ .

### Definition 3

- The linear subspace

$$\mathcal{J}(\Lambda^{F\perp}) \subset \Lambda^\perp \subset \mathcal{H},$$

is called the **DR linear space**.

- Explicitly,

$$\mathcal{J}(\Lambda^{F\perp}) = \left\{ \mathcal{J}(\varphi^F) : \varphi^F(Z) \in \Lambda^{F\perp} \right\},$$



## **10.2 Improving Efficiency with Two Levels of Missingness**

---

## Set-up: Two-level missingness

- Partition  $Z = (Z_1^\top, Z_2^\top)^\top$  with  $Z_1$  always observed and  $Z_2$  sometimes missing.
- Complete-case indicator  $R \in \{0, 1\}$  with MAR:  
 $P(R = 1 \mid Z) = P(R = 1 \mid Z_1) = \pi(Z_1, \psi)$  (e.g., logit model).
- Observed data:  $O = (R, Z_1, RZ_2)$ .
- In this case,  $\frac{I(C = \infty) \varphi^{*F}(Z)}{\varpi(\infty, Z)} = \frac{R\varphi^{*F}(Z)}{\pi(Z_1)}$  for  $\varphi^{*F}(Z) = m(Z, \beta_0) \in \Lambda^{F\perp}$ ,  
hence, we want to find its projection onto  $\Lambda_2$ .
- Also,  $\Lambda_2 = \left\{ \frac{R - \pi(Z_1)}{\pi(Z_1)} h_2(Z_1, \beta) : \forall h_2 : Z_1 \rightarrow \mathbb{R}^q \right\}$

# Finding the Projection onto the Augmentation space

## Theorem 10.2

- The projection of

$$\frac{R \varphi^{*F}(Z)}{\pi(Z_1)}$$

onto the augmentation space  $\Lambda_2$  is the unique element

$$\left( \frac{R - \pi(Z_1)}{\pi(Z_1)} \right) h_2^0(Z_1) \in \Lambda_2,$$

where

$$h_2^0(Z_1) = \mathbb{E}[\varphi^{*F}(Z) \mid Z_1].$$

- For implementation, we just compute  $\mathbb{E}[\varphi^{*F}(Z) \mid Z_1]$ .

## Adaptive estimation and its algorithm

- Consider the case where we can **choose** an estimating function  $m(Z, \beta)$ .
- **Optimal** augmentation:  $h_2^*(Z_1, \beta) = \mathbb{E}[m(Z, \beta) \mid Z_1]$ .
- So, we posit a (possibly misspecified) model  $p_{Z_2|Z_1}^*(\cdot \mid z_1; \xi)$  and compute CEF

$$h_2^*(Z_1, \beta, \xi) = \int m(z_1, u, \beta) p_{Z_2|Z_1}^*(u \mid z_1; \xi) du.$$

- Estimation: fit  $\hat{\psi}_n$  by MLE for  $\pi(\cdot, \psi)$ ; fit  $\hat{\xi}_n^*$  by likelihood for the posited model; solve

$$\sum_{i=1}^n \left[ \frac{R_i}{\pi(Z_{1i}, \hat{\psi}_n)} m(Z_i, \beta) - \frac{R_i - \pi(Z_{1i}, \hat{\psi}_n)}{\pi(Z_{1i}, \hat{\psi}_n)} h_2^*(Z_{1i}, \beta, \hat{\xi}_n^*) \right] = 0.$$

- This estimation strategy is relatively easy to implement.

## AIPWCC estimation which is ignorant of $\xi$

- While, using posit model, we can think of the limiting value  $h_2^*(Z_{1i}, \beta, \xi^*)$ .
- Even if  $h_2^*(Z_{1i}, \beta, \xi^*) \neq h_2^0(Z_{1i}, \beta)$ , it is a function of  $Z_{1i}$ , which satisfies the condition that can construct the IF.

$$- \left\{ \frac{R_i - \pi(Z_{1i}, \psi_0)}{\pi(Z_{1i}, \psi_0)} \right\} h_2^*(Z_{1i}, \beta_0, \xi^*) \in \Lambda_2$$

- Therefore, the estimating equation for IF is

$$\sum_{i=1}^n \left[ \frac{R_i m(Z_i, \beta)}{\pi(Z_{1i}, \hat{\psi}_n)} - \left\{ \frac{R_i - \pi(Z_{1i}, \hat{\psi}_n)}{\pi(Z_{1i}, \hat{\psi}_n)} \right\} h_2^*(Z_{1i}, \beta_0, \xi^*) \right] = 0$$

- The only difference is to estimate  $\xi_n^*$  or not (AIPWCC requires integration).

# Asymptotic equivalence between adaptive estimation and AIPWCC

## Theorem 10.3

Let  $\hat{\beta}_n$  be the estimator obtained by the adaptive estimation, and let  $\hat{\beta}_n^*$  be the AIPWCC estimator . Under suitable regularity conditions,

$$n^{1/2} \left( \hat{\beta}_n - \hat{\beta}_n^* \right) \xrightarrow{P} 0.$$

Hence, the adaptive estimator is asymptotically equivalent to the AIPWCC estimator.

- Thus, we think misspecification of  $\xi$  and its estimation (=adaptive estimation) as a kind of negligible problem. (BUT no efficiency !)

## Double robustness (DR)

- Under regularity, the adaptive AIPWCC estimator is **consistent** if

*either*  $\pi(Z_1, \psi)$  is correctly specified **or**  $p_{Z_2|Z_1}^*(\cdot | Z_1; \xi)$  is correct.

- The estimation function holds, then  $\beta$  is consistent and asymptotically normal.

$$\mathbb{E} \left[ \frac{R m(Z, \beta_0)}{\pi(Z_1, \psi^*)} - \left\{ \frac{R - \pi(Z_1, \psi^*)}{\pi(Z_1, \psi^*)} \right\} h_2^*(Z_1, \beta_0, \xi^*) \right] = 0,$$

where  $\hat{\psi}_n \xrightarrow{P} \psi^*$  and  $\hat{\xi}_n^* \xrightarrow{P} \xi^*$ .

## **10.5 Improving Efficiency when Coarsening is Nonmonotone**

---



## Generalization idea: operators $L$ and $M$ (you'll need this)

### Definition 4 and 5

- Define two linear operators  $\mathcal{L} : \mathcal{H}^F \rightarrow \mathcal{H}$  and  $\mathcal{M} : \mathcal{H}^F \rightarrow \mathcal{H}^F$

$$\mathcal{L}\{h^F\} = \mathbb{E}[h^F(Z) \mid C, G_C(Z)], \quad \mathcal{M}\{h^F\} = \mathbb{E}[\mathcal{L}\{h^F\} \mid Z].$$

where  $h^F$  is an arbitrary element in  $\mathcal{H}^F$ .

$$L\{h_F(\cdot)\} = \sum_{r=1}^{\infty} I(C = r) \mathbb{E}\{h_F(Z) \mid G_r(Z)\},$$

and

$$M\{h_F(\cdot)\} = \sum_{r=1}^{\infty} \pi\{r, G_r(Z)\} \mathbb{E}\{h_F(Z) \mid G_r(Z)\}.$$

## How to derive the projection : $\Pi \left[ \frac{1(C = \infty)h_F(Z)}{\pi(\infty, Z)} \middle| \Lambda_2 \right]$

### Theorem 10.6

- Inverse operator  $\mathcal{M}^{-1}$  exists and is uniquely defined.
- Moreover, the projection is given by

$$\Pi \left[ \frac{1(C = \infty)h_F(Z)}{\pi(\infty, Z)} \middle| \Lambda_2 \right] = \frac{\infty(C = \infty)h_F(Z)}{\pi(\infty, Z)} - \mathcal{L}[\mathcal{M}^{-1}\{h_F(\cdot)\}],$$

- By **Definition 1**,  $\mathcal{L}[\mathcal{M}^{-1}\{\varphi^F(Z)\}] = \mathcal{J}\{\varphi^F(Z)\}$ .
- Therefore, DR linear space is written by  $\mathcal{J}(\Lambda^{F\perp}) = \mathcal{L}[\mathcal{M}^{-1}\{\Lambda^{F\perp}\}]$ .

## Proof sketch for the uniqueness of $\mathcal{M}^{-1}$

- $I - \mathcal{M}$  is a contraction mapping

$$\Leftrightarrow \forall h^F \in \mathcal{H}^F, 0 \leq \exists L < 1, \|(I - \mathcal{M})h^F\| \leq L\|h^F\|$$

- Existence of  $\mathcal{M}^{-1} : \sum_{k=0}^{\infty} (I - \mathcal{M})^k$  (Completeness of linear operator space)

$$\because \sum_{k=0}^n (I - \mathcal{M})^k \circ \mathcal{M} = I - (I - \mathcal{M})^{n+1} \rightarrow I \quad (\text{as } n \rightarrow \infty)$$

- Uniqueness of  $\mathcal{M}^{-1} : \mathcal{M}^{-1} = \mathcal{M}^{-1}(\mathcal{M} \circ \mathcal{M}'^{-1}) = (\mathcal{M}^{-1} \circ \mathcal{M})\mathcal{M}'^{-1} = \mathcal{M}'^{-1}$

## Constructing Strategy

- we only focus on the operators  $\mathcal{L}, \mathcal{M}^{-1}$  to estimate  $\beta$ .

$$\sum_{i=1}^n \mathcal{L}_i [\mathcal{M}_i^{-1} \{ m(Z_i, \beta) \}] = 0.$$

- $\mathcal{L}$  : posit model  $\xi$  for conditional probability of  $Z$  given  $G_r(Z) \rightarrow \mathcal{L}(\cdot, \xi)$
- $\mathcal{M}^{-1}$  :  $\psi$  for coarsening probability, posit model  $\xi$  (same above)  $\rightarrow \mathcal{M}^{-1}(\cdot, \psi, \xi)$
- To construct  $\beta$ , after estimating  $\psi$  for  $\hat{\psi}_n$ , and  $\xi$  for  $\hat{\xi}_n^*$ , solve the equation

$$\sum_{i=1}^n \mathcal{L}_i [\mathcal{M}_i^{-1} \{ m(Z_i, \beta), \hat{\psi}_n, \hat{\xi}_n^* \}, \hat{\xi}_n^*] = 0$$

## Equivalence of $\mathcal{L} \circ \mathcal{M}^{-1}$ and AIPWCC

### Theorem 10.7 - (i)

- Let  $d^F(Z, \beta, \psi, \xi) = \mathcal{M}^{-1}\{m(Z, \beta), \psi, \xi\}$ , then, the residual can separate

$$\begin{aligned}\mathcal{L}[\mathcal{M}^{-1}\{m(Z, \beta), \psi, \xi\}, \xi] &= \mathcal{L}\{d_F(Z, \beta, \psi, \xi), \xi\} \\ &= \underbrace{\frac{I(C = \infty) m(Z, \beta)}{\pi(\infty, Z, \psi)}}_{\text{IPWCC space}} + \underbrace{L_2^*\{C, G_C(Z), \beta, \psi, \xi\}}_{\text{Augmentation space}}.\end{aligned}$$

- This means the solution to this estimating equation is an AIPWCC estimator

$$\sum_{i=1}^n \frac{I(C_i = \infty) m(Z_i, \beta)}{\varpi(\infty, Z_i, \hat{\psi}_n)} + L_2^*\{C_i, G_{C_i}(Z_i), \beta, \hat{\psi}_n, \hat{\xi}_n^*\} = 0.$$

## The part of augmentation space

### Theorem 10.7 - (ii)

- The element of the augmentation space is written by

$$\begin{aligned} L_2^*\{C, G_C(Z), \beta, \psi, \xi\} = \\ - \frac{I(C = \infty)}{\pi(\infty, Z, \psi)} \left( \sum_{r \neq \infty} \pi(r, G_r(Z), \psi) \mathbb{E}[d_F(Z, \beta, \psi, \xi) \mid G_r(Z), \xi] \right) \\ + \sum_{r \neq \infty} I(C = r) \mathbb{E}[d_F(Z, \beta, \psi, \xi) \mid G_r(Z), \xi] \end{aligned}$$

- Let  $L_{2r}\{G_r(Z)\} = -\mathbb{E}[d_F(Z, \beta, \psi, \xi) \mid G_r(Z), \xi]$ , we can obtain the same representation as a typical element of  $\Lambda_2$ . (see (7.37))

## Construct $\mathcal{M}^{-1}$ and $\mathcal{L}$

- $\mathcal{M}^{-1}$  is represented by infinite series, so be constructed by successive approximation.

$$d_{(j)}^F(Z, \beta, \psi, \xi) = \hat{\mathcal{M}}^{-1}\{m(Z, \beta), \psi, \xi\}$$

- By construction,  $L_2^*\{C, G_C(Z), \beta, \psi, \xi\} \in \Lambda_2$  in any cases.
- This means that when  $\psi_0$  (coarsening probability model) is correctly specified,

$$\frac{I(C = \infty) m(Z, \beta_0)}{\varpi(\infty, Z, \psi_0)} + L_{2(j)}^*\{C, G_C(Z), \beta_0, \psi_0, \xi^*\} \in \Lambda^\perp$$

- Hence, we can obtain an AIPWCC estimator by solving

$$\sum_{i=1}^n \frac{I(C_i = \infty) m(Z_i, \beta)}{\varpi(\infty, Z_i, \hat{\psi}_n)} + L_{2(j)}^*\{C_i, G_{C_i}(Z_i), \beta, \hat{\psi}_n, \hat{\xi}_n^*\} = 0.$$

- The estimator is RAL (i.e. consistent and asymptotically normal)

- In addition to correct specification of  $\psi$ , the posit model  $p_Z^*(z, \xi)$  is correctly specified, its influence function is **efficient**. (double-robustness)
- Also, this AIPWCC estimator is consistent and asymptotically normal even if  $\psi$  is misspecified. (Actually, we do not have to consider this pattern **thanks to CAR**)
- Recall this two probabilistic model
  - posit model  $p_Z^*(z, \xi)$ ; estimate  $\hat{\xi}_n^* \rightarrow \xi^* (\neq \xi_0)$  (if misspecified)
  - coarsening model  $\pi\{r, G_r(Z), \psi\}$ ; estimate  $\hat{\psi}_n^* \rightarrow \psi^* (\neq \psi_0)$  (if misspecified)



## Theorem 10.8

- This theorem establishes the **double robustness** property.
- Specifically:

$$\mathbb{E}_{\xi_0, \psi_0} [L(M^{-1}\{m(Z, \beta_0), \psi_0, \xi^*\}, \xi^*)] = 0,$$

and

$$\mathbb{E}_{\xi_0, \psi_0} [L(M^{-1}\{m(Z, \beta_0), \psi^*, \xi_0\}, \xi_0)] = 0.$$

- Hence, the estimating equation remains unbiased if either the coarsening model ( $\psi$ ) or the marginal model of  $Z$  ( $\xi$ ) is correctly specified.

## **11. Locally Efficient Estimators for Coarsened-Data Semiparametric Models**

---

## Roadmap for Chapter 11

- For a full-data estimating function  $m(Z, \beta) \in \Lambda^{F\perp}$ , applying  $\mathcal{J}$ , we obtain an observed-data estimating function.

$$\sum_{i=1}^n \left[ \frac{I(C_i = \infty)m(Z_i, \beta)}{\varpi(\infty, Z_i, \hat{\psi}_n)} - \Pi \left( \frac{I(C_i = \infty)m(Z_i, \beta)}{\varpi(\infty, Z_i)} \mid \Lambda_2 \right) \right] = 0$$

- For computation, we need to estimate  $\psi$  in the former,  $\xi$  (posit model) in the latter.
- We propose a locally efficient estimator, which achieves the semiparametric efficiency bound if the posited model  $p_Z^*(z, \xi)$  is correct but still be consistent and asymptotically normal and RAL even if misspecified.
- Recall the essential framework : Efficient score  $\rightarrow$  Efficient IF (proportional relationship)

## **11.1 The Observed-Data Efficient Score**

---

## Efficient score: Representation 1 (likelihood-based) : theoretical

- Basiccally, the efficient score is given by

$$S_{\text{eff}}\{C, G_C(Z)\} = S_{\beta}\{C, G_C(Z)\} - \Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda],$$

where  $S_{\beta} = \mathbb{E}[S_{\beta}^F(Z) \mid C, G_C(Z)]$ , and  $\Lambda = \Lambda_{\psi} \oplus \Lambda_{\eta}$  ( $\Lambda_{\psi} \perp \Lambda_{\eta}$ ).

$$\begin{aligned}\Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda] &= \Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda_{\psi}] + \Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda_{\eta}] \\ &= \Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda_{\eta}] \quad (\because S_{\beta}\{C, G_C(Z)\} \perp \Lambda_2)\end{aligned}$$

- Recall  $\Lambda_{\eta} = \{\mathbb{E}[\alpha^F(Z) \mid C, G_C(Z)] : \alpha^F(Z) \in \Lambda^F\}$ , there exists  $\alpha_{\text{eff}}^F(Z) \in \Lambda^F$ ,

$$\Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda_{\eta}] = \mathbb{E}[\alpha_{\text{eff}}^F(Z) \mid C, G_C(Z)]$$

$$\begin{aligned}\therefore S_{\text{eff}}\{C, G_C(Z)\} &= S_{\beta}\{C, G_C(Z)\} - \Pi[S_{\beta}\{C, G_C(Z)\} \mid \Lambda_{\eta}] \\ &= E[\{S_{\beta}^F(Z) - \alpha_{\text{eff}}^F(Z)\} \mid C, G_C(Z)]\end{aligned}$$

## Efficient score: Representation 2 (AIPWCC-based) : practical

- Since the full-data efficient score  $B^F(Z) \in \Lambda^{F\perp}$ , we can apply  $\mathcal{J}$ .

$$S_{\text{eff}}\{C, G_C(Z)\} = \mathcal{J}\{B_F^{\text{eff}}(Z)\} = I(C = \infty) \frac{B_F^{\text{eff}}(Z)}{\pi(\infty, Z)} - \Pi \left[ I(C = \infty) \frac{B_F^{\text{eff}}(Z)}{\pi(\infty, Z)} \middle| \Lambda_2 \right].$$

- Clearly, this element lives in DR linear space  $\mathcal{J}(\Lambda^{F\perp})$
- Now, we obtain two equivalent representations about the efficient score

$$\mathbb{E}[S_{\beta}^F(Z) - \alpha_{\text{eff}}^F(Z) \mid C, G_C(Z)] = \frac{I(C = \infty) B_{\text{eff}}^F(Z)}{\varpi(\infty, Z)} - \Pi \left[ \frac{I(C = \infty) B_{\text{eff}}^F(Z)}{\varpi(\infty, Z)} \middle| \Lambda_2 \right]$$

- For computation, this AIPWCC estimator is far more practical but still be difficult to construct.  $\rightarrow$  Efficiency problem boils down to find  $B_{\text{eff}}^F(Z)$ .

## How to derive $B_{\text{eff}}^F(Z)$

### Theorem 11.1

- $B_{\text{eff}}^F(Z) \in \Lambda^{F\perp}$  is the unique solution to

$$\Pi\left[\mathcal{M}^{-1}\{B^F(Z)\} \mid \Lambda^{F\perp}\right] = S_{\text{eff}}^F(Z),$$

- Therefore, solving this equation, we obtain  $B_{\text{eff}}^F(Z)$ , then we can construct the efficient score (Note : This is revisited)

$$S_{\text{eff}}\{C, G_C(Z)\} = \frac{I(C = \infty) B_{\text{eff}}^F(Z)}{\varpi(\infty, Z)} - \Pi\left[\frac{I(C = \infty) B_{\text{eff}}^F(Z)}{\varpi(\infty, Z)} \mid \Lambda_2\right],$$

## Proof sketch of Lemma 11.1 : successive approximation

- To construct  $B_{\text{eff}}^F(Z)$ , define  $D_{\text{eff}}^F(Z) = \mathcal{M}^{-1}\{B_{\text{eff}}^F(Z)\}$ .
- Then  $D_{\text{eff}}^F(Z)$  is the solution to

$$S_{\text{eff}}(Z) = \Pi[h^F(Z) \mid \Lambda^{F\perp}] + \Pi[\mathcal{M}\{h^F(Z)\} \mid \Lambda^F] = (I - \mathcal{Q})\{h^F(Z)\}$$

$$\text{where } \mathcal{Q}\{D_{\text{eff}}^F(Z)\} = \Pi[(I - \mathcal{M})\{D_{\text{eff}}^F(Z)\} \mid \Lambda^F]$$

- $\mathcal{Q}$  is a contraction mapping, so  $(I - \mathcal{Q})^{-1}$  uniquely exists.
- Finally, we can successively approximate  $D_{\text{eff}}^F(Z) = (\sum_{i=0}^{\infty} \mathcal{Q}^i) S_{\text{eff}}^F(Z)$



## Theorem 11.2

- Suppose the coarsening mechanism is **monotone**.
- For any  $h^F(Z) \in \mathcal{H}^F$ , the inverse operator  $M^{-1}$  has the closed form

$$M^{-1}\{h^F(Z)\} = \frac{h^F(Z)}{\varpi(\infty, Z)} - \sum_{r \neq \infty} \frac{\lambda_r\{G_r(Z)\}}{K_r\{G_r(Z)\}} \mathbb{E}[h^F(Z) \mid G_r(Z)],$$

where

$$K_r(Z) = \Pr(C \geq r + 1 \mid Z), \quad \lambda_r\{G_r(Z)\} = \Pr(C = r \mid C \geq r, Z).$$

- Hence,  $M^{-1}$  can be computed explicitly in terms of the hazard and survival functions of the coarsening distribution.

## Lemma 11.2

- Let  $\{D^{(k)}\}$  be defined recursively by

$$D^{(k+1)}(Z) = \Pi\left((I - M)D^{(k)}(Z) \mid \mathcal{H}^F\right) + S_{\text{eff}}^F(Z),$$

with  $D^{(0)}$  arbitrary.

- Then

$$D^{(k)} \longrightarrow D_{\text{eff}}^F \quad \text{in } \mathcal{H}^F \text{ norm,}$$

and

$$B^{(k)}(Z) = \Pi(MD^{(k)}(Z) \mid \Lambda^{F\perp}) \longrightarrow B_{\text{eff}}^F(Z).$$

- Thus, the successive approximation scheme converges to the efficient element  $B_{\text{eff}}^F(Z)$ .

## **11.2 Strategy for Obtaining Improved Estimators**

---

- Consider finding a full-data estimating function  $m(Z, \beta)$ .
- Firstly, by MLE (or likelihood methods), we estimate  $\hat{\psi}_n, \hat{\xi}_n^*$ .
- Secondly, we calculate full-data efficient score  $S_F^{\text{eff}}(Z)$ .
- By successive approximation,  $\hat{B}_{\text{eff}}^F(Z) = m(Z, \hat{\psi}_n, \hat{\xi}_n^*)$ .
- Finally, we can estimate observed-data efficient score using

$$S_{\text{eff}}\{C, G_C(Z)\} = \frac{I(C = \infty) B_{\text{eff}}^F(Z)}{\varpi(\infty, Z)} - \Pi \left[ \frac{I(C = \infty) B_{\text{eff}}^F(Z)}{\varpi(\infty, Z)} \mid \Lambda_2 \right]$$

- Now, we focus on the projection  $\Pi \left[ \frac{I(C = \infty) m(Z, \beta)}{\varpi(\infty, Z)} \mid \Lambda_2 \right]$ .  
 $\rightarrow$  In some special coarsening mechanism, this projection is simply written down.
- In actual implementation, we solve this adaptive estimating equation.

$$\begin{aligned}
& \sum_{i=1}^n \left( \frac{I(C_i = \infty) m(Z_i, \beta, \hat{\psi}_n, \hat{\xi}_n^*)}{\varpi(\infty, Z_i, \hat{\psi}_n)} \right. \\
& \quad - \frac{I(C_i = \infty)}{\varpi(\infty, Z_i, \hat{\psi}_n)} \left[ \sum_{r \neq \infty} \varpi\{r, G_r(Z_i), \hat{\psi}_n\} \mathbb{E} \left\{ D^{(j)}(Z, \beta, \hat{\psi}_n, \hat{\xi}_n^*) \mid G_r(Z_i), \hat{\xi}_n^* \right\} \right] \\
& \quad \left. + \sum_{r \neq \infty} I(C_i = r) \mathbb{E} \left\{ D^{(j)}(Z, \beta, \hat{\psi}_n, \hat{\xi}_n^*) \mid G_r(Z_i), \hat{\xi}_n^* \right\} \right) = 0
\end{aligned}$$

## locally efficient estimator for restricted moment model with monotone coarsening

- Full data:  $Z = (Y, X)$ , Model:  $Y = \mu(X, \beta) + \varepsilon$  with  $\mathbb{E}(\varepsilon \mid X) = 0$ .
- Full-data efficient score:  $S_{\text{eff}}^F(Z) = D^\top(X)V^{-1}(X)\varepsilon$ ,  
 $D = \partial\mu/\partial\beta^\top$ ,  $V(X) = \text{Var}(Y|X)$ .
- Seek  $B_F^{\text{eff}}(Z) = A(X)\varepsilon \in \mathcal{F}^\perp$  solving

$$\Pi[M^{-1}\{A(X)\varepsilon\} \mid \mathcal{F}^\perp] = S_{\text{eff}}^F(Z).$$

- Leads to an integral equation for  $A(X)$ ; exact solution rare; use numerical/approximate  $A_{\text{imp}}$ .

## Approximate locally efficient estimator (practice)

- Compute a numerically feasible  $A_{\text{imp}}(X, \hat{\beta}, \hat{\psi}, \hat{\xi}^*)$ .
- Plug into the AIPWCC efficient-score form:

$$\sum_{i=1}^n \left[ I(C_i = \infty) \frac{A_{\text{imp}}(X_i, \beta, \hat{\psi}, \hat{\xi}^*) \{Y_i - \mu(X_i, \beta)\}}{\pi(\infty, Y_i, X_i; \hat{\psi})} + L_2\{C_i, G_{C_i}(Y_i, X_i), \beta, \hat{\psi}, \hat{\xi}^*\} \right] = 0.$$

- Here  $L_2$  is the projection term (closed form under monotone coarsening).

### Robustness note

The  $A_{\text{imp}}$  route targets local efficiency but may be numerically heavy; when it's too hard, fall back to the DR AIPWCC from Ch. 10.

## **11.3 Concluding Thoughts**

---



- Adaptive estimation : Specified coarsening model  $\rightarrow$  using posit model  $\xi \rightarrow$  fit  $\xi_n^*$
- Inverse weighted method balances between simplicity of implementation and relative efficiency.
- AIPWCC estimator is more improved one in double robustness and gaining considerable efficiency at the cost of numerical implementation. (successive approximation)
- Locally efficient estimator aims to find optimal full-data estimating function  $B_{\text{eff}}^F(Z)$  as well as optimal augmentation. (difficult to implement)

## References

---

- Tsiatis, A. (2006). *Semiparametric Theory and Missing Data*. Springer (Chs. 10–11).
- A.W. van der Vaart (1998). *Asymptotic Statistics*. Cambridge University Press.