

Generic machine learning inference on heterogeneous treatment effects in randomized experiments, with an application to immunization in India

Victor Chernozhukov et al. (2024), Econometrica.

Naoki Eguchi

2025.7.9 ミクロ計量経済学

Faculty of Medicine, Kyoto University

Introduction

- When conducting RCT (Randomized Controlled Trials), researchers and policy makers are often curious about not only ATE but also HTE (Heterogeneous Treatment Effects).
- In such cases, ML methods are good for estimating

Proposed estimator

-

Main identification results and estimation strategies

BLP (Best Linear Predictor)

- Firstly, we obtain the estimator $S(Z)$ for CATE $s_0(Z)$ by some ML method using the auxiliary samples \mathcal{A} .
- BLP is defined as the projection of $s_0(Z)$ on the linear span of 1 and $S(Z)$ in $L^2(P)$.

$$\text{BLP}[s_0(Z)|S(Z)] = \arg \min_{f(Z) \in \text{Span}(1, S(Z))} \mathbb{E}[(s_0(Z) - f(Z))^2]$$

- This equals to the solution of $\arg \min_{b_1, b_2} \mathbb{E}[(s_0(Z) - b_1 - b_2 S(Z))^2]$.

$$\rightarrow \beta_1 = \mathbb{E}[s_0(Z)], \beta_2 = \frac{\text{Cov}(s_0(Z), S(Z))}{\text{Var}(S(Z))}$$

First strategy : Weighted Residual BLP

- Consider the regression model with the moment condition as follows.

$$Y = \alpha' X_1 + \beta_1(D - p(Z)) + \beta_2(D - p(Z))(S(Z) - \mathbb{E}[S(Z)]) + \epsilon, \mathbb{E}[w(Z)\epsilon X] = 0$$

$$\text{where } w(Z) = \frac{1}{p(Z)(1 - p(Z))}, X = (X_1, X_2),$$

$$X_1 = (1, B(Z)), X_2 = (D - p(Z), (D - p(Z))(S(Z) - \mathbb{E}[S(Z)]))).$$

Theorem 1

- Consider $z \mapsto S(z)$ and $z \mapsto B(z)$ as fixed maps. (known function)
- Assume that Y and X have finite second moments, $\mathbb{E}[X X']$ is full rank, and $\text{Var}(S(Z)) > 0$.
- Then, $(\beta_1, \beta_2)' = \arg \min_{b_1, b_2} \mathbb{E}[(s_0(Z) - b_1 - b_2 S(Z))^2]$ (Identified)

Second strategy : Horvitz-Thompson BLP

Theorem 2

- Consider $z \mapsto S(z)$ and $z \mapsto B(z)$ as fixed maps. (known function)
- Assume that Y has finite second moments, $\mathbb{E}[\tilde{X}\tilde{X}']$ is finite and full rank, and $\text{Var}(S(Z)) > 0$.
- Then, $(\beta_1, \beta_2)' = \arg \min_{b_1, b_2} \mathbb{E}[(s_0(Z) - b_1 - b_2 S(Z))^2]$ (Identified)

GATES (sorted Group Average Treatment Effects)

- Firstly, we build the groups by the estimated value $S(Z)$ of $s_0(Z)$.

$$G_k = \{S(Z) \in I_k\}, k = 1, \dots, K, I_k = [l_{k-1}, l_k) : \text{disjoint}$$

- The estimand "GATES" is defined as $\mathbb{E}[s_0(Z)|G_k]$ for $k = 1, \dots, K$.

Two strategies : Weighted Residual and Horvitz-Thompson GATES

- Consider the regression model with the moment condition as follows.

$$Y = \alpha' X_1 + \sum_{k=1}^K \gamma_k (D - p(Z)) \mathbf{1}_{G_k} + \nu, \mathbb{E}[w(Z) \nu W] = 0$$

$$\text{where } W = (X_1, W_2')', W_2 = \{(D - p(Z)) \mathbf{1}_{G_k}\}_{k=1}^K.$$

Theorem 3

- Consider $z \mapsto S(z)$ and $z \mapsto B(z)$ as fixed maps. (known function)
- Assume that Y has finite second moments and both $\mathbb{E}[WW']$ and $\mathbb{E}[\tilde{W}\tilde{W}']$ are finite and full rank.
- Then, $\gamma = \{\gamma_k\}_{k=1}^K$ defined in two different ways are equivalent and identified.

$$\gamma_k = \mathbb{E}[s_0(Z)|G_k]$$

CLAN (Classification Analysis)

- When BLP and GATES reveal substantial heterogeneity, it is interesting to know **the properties of the subpopulations** that are most and least affected.
 - We focus on the "least affected group" G_1 and "most affected group" G_K .
- Let $g(Y, Z)$ be a vector of characteristics of an observational unit. The estimands are

$$\delta_1 = \mathbb{E}[g(Y, Z)|G_1] \quad \text{and} \quad \delta_K = \mathbb{E}[g(Y, Z)|G_K]$$

- These parameters are identified with no assumption because they are just average of observed variables.
- We compare δ_1 and δ_K to detect (single out) the covariates which causes the heterogeneity.
 - We can extend the comparison of not only averages but also variances or distributions.

”Variational” estimation and inference methods

Applicatoion

Final marks

References

- Victor Chernozhukov, Mert Demirer, Esther Duflo, and Iván Fernández-Val (2024), *Generic machine learning inference on heterogeneous treatment effects in randomized experiments, with an application to immunization in India*, Econometrica.
- Kosuke Imai and Michael Lingzhi Li (2025), *Statistical Inference for Heterogeneous Treatment Effects Discovered by Generic Machine Learning in Randomized Experiments*, Journal of Business and Economic Statistics.
- 末石直也 (2024), データ駆動型回帰分析, 日本評論社.