

Recommended Clinical Targets for Glycemic Control for Adults with Type 1 or Type 2 Diabetes

Targets	Type 1/Type 2 Diabetes	Older/High Risk Individuals with Type 1/Type 2 Diabetes
Time in range (TIR)		
% of readings	› 70%	› 50%
Time per day	› 16 hr, 48 min	› 12 hr
Glucose target range	70–180 mg/dL	70–180 mg/dL
Time below range (TBR)		
LEVEL 1 Hypoglycemia		
% of readings	‹ 4%	‹ 1%
Time per day	‹ 1 hr	‹ 15 min
Glucose range	‹ 70 mg/dL	‹ 70 mg/dL
LEVEL 2 Hypoglycemia		
% of readings	‹ 1%	0%
Time per day	‹ 15 min	0 min
Glucose range	‹ 54 mg/dL	‹ 54 mg/dL
Time above range (TAR)		
LEVEL 1 Hyperglycemia		
% of readings	‹ 25%	‹ 50%
Time per day	‹ 6 hr	‹ 12 hrs
Glucose range	› 180 mg/dL	› 180 mg/dL
LEVEL 2 Hyperglycemia		
% of readings	‹ 5%	‹ 10%
Time per day	‹ 1 hr, 12 min	‹ 2 hrs, 24 min
Glucose range	› 250 mg/dL	› 250 mg/dL

Adapted from Battelino T, et al. Clinical targets for continuous glucose monitoring data interpretation: recommendations from the international consensus on time in range. *Diabetes Care*. 2019;42:1593-1603.