

SAD OR HOPELESS & SUICIDE PLAN

YRBS 2007 . N=13,666

PRIMARY

Does the manifestation of a clinical depressive mood significantly alter the probability of a student formulating an active suicide plan?

EXTENDED

Are depression and suicidal tendencies related to gender or grade level

VARIABLES

Independent :
X1 : Sad_Or_Hopeless
X2 : Sex

Dependent :
Y : Suicide_Score

Covariate :Grade

METHOD

Two-Way ANOVA

DEPRESSIVE MOOD / SUICIDE





H₀: no significant different. H₁:have a significant different.
F=1591, P<0.001 → **Reject H₀ ✓**

GENDER / SUICUDE

H₀: no significant different. H₁:have a significant different.
F=5.666, P=0.0173 → **Reject H₀ ✓**

GENDER&DEPRESSIVE MOOD / SUICIDE

H₀: no significant different. H₁: have a significant different.
F=0.516, P=0.4725 → **Fail to Reject H₀ ✗**



CONCLUSION

A persistent depressive mood is the single most dominant predictor of adolescent suicide planning, Furthermore, while females are more susceptible to falling into depression than males, the interaction between sex and depression is completely non-significant. This establishes the "Universal Injury Principle": once depression sets in, its psychological escalation into an active suicide plan operates with identical, destructive velocity across both genders.