

a
NWF-C clast
M / Ms
B (mT)
fresh interior
weathered rind
b
PKS-B flow (1155 ka)
B (mT)
c
SD
PSD
filled = fresh
open = weathered
MD →
(Mr/Ms < 0.05)
Mr / Ms
Hcr / Hc
PKS-B (flow)
PKS-H (flow)
PKS-I (corestone)
NWF-C (clast)
PKQ-A 2A (clast)
PKQ-A 5A (clast)
PKQ-A 5C (clast)
d
NWF-C, low temperature
Verwey (~120 K)
M / Mmax
T (K)
ZFC LT-SIRM
FC LT-SIRM
RT-SIRM cooling
RT-SIRM warming
e
PKS-H (in situ flow)
peak Bc ≈ 11 mT
f
PKS-I (corestone)
peak Bc ≤ 5 mT
g
PKS-B flow / NWF-C clast
peak Bc ≈ 17 mT
PKS-B, fresh interior
Fresh interior Bu (mT)
h
peak Bc ≈ 10 mT
i
peak Bc ≈ 15 mT
j
peak Bc ≈ 12 mT
NWF-C, weathered rind
Weathered rind Bu (mT)
Bc (mT)
ρ (normalised)
k
500 µm
titanomagnetite spots
BSE, fresh basalt LS-H18
l
FeO
TiO2
Cr2O3
V2O5
MnO
MgO
Al2O3
SiO2
CaO
Na2O
K2O
Titanomagnetite (Oxide 1)
Cr-titanomagnetite (Oxide 2)
Titanomagnetite (Oxide 3)
Titanomagnetite (Oxide 5)
Fe–Mn oxide (secondary) (Oxide 4)
Olivine (Fo53) (Pyx 1)
Olivine (Fo78) (Pyx 2)
Augite (Ukn 1)
Plagioclase (An59) (Plag 1)
Plagioclase (An62) (Plag 2)
Oxide (wt%, EDS, normalised)