

a
NWF-A · NWFA-4C
in situ flow, 518 ka · normal
ChRM D = 2°, I = 23°, MAD = 0.5°
E, Up
N
NRM
×10⁻⁵ A m²
J₀ = 3.3e−5 A m²
J / J₀
T (°C)
b
PKS-J · PKSJ-2E
in situ flow, 601 ka · reversed
ChRM D = 173°, I = -32°, MAD = 1.5°
J₀ = 8.9e−6 A m²
c
PKS-B · PKSB-1F
in situ flow, 1155 ka · reversed
ChRM D = 192°, I = -9°, MAD = 2.6°
J₀ = 7.2e−5 A m²
d
WBE-A · WBEA-2B
basal flow, 1397 ka · reversed
ChRM D = 170°, I = -30°, MAD = 2.8°
J₀ = 1.5e−5 A m²
e
PKS-H, core HA1 · PKSH-1I
upper horizon · reversed VGP
ChRM D = 142°, I = −34°, MAD = 5.1°
×10⁻⁶ A m²
J₀ = 6.7e−6 A m²
f
PKS-H, core HA2 · PKSH-2H
upper horizon · 64° from HA1
ChRM D = 192°, I = 11°, MAD = 3.3°
J₀ = 9.5e−6 A m²
g
PKS-H, core HB1 · PKSH-3B
lower flow · reversed VGP
ChRM D = 207°, I = 7°, MAD = 4.3°
J₀ = 9.2e−6 A m²
h
NWF-C · NWFC-1C
diamicton clast (conglomerate test)
ChRM D = 281°, I = 7°, MAD = 1.0°
J₀ = 5.5e−5 A m²
i
ChRM inclination vs age
Brunhes (N)
PKS-K
NWF-A (a)
PKS-I
PKS-J (b)
B–M 773 ka
PKS-HA (e, f)
impact ~790 ka
PKS-HB (g)
NWF-B
Jaramillo
PKS-B (c)
Cobb Mtn
WBE-A (d)
Matuyama (R)
⁴⁰Ar/³⁹Ar age (ka)
ChRM inclination (°, geographic)
horizontal (N–E)
vertical (N–Up)
step rejected in fitting
LT fit (arrow = demagnetization direction)
MT fit (arrow = demagnetization direction)
HT (ChRM) fit (arrow = demagnetization direction)
site mean, normal (i)
reversed (i)
disturbed PKS-HA cores (i)
clast locality (i)
GAD inclination (±29°)
NRM (not used in fits)
after LN₂ (low-T demag.)